JUN 1 1 1996

John S. Thornton
Bayer Corporation
P.O. Box 4913
8400 Hawthorn Road
Kansas City, MO 64120-0013

Dear Mr. Thornton:

Subject: Add sidedress application to potato
Admire(TM) 2 Flowable

EDA Posistration No. 3125-422

EPA Registration No. 3125-422 Your resubmission of May 5, 1996

The amendment referred to above, submitted in connection with registration under FIFRA, is acceptable provided that you:

Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours

DHE

Dennis H. Edwards, Jr. Product Manager (19) Insecticide Rodenticide Branch Registration Division (7505C)

Enclosure

U.S. LABEL

Reason to Issue: To add sidedress applications to potatoes. 05/09/96 Draft: To remove 80-day PHI restriction.

Date of Draft: 05/09/96 (Pre-Reg) (C) Supersedes Pre-Reg Draft: 02/28/96

Admire™ 2 Flowable

Systemic Insecticide

For control of certain insects infesting vegetables.

ACTIVE INGREDIENT:

!midacloprid, 1-[(6-Chloro-3-pyridinyl)methyl] -N-nitro-2-imidazolidinimine INERT INGREDIENTS 100.0%

Contains 2 pounds of imidacloprid per gallon EPA File Symbol No. 3125-422

Net Contents: Gallons

STOP - Read The Label Before Use KEEP OUT OF THE REACH OF CHILDREN

AMENDMENT To Currently Registered Labeling

RECOMMENDED APPLICATIONS			
CROP	PEST	RATE PER APPLICATION	REMARKS
Potato	Aphids Colorado potato beetle Flea beetles Potato leafhopper Psyllids	Soil Application 0.9 to 1.3 fl oz per 1000 row feet (see rate conversion chart for rates per acre)	Apply specified dosage in one of the following methods: 1. As a narrow band directly below the seed row in a bedding operation 7 or fewer days before planting. 2. As an in-furrow spray during planting. For best results, direct spray on the seed pieces or seed potatoes in the furrow. 3. As a sidedress on both sides of the potato row of as a spray at ground cracking directly over the row during hilling: In either case. ADMIRE must be covered immediately with 3 or, more inches of soils Make applications no closer than 80 days before harvest. The higher rate is recommended when planting into muck soils, for seed potato production, or when extended length of control is needed. Do not apply more than 0.31-lb active ingredient of ADMIRE per acre per season as a soil application regardless of formulation.

NOTE: Regardless of formulation or type of application (soil or foliar), do not apply more than à total 0.31 lb active ingredient of ADMIRE or PROVADO per acre per season.

Text being added is highlighted by redlining.

Text being deleted is highlighted by strikeout.

Bayer Corporation Crop Protection Products Kansas City, MO 64120-0013

