



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

3125-427

Date of Issuance:

APR 28 1993

NOTICE OF PESTICIDE:
 X Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: Until
Reregistration

Name of Pesticide Product:

Guthion 3 Flowable
Insecticide

Name and Address of Registrant (include ZIP Code):

Miles Inc.
P.O. Box 4913, 4800 Hawthorne Road
Kansas City, MO 64120-0013

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. The registrant continues to be subject to any obligation regarding data submission or compensation under FIFRA Sections 3(c)(1)(F), 4(H) and 3(c)(2)(B) that pertains to EPA Reg. No. 3125-338.
3. Add the phrase "EPA Registration No. 3125-427" to your label before you release the product for shipment.

Signature of Approving Official:

DENNIS H. EDWARDS

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division

Date:

4. Submit five copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

DENNIS H. EDWARDS

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

Reason to Issue: To delete sugarcane use
in Louisiana.

Date of Draft: 04/12/93(C)
Supersedes Draft Dated: 02/19/92

RESTRICTED USE PESTICIDE

Due to Acute Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

GUTHION 3 Flowable INSECTICIDE

FOR EFFECTIVE ECONOMICAL INSECT CONTROL

ACTIVE INGREDIENT:

O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate 30%

INERT INGREDIENTS: 70%
100%

Contains 3 lb O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate per gallon.

EPA Reg. No. 3125-XXX
EPA Est. 3125-MO-1

Net Contents: ___ Gallons

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

DANGER



POISON

(See rear panel for Statements of Practical Treatments and Other Precautionary Statements)
PROTECT FROM HEAT. SHAKE WELL BEFORE USING.

PELIGRO


PRECAUTION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 28 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 3125-427

MILES



GUTHION 3 Flowable Insecticide

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER



POISON

Poisonous if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Do not breathe fumes or spray mist. Spray operator should work to windward to stay out of drift or mist.

When loading spray tank or handling the concentrate at any other time, wear protective clothing, natural rubber gloves, and goggles. When handling the concentrate, wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U. S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11.

Do not contaminate feed or foodstuffs. Keep out of reach of children and domestic animals. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking.

HANDLING OF CONTAMINATED ARTICLES: Clothing and rags receiving undiluted GUTHION 3 Flowable as a result of leaks, spills, or mishaps should be removed immediately and not reused. These contaminated articles should be buried in the same manner as the empty container; see container disposal section for directions. (Contaminated articles which should be discarded would include: hats, gloves, aprons, coats, boots, etc. not made of rubber or coated with rubber or other similar materials). Clothing receiving spray mist or droplets from the mixed, diluted GUTHION 3 Flowable should be removed and decontaminated before reuse. This clothing can be decontaminated by machine washing separately from other items with soap or detergent and bleach in hot water twice before reuse. Protective clothing and equipment should be washed down with detergent or soap and bleach in water. Wash water from the cleaning of protective clothing or equipment should not be allowed to run off or otherwise enter water supplies.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

STATEMENTS OF PRACTICAL TREATMENT

In case of poisoning call a physician or Poison Control Center immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz or 15 ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Professional medical assistance should be secured immediately. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

NOTE TO PHYSICIAN:

ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and wildlife.

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water

by cleaning of equipment or disposal of wastes. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PROTECTIVE CLOTHING AND WORK SAFETY STATEMENTS

If handled indoors provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See First Aid section of the label.

For questions regarding further treatment, or information about the product, call the National Pesticide Telecommunications Network at 800-858-7378.

HANDLE THE CONCENTRATE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT: Wear a protective suit of one or two pieces that covers all parts of the body except the head, hands, and feet. Wear chemical resistant gloves, chemical resistant apron, and chemical resistant shoes, shoe coverings, or boots. Wear goggles or a face shield and a pesticide respirator approved by the National Institute for Occupational Safety and Health under the provision of 30 CFR Part 11.

If handling the concentrate with a closed system, long sleeved shirt and long-legged pants may be substituted for the protective suit and the respirator requirement is waived.

WEAR THE FOLLOWING PROTECTIVE CLOTHING DURING APPLICATION, EQUIPMENT REPAIR, EQUIPMENT CLEANING, DURING REENTRY TO TREATED AREAS, AND DISPOSAL OF THE PESTICIDE: Wear a protective suit of one or two pieces that covers all parts of the body except the head, hands, and feet. Wear chemical resistant gloves and chemical resistant shoes, shoe coverings, or boots. Wear a chemical resistant hat during airblast application.

During application from an enclosed tractor cab or airplane cockpit, or other suitable vehicle in which the windows are rolled up, long sleeved shirt and long pants may be worn in place of the above protective clothing. Chemical resistant gloves must be available in the cab or cockpit and must be worn while exiting. This clothing is inadequate to protect you during equipment repair, equipment cleaning, re-entry, or during pesticide disposal.

IMPORTANT! BEFORE REMOVING GLOVES, WASH THEM WITH SOAP AND WATER. ALWAYS WASH HANDS, FACE, AND ARMS WITH SOAP AND WATER BEFORE SMOKING, DRINKING, EATING OR TOILETING.

After work, take off all clothing and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirators must be cleaned and filters replaced according to instructions included with the respirators. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with GUTHION azinphos-methyl must be destroyed according to state and local regulations. **HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.**


During aerial application, human flaggers are prohibited unless in totally enclosed vehicle.

RE-ENTRY STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas 24 hours after application unless protective clothing is worn, as described in the product labeling. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or an area about to be treated with this product. Oral warnings should include those statements found under the label heading

"Precautionary Statements" and Re-entry Statements" included on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER, area treated with GUTHION 3 Flowable on (date of application). Do not enter treated areas for 24 hours after application unless protective clothing is worn, as described in the product labeling. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes. If swallowed, vomiting should be induced. (See Precautionary Statements on label).



ROTATIONAL CROP STATEMENT

Do not plant root crops other than those with registered azinphos-methyl uses in azinphos-methyl treated soil sooner than 6 months after the last application. Do not

plant any other crop other than those with registered azinphos-methyl uses in treated soil sooner than 30 days after last applications.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using GUTHION 3 Flowable insecticide.


CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. MILES OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MILES AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Mixing: GUTHION 3 Flowable insecticide mixes readily with water and is suitable for use in all power-operated ground sprayers and aircraft sprayers. Fill the spray tank 1/4 to 1/3 full with clean water and add GUTHION 3 Flowable at the recommended rate. Mix thoroughly and add water to fill spray tank. GUTHION 3 Flowable may also be applied in a 1:1 ratio with water as an ultra-low volume spray with either ground or aircraft equipment that has been adapted and calibrated for ultra-low volume spraying as described below under "Recommended Applications" for those crops specified.


Do not apply this product through any type of irrigation system.

DOSAGE: Use specified dosage of GUTHION 3 Flowable in the amount of water necessary to give complete coverage of foliage. The type of equipment used will determine the concentration required.

SPRAYING: Work to windward. Protect sprayer operators from drift or mist. When low volumes of spray are applied, complete coverage and thorough application are essential for most effective results. Schedule applications in accordance with local conditions. Consult your State Agricultural Extension Service or Experiment Station for specific use information in your area.




RECOMMENDED APPLICATIONS			
CROP	INSECT	PINTS GUTHION 3 FLOWABLE	REMARKS
FIELD CROPS Alfalfa	Alfalfa weevil	2/3 to 2	Apply specified dosage per acre using a minimum of 10 gallons of water per acre when applying with ground equipment, and a minimum of 5 gallons per acre for aerial applications for alfalfa weevils and at least 1 gallon per acre for the other insects listed. It may be necessary to use 20 to 25 gallons of water per acre on heavy growth for control of alfalfa weevil or Egyptian alfalfa weevil with ground equipment.
	Leafhoppers Mites	2/3 to 1-1/3	
	Egyptian alfalfa weevil	1 to 2	Apply twice per cutting at the 2/3 pint rate at intervals of 10 to 11 days. Do not apply more than twice per cutting at the 2/3 pint rate. Apply only once per cutting at rates above 2/3 pint per acre. Do not apply within 14 days of harvest at the rates of 2/3 to 1 pint per acre or within 16 days at the rate of 1-1/3 pints per acre. Do not apply rates above 1-1/3 pints per acre within 21 days of harvest.
	Alfalfa plant bug Fleahopper Grasshoppers Lygus bugs Spittlebug	1-1/3 to 2	
Cotton (Conventional or Low Volume Spray)	Boll weevil	1/3 to 2/3	Apply specified dosage per acre by air ground equipment in at least 1 gallon of water. For early season control of bollworm and pink bollworm in states west of Rocky Mountains use 1-1/3 to 2 pints per acre and for mid-to-late season applications, use 2 to 2-2/3 pints per acre. Repeat as necessary. Do not apply within 1 day of picking at rates up to 1-1/3 pints per acre or within 7 days of picking at rates above 1-1/3 pints per acre. Do not graze treated fields.
	Brown cotton leafworm Cotton fleahopper Cotton leafworm Lygus bugs Thrips	2/3	
	Rapid plant bug Tarnished plant bug	2/3 to 1-1/3	
	Stink bug	1-1/3	
	Bollworm Pink bollworm	East of Rocky Mountains 1-1/3	
		West of Rocky Mountains 1-1/3 to 2-2/3	



RECOMMENDED APPLICATIONS

CROP	INSECT	PINTS GUTHION 3 FLOWABLE	REMARKS
<p>Sugarcane (Conventional Spray)</p> <p>(Florida and Texas only)</p> <p>Not for use in Louisiana</p>	<p>Sugarcane borer</p>	<p align="center">2</p>	<p>Apply specified dosage per acre by air application using a minimum of 2 gallons of water per acre. For best results applications should be made at approximately cane top level and not more than 10 feet above cane top level. Do not apply more than 5 times per season. Allow at least 30 days between the last application and harvest. Bagasse from sugarcane treated with GUTHION may be used for feed of cattle, goats, and sheep. Do not graze treated fields. Consult your State Agricultural Extension Service or Experiment Station for specific use information.</p>
<p>Sugarcane¹ (Ultra Low Volume Spray)</p> <p>(Florida and Texas only)</p> <p>Not for use in Louisiana</p>	<p>Sugarcane borer</p>	<p align="center">2</p>	<p>Apply specified dosage per acre. GUTHION 3 Flowable may be used in a 1:1 ration with water in any aerial spray equipment that has been adapted and calibrated for ultra-low volume spraying. Planes must be equipped with accepted low volume devices that will produce droplets within the range of 30 to 100 microns in size. For best results applications should be made at approximately cane top level and not more than 1 foot above cane top level. Do not apply more than 5 times per season. Allow at least 30 days between the last application and harvest. Bagasse from sugarcane treated with GUTHION may be used for feed of cattle, sheep and goats. Do not graze treated fields. Consult your State Agricultural Extension Service or Experiment Station for specific use information.</p>

¹ NOTE: This formulation, when used ultra-low volume, may cause spotting of automobile finishes if prolonged exposure is permitted. Do not spray directly over automobiles. If accidental exposure does occur, automobiles should be washed immediately.



RECOMMENDED APPLICATIONS

CROP	PEST	GUTHION 3F		REMARKS															
		PINTS/ ACRE	PINTS/ 100 GAL																
FRUIT Apples Pears (Except California)	Aphids* Apple maggot Codling moth Eyespotted bud moth Leafhoppers* Plum curculio Redbanded leafroller San Jose Scale* Tarnished plant bug* Green fruitworm* Fruit tree leafroller European apple sawfly Forbes scale Mealybug Putnam scale Oblique-banded leafroller*	2	1/2	Apply specified dosage as a full coverage spray. Up to 8 pints GUTHION 3F may be applied per acre per crop season. Allow at least 7 days between applications and 7 days between last application and harvest. * In some areas, these species may have developed resistance to organophosphate insecticides. GUTHION 3F used alone may not provide satisfactory control in those areas. Consult your local agricultural advisor or Cooperative Extension Service for recommendations.															
					Cherries (East of Rocky Mountains)	Eye-spotted bud moth Forbes scale Fruit flies Fruit tree leafroller Lesser peach tree borer Plum curculio San Jose scale* Cherry leaf miner Mineola moth	2	2/3	Apply specified dosage per acre as a full coverage spray. Up to four applications may be applied per acre per crop season. Allow at least 14 days between applications and 21 days between the last application and harvest.										
										Cherries (West of Rocky Mountains)	Leafhoppers Fruit flies Fruit tree leafroller Lesser peach tree borer Cherry leaf miner	PINTS GUTHION 3F <hr/> 1-1/3 (7 fl oz per 100 gal)		Apply specified dosage per acre as a full coverage spray. Up to two applications may be applied per acre per crop season. Allow at least 14 days between applications and 7 days between last application and harvest.					
															*In some areas, these species may have developed resistance to organophosphate insecticides. GUTHION 3 Flowable used alone may not provide satisfactory control in those areas. Consult your local agricultural advisor or Cooperative Extension Service for recommendations.				
															VEGETABLES Potatoes	Colorado potato beetle	1		Apply specified dosage per acre in sufficient water for complete coverage. A total of 3 applications may be made per crop season regardless of rate, formulation or method of application. Allow at least 7 days between applications and 7 days between last application and harvest. NOTE: Resistance of Colorado potato beetle has occurred in some areas. Consult your local Extension Service or Miles Sales Representative for details.
																Banded cucumber beetle Leafminers	1 to 1-1/3		
																European corn borer Flea beetle Leafhoppers Spittlebugs Tarnished plant bug	1-1/3 to 2		
																Tuberworm	1-1/2 to 2		

RECOMMENDED APPLICATIONS

CROP	INSECT	PINTS GUTHION 3 FLOWABLE	REMARKS
ORNAMENTALS Christmas Trees	Scale spp. Sawfly spp. European pine shoot moth Eastern pine shoot borer Nantucket pine tip moth	1 - 2	Apply specified dosage per acre by air or ground equipment in sufficient water to give complete coverage but not less than 1 gallon per acre. Time applications to coincide with susceptible pest development.
Southern Pine Seed Orchards	Coneworm Seedworm	See Remarks	Use 4 pints per 100 gallons of water (0.2% dilution) as a full coverage spray.** Use 2 pints per 10 gallons of water (1% dilution) for low volume sprayers. Apply first application within 30 days following conelet closure, followed by 3 to 5 applications (at least 30 days apart). Apply approximately 5 to 10 gallons of the 0.2% dilution with high volume sprayers or approximately 1 to 2 gallons of the 1% dilution with low volume sprayers per tree. Thorough coverage of cones is necessary for maximum control.

*In some areas, these species may have developed resistance to organophosphate insecticides. GUTHION 3 Flowable used alone may not provide satisfactory control in those areas. Consult your local agricultural advisor or Cooperative Extension Service for recommendations.

** This concentration is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per acre with concentrate sprayers, increase the concentration of GUTHION 3 Flowable in the spray mixture in order to apply amount of GUTHION 3 Flowable per acre equivalent to a full coverage spray. In Southern Pine Seed Orchards where conditions dictate an air application, apply dosage per acre equivalent to a full coverage ground spray in not less than one gallon of water per acre.

Ornamentals Nursery Plants	Aphids <i>Cercococcus</i> scale Euonymus scale Juniper scale Lace bugs Leafhoppers Mites Olive scale Oystershell scale <i>Pulvinaria</i> scale Thrips	1 to 1-1/3	Apply specified dosage per 100 gallons of water (1-1/3 teaspoonfuls per gallon). Spray all foliage surfaces including the underside of leaves for complete coverage. For control of black pine leaf, brown soft, European elm, and Putnam scales, use 2/3 tablespoonful per gallon. Repeat as necessary.
	Brown soft scale Putnam scale	2-2/3	
	European elm scale Black pine leaf scale	2 to 2-2/3	
	Cone midge Cone moth	5-1/3 to 10-2/3	
	European pine shoot moth Nantucket pine tip moth	1 to 2	

Injury to Hawthorn or American linden may occur under some conditions. Do not allow children or pets on treated area until material has been washed into soil and treated area is dry.

RESTRICTIONS

Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues.

Do not graze livestock in tted orchards or groves for 21 days after treatment. Do not treat food crops grown in the greenhouse.

STORAGE AND DISPOSAL

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides above. In spill or leak incidents, keep unauthorized people away. You may contact the Miles Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Miles Kansas City Emergency Response Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

Miles, Inc.
Crop Protection Products
Box 4913, Kansas City, MO 64120-0013

