

BAYLETON 25 WP

436-8697.YLD

4/11/22

3125-436

U. S. LABEL

Base Pre-Reg (8697)

Reason to Issue: To revise directions for use on bermudagrass decline and take all patch.

Date of Draft: 01/03/96(T)

BAYLETON® 25

Wettable Powder

Nursery and Greenhouse Systemic Fungicide

For control of certain diseases on flowers, foliage plants, shrubs, turf and shade trees in commercial nurseries, garden centers, greenhouses and sod farms.

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone 25%

INERT INGREDIENTS 75%
100%


U.S. Patent No. 3,912,752

EPA Reg. No. 3125-436

EPA Est. 3125-MO-1

Net Weight: _____ Pounds

AMENDMENT TO PREVIOUSLY
REGISTERED LABELING



Unregistered label text is highlighted in redlining
Text to be deleted is indicated by ~~strikeout~~

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL			
DISEASE	OZ OF BAYLETON 25 WP PER 1,000 SQUARE FEET *		REMARKS
Bermudagrass decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>) Take all patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>) (Except California)	Preventive 2 - 4	Curative 4	Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone. PREVENTIVE RATE: Begin applications 2 to 4 weeks prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21- to 28-day intervals. For take all patch, applications in both spring and fall may be necessary. CURATIVE RATE: To control existing infections, apply 4 oz at 14 to 21 day intervals for the initial 4 or 2 treatments followed by 2 to 4 oz at 21- to 28-day intervals. Cultural control practices such as aerification, topdressing, reseeding, and fertilization should must be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.

Bayer Corporation
 Specialty Products
 Box 4913, Kansas City, Mo. 64120-0013