

PM 22  
BAYLETON 25 WSP

438-8542.BLD

3125-438  
U. S. LABEL

Base Pre-Reg (8542)

Reason to Issue: To revise name for bermudagrass decline and take all patch.

Date of Draft: 06/07/95(Pre-Reg)(T)

# BAYLETON® 25

## Nursery and Greenhouse Systemic Fungicide in Water Soluble Packets

For control of certain diseases on flowers, foliage plants, shrubs, turf and shade trees in commercial nurseries, garden centers, greenhouses and sod farms.

### ACTIVE INGREDIENT:


1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone	25%
INERT INGREDIENTS	75%
	100%

Keep water soluble packets in this container and store in a cool place but not below freezing (32°F).

U. S. Patent No. 3,912,752  
EPA Reg. No. 3125-438  
EPA Est. 3125-MO-1

Sell in Original Packages Only  
Contains Four 11 Ounce Water Soluble Packets  
Total Net Contents: 2.75 Pounds

### AMENDMENT TO PREVIOUSLY REGISTERED LABELING



Bayer

Unregistered label text is highlighted in redlining.  
Text to be deleted is indicated by strikeout.

**BEST COPY AVAILABLE**

1 of 2

2 of 2

RECOMMENDED APPLICATIONS - TURF GRASS DISEASE CONTROL		
DISEASE	OZ OF BAYLETON 25 NURSERY AND GREENHOUSE PER 1,000 SQUARE FEET*	REMARKS
<p><i>Gaeumannomyces graminis</i> var. <i>graminis</i></p> <p>(Such as Bermudagrass decline, take-all patch on centipede grass)</p> <p>Bermudagrass decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)</p> <p>Take-all patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)</p> <p>(Except California)</p>	<p>Preventive    Curative</p> <p>2                4</p>	<p>Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone.</p> <p><b>PREVENTIVE RATE:</b> Begin applications 2 to 4 weeks prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21 to 28-day intervals.</p> <p><b>CURATIVE RATE:</b> To control existing infections, apply 4 oz at 14- to 21-day intervals for the initial 1 or 2 treatments followed by 2 oz at 21- to 28-day intervals. Cultural control practices such as aerification, top-dressing, reseeding, and fertilization must be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.</p>

2