

# BAYLETON 25 WSP

438-8698.YLD

pm 22

3125-438

1/2

Base Pre-Reg (8698)

U. S. LABEL

Reason to Issue: To revise directions for use on bermudagrass decline and take all patch.

Date of Draft: 01/03/96(T)

## BAYLETON® 25

### Nursery and Greenhouse

### Systemic Fungicide in Water Soluble Packets

For control of certain diseases on flowers, foliage plants, shrubs, turf and shade trees in commercial nurseries, garden centers, greenhouses and sod farms.

**ACTIVE INGREDIENT:**


1-(4-Chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone .....	25%
INERT INGREDIENTS .....	75%
	100%

Keep water soluble packets in this container and store in a cool place but not below freezing (32°F).

U. S. Patent No. 3,912,752  
EPA Reg. No. 3125-438  
EPA Est. 3125-MO-1

Sell in Original Packages Only  
Contains Four 11 Ounce Water Soluble Packets  
Total Net Contents: 2.75 Pounds

#### AMENDMENT TO PREVIOUSLY REGISTERED LABELING



Unregistered label text is highlighted in red in original.  
Text to be deleted is indicated by ~~strikeout~~

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL			
DISEASE	OZ OF BAYLETON 25 NURSERY AND GREENHOUSE PER 1,000 SQUARE FEET*		REMARKS
Bermudagrass decline <i>(Gaeumannomyces graminis var. graminis)</i>  Take all patch <i>(Gaeumannomyces graminis var. avenae)</i>  (Except California)	Preventive  2 - 4	Curative  4	<p>Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone.</p> <p><b>PREVENTIVE RATE:</b> Begin applications <del>2 to 4 weeks</del> prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21- to 28-day intervals. For take all patch, applications in both spring and fall may be necessary.</p> <p><b>CURATIVE RATE:</b> To control existing infections, apply 4 oz at <del>14 to 21 day intervals</del> for the initial <del>1 or 2</del> treatments followed by 2 to 4 oz at 21- to 28-day intervals. Cultural control practices such as aerification, topdressing, reseeding, and fertilization should <b>must</b> be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.</p>

Bayer Corporation  
 Specialty Products  
 Box 4913, Kansas City, Mo. 64120-0013

