3125-439 lesse read instructions on reverse before completing form Form Approved. OMB No. 2070-0060. Approval expires 05-31-98 OPP Identifier Number Registration **United States Environmental Protection Agency** Amendment 252431 Washington, DC 20460 Other Application for Pesticide - Section I 2. EPA Product Manager 3. Proposed Classification 1. Company/Product Number 3125-439 Dennis H. Edwards, Jr. X None Restricted 4. Company/Product (Name) Merit 75 WSP Insecticide 19 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling Bayer Corporation 8400 Hawthorn Road, P.O. Box 4913 EPA Reg. No. Kansas City, MO 64120 Check if this is a new address Product Name Section - II Amendment - Explain below. Final printed labels in response to Agency letter dated Resubmission in response to Agency letter dated \_ "Me Too" Application. Notification - Explain below. Other - Explain below. NOTIFICATION Explanation: Use additional page(s) if necessary. (For section I and Section II.) APR 2 5 1997 Adding a Pest Section - III 1. Material This Product Will Be Packaged In: Child-Rasistant Packaging Unit Packaging Water Soluble Packaging 2. Type of Container Metal Yes' Yes Yes Plastic Glass Paper No. per If "Yes" No. per \* Certification must Unit Packaging wgt. Package wgt Other (Specify) be submitted 3. Location of Net Contents Information 4. Size(s) Retail Container 5. Location of Label Directions Lahal Container On Labeling accompanying product Lithograph Paper glued Stenciled Other 6. Manner in Which Label is Affixed to Product Section - IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Telephone No. (include Area Code) (816) 242-2457 Charles W. Boyd Senior Registrations Scientist 6. Date Application Certification bevieag?... l certify that the statements I have made on this form and an accommendation of the statement of a cknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment of a cknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment of a cknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment of a cknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment of a cknowledge that any knowledge that an I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. (Stamped) 2. Signature Director, Product Registrations and Regulatory Affairs 5. Date April 4, 1997 John S. Thornton

Merit<sup>3</sup> 75 WSP Insecticide EPA Reg. No. 3125-439 April 3, 1997 Page 1 of 1


Section II - Explanation

Bayer Corporation is submitting a Notification of the addition of a pest to the label of MERIT® 75 WSP Insecticide, EPA Reg. No. 3125-439.

Per PR Notice 95-2, (paragraph II(A)(2)) we are notifying the Agency that **Cutworm** is being added to the pest listing under the recommendations for use on Turfgrasses. We are also changing the pest table to allow the use of a rate range on most of the pests labelled. This rate range was already approved for Annual bluegrass weevil and Billbugs.

This notification is consistent with the provision of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Enclosed is one copy of the product label, draft dated 3/17/97.



U.S. LABEL

Base Reg. (8931)

Reason to Issue: To add cutworm suppression and to allow a rate range on turfgrass pests.

Date of Draft: 03/17/97 (S) Supercedes Draft Dated: 03/23/95

## Merit® 75 WSP

#### Insecticide

For foliar and systemic insect control in turfgrass, landscape ornamentals and interior plantscapes.

#### **ACTIVE INGREDIENT:**

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]- N-nitro-2-imidazolidinimine	75.0%
INERT INGREDIENTS	25.0%
	100.0%

Keep water soluble packets in this container and store in a cool dry place but not below freezing (32°F). Do Not Remove Packets From Container Except For Immediate Use.

EPA Reg. No. 3125-439

Net Contents: \_\_\_ Pounds

# STOP - Read The Label Before Use KEEP OUT OF REACH OF CHILDREN NOTIFICATION

#### CAUTION

APR 2 5 1997

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with

soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

#### STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. If syrup of ipecac is available, administer 1 tablespoonful (15 mL) of syrup of ipecac followed by 1 to 2 glasses of water. If vomiting does not occur within 20 minutes, repeat the dose once. Do not induce

vomiting or give anything by mouth to an unconscious person. If on skin: Wash thoroughly with soap and water. Get medical attention if irritation occurs. If in eyes: Hold eyelids open and flush with plenty of water.

To Physician: No specific antidote savailable. Treat the patient symptomatically.

#### **ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical demonstrates the properties and characteristics

Merit 75 WSF

associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable,

particularly where the water table is shallow, may result in groundwater contamination.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire DIRECTIONS FOR USE, GENERAL INFORMATION, AND CONDITIONS OF SALE before using MERIT 75 WSP Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY RESULT FROM THE OCCURRENCE MAY EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Do not formulate this product into other end-use products.

#### **APPLICATION TO TURFGRASS:**

MERIT 75 WSP Insecticide can be used for the control of soil inhabiting pests of turfgrass, such as Northern & Southern masked chaters, Cyclocephala borealis, C. immaculata, and/or C. lurida; Asiatic garden beetle, Maladera castanea; European chafer, Rhizotrogus majalis; May or June beetle. Phyllophaga spp.; Japanese beetle, Popillia japonica; Oriental beetle, Anomala orientalis; Billbugs, Spherophorus spp.; Annual bluegrass weevil, Hyperodes spp.; Black turfgrass ataenius, Ataenius spretulus and Aphodius spp.; and mole crickets, Scapteriscus spp. MERIT 75 WSP Insecticide can also be used for the suppression of cutworms in turigrass areas. MERIT 75 WSP insecticide can be used as directed on turigrass in sites such as home lawns, business and office complexes, shopping complexes, multifamily residential completes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields. MERIT 75 WSP insecticide can not be used on commercial sod farms.

The active ingredient in MERIT 75 WSP Insecticide has sufficient residual activity so that applications can be made

preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Applications should not be made when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Applications cannot exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

#### Application Equipment for Use on Turfgrass

Apply MERIT 75 WSP Insecticide in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

#### APPLICATION TO ORNAMENTALS:

MERIT 75 WSP Insecticide is for use on ornamentals in commercial and residential landscapes and interior plantscapes. MERIT 75 WSP Insecticide is a systemic product and will be translocated upward into the plant system. To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution will enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. In some cases, this translocation delay can take up to 60 days. For this reason, applications should be made prior to anticipated pest infestation to achieve optimum levels of control.

For outdoor ornamentals, broadcast applications cannot exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

NOTE: Not for use in commercial greenhouses, nurseries, on sod farms or on grass grown for seed. For use on plants



intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

#### Application Equipment for Ornamental Uses

MERIT 75 WSP Insecticide mixes readily with water and may be used in many types of application equipment. Mix product with the required amount of water and apply as desired dependent upon the selected use pattern.

When making foliar applications on hard to wet foliage such as holly, pine, or ivy, the addition of a spreader/ sticker is recommended. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed, as would be used in a dilute application.

MIXING: Within each foil pouch is a clear inner packet containing MERIT 75 WSP Insecticide. The clear inner packet is water soluble. Do not allow packets to become wet prior to adding to the spray tank. Do not handle the clear inner packets with wet hands or wet gloves. Rough handling may cause breakage. Reseal outer carton to protect remaining packets.

To prepare the spray mixture, remove the outer foil pouch and drop the required number of unopened clear water soluble packets, as determined under "Recommended Applications", into the spray tank while filling with water to the desired level. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within a few minutes from the time they are added to the water. Cooler water temperatures increase the time needed for the inner packet to dissolve completely.

CAUTION: Do not use MERIT WSP packets in a tank-mix with products that contain Boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic which is not soluble in water or solvents such as diesel oils, kerosene, gasoline or alcohol. Do not attempt to use the WSP packets directly in diesel oils or summer spray type oils as in ULV or LV uses. PVA packets are water soluble not oil soluble.

MERIT 75 WSP Insecticide has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides.

Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

Do not apply through any irrigation system.

RECOMMENDED APPLICATIONS				
CROP	PEST	DOSAGE MERIT 75 WSP	REMARKS	
Turfgræsses	Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Cutworm (suppression) European chafer Japanese beetle Northern masked chafer Oriental beetle Phyllophaga spp. Southern masked chafer	1.6 oz (1 Packet) per 8,250 to 11,000 sq ft	For optimum control of grubs, billbugs and annual bluegrass weevil, make application prior to egg hatch of the target pest.  Be sure to read "APPLICATION EQUIPMENT" Section of this label.	
	Mole crickets	1.6 oz (1 packet) per 8,250 sq ft	For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, MERIT application should be accompanied by a curative insecticide. Follow label instructions for other insecticides when tank-mixing.	

Consult your local State Agricultural Experiment Station, or State Extension Turf Specialists for more specific information regarding fiming of application.

NOTE: For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 8.6 oz (0.4 ib of active ingredient) per acre per year. Avoid mowing turf or lawners a until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

RECOMMENDED APPLICATIONS FOR USE ON or IN ORNAMENTALS  For use only in and around industrial and commercial buildings and residential areas				
CROP	PEST	DOSAGE MERIT 75 WSP	REMARKS	
Trees Shrubs Evergreens Flowers Foliage plants Ground covers	Adelgids Aphids Elm leaf beetle Lacebugs Leafhoppers Leatminers Meaiybugs Sawfly larvae Thrips White Flies	1.6 oz (1 packet) per 300 gal of water	Foliar Applications: Start treatments prior to establishment of high pest populations and reapply on an as needed basis.	
	White grub larvae (such as Japanese beetle larvae, Chafers, Phyllophaga spp. Aslatic garden beetle, Oriental beetle)	1.6 oz (1 packet) per 8,250 to 11,000 sq ft	Broadcast Applications: Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of water per 1000 sq ft. For optimum control, irrigate thoroughly to incorporate MERIT 75 WSP insecticide into the upper soil profile.  Refer to REMARKS section for use directions specific for FLOWERS and GROUND COVERS concerning additional use directions.	
Trees Shrubs Flowers & Ground Covers	Adelgids Aphids Lacebugs Leafminers Mealybugs Scale insects Thrips Whiteflies Pine Tip moth larvae Elm Leaf beetle Sawfly larvae Japanese beetle Leafhopper White grub larvae	For Trees:  1.6 oz (1 packet) per 24 to 48 inches of cumulative trunk diameter	Soil Injection: GRID SYSTEM: Holes should be spaced on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. CIRCLE SYSTEM: Apply in holes eventy spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.  Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.  Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.	
		For Shrubs:  1.6 oz (1 packet) per 24 to 48 ft of cumulative shrub height	Soil Injection: Apply to individual plants using dosage indicated.  Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.  Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.	
		For Flowers & Ground Covers:  1.6 oz (1 packet) per 8,250 to 11,000 sq ft	Apply as a broadcast treatment and incorporate into the soll before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application.	



#### RESTRICTIONS

Do not graze treated areas or use clippings from treated areas for feed or forage. Avoid runoff or puddling of irrigation water following application. Keep children and pets off treated area until dry. Avoid application of MERIT 75 WSP Insecticide to areas which are water logged or saturated,

which will not allow penetration into the root zone of the plant. Do not apply more than 8.6 oz (0.4 lb of active ingredient) per acre per year. Do not plant any food crop within one year of a treatment with MERIT 75 WSP Insecticide.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill, by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides above. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response telephone number is 800-414-0244 or contact Chemtrec at 800-424-9300.

