3125-452 4-14-2000 orm Approved. OMB No. 2070-0060. Approval expires 05-31-98 Mease read bistruction on reverse before completing for **OPP Identifier Number United States** Registration **Environmental Protection Agency Amendment** 257730 Washington, DC 20460 Other Application for Pesticide - Section I 2. EPA Product Manager 1. Company/Product Number 3. Proposed Classification 3125-452 Dani Daniel Restricted None 4. Company/Product Name PM# 0 4 Merit 1G Greenhouse and Nursery Insecticide 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) **Bayer Corporation** (b)(i), my product is similar or identical in composition and labeling 8400 Hawthorn Road, P.O. Box 4913 EPA Reg. No. Kansas City, MO 64120-0013 Check. If this is Is a new address Product Name Section - II Final printed labels in response to Amendment - Explain below. Agency letter dated NOTIFICATION "Me Too" Application. Resubmission in response to Agency letter dated APR 1 2000 Notification - Explain below. Other - Explain below. Explanation: Use additional page(s) if necessary. (For Section I and Section II.) See Attached Section - III 1. Material This Product Will Be Packaged in: Child-Resistant Packaging **Unit Packaging** Water Soluble Packaging 2. Type of Container Yes* Yes Yes Metal No No Plastic Glass If "Yes" No Per If "Yes No. Per Paper Unit Packaging wgt.. Container Package wgt. Containe ertification must Other (Specify) be submitted 3. Location of Net Contents Information 4. Size(s) Retail Container 5. Location of Label Directions On Label ☐ Label Container On Labeling accompanying product 6. Manner in Which Label is Affixed to Product Lithograph Other Paper Glued Stenciled Section - IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Name Telephone No. (Include Area Code) Charles W. Boyd Senior Registrations Scientist (816) 242-2457 Certification 6. Date Application Received I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. (Stamped)',, I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. 2. Signature 3. Title Manager, Federal Regulatory Affairs 5. Date Julie M. Spagnoli April 11, 2000

2/6

Page 1 of 1

Merit® 1 G Greenhouse and Nursery Insecticide EPA Reg. No. 3125-452 April 11, 2000

Section II - Explanation

Bayer Corporation is submitting a **Notification** to add leafhoppers to the Merit® 1 G Greenhouse and **Nursery Insecticide label** for ornamental plant application.

Enclosed is one copy of the product label, draft number M - 8944b.

This notification is consistent with the provision of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

POTIFICATION

APR 1 4 2000

Merit® 1 G

Greenhouse and Nursery Insecticide

For systemic insect control in ornamental crops.

ACTIVE INGREDIENT:

Imidacloprid,1-[(6-Chloro-3-pyridinyl)methyl]-	
-N-nitro-2-imidazolidinimin	.1.0%
OTHER INGREDIENTS	. <u>99.0%</u>
1	00.0%

EPA Reg. No. 3125-452

Net Contents: ___ Ounces

STOP - Read the label before use. Keep out of reach of children.

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS** AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- · Water-proof gloves
- · Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

· When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker

Protection Standard (WPS) for agricultural pesticides [40] CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENTS OF PRACTICAL TREATMENT

If in eyes: Hold eyelids open and flush with plenty of water. Call a physician if irritation persists. If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger, or, if available, administer 1 tablespoonful. (15 mL) of syrup of ipecac followed by 1 to 2 glasses of water. If vomiting does not occur within 20 minutes, repeat the dose once. Do not induce vomiting or give anything by mouth to an unconscious person. If on skin: Wash thoroughly with soap and water. Get medical attention if irritation occurs.

To Physician: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water

Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. ' b'nly' protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consulted as the agency responsible for pesticide regulation.



AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- · Waterproof gloves
- · Shoes plus socks

IMPORTANT: Read these entire DIRECTIONS FOR USE, GENERAL INFORMATION, AND CONDITIONS OF SALE before using MERIT 1 G Greenhouse and Nursery Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

GENERAL INFORMATION

MERIT 1 G Greenhouse and Nursery Insecticide is for insect control on ornamental plants in nurseries and greenhouses. MERIT 1 G Greenhouse and Nursery Insecticide is a systemic product and will be translocated upward within the plant. To assure optimum effectiveness, the product must be placed by incorporation where the roots of the plant can

absorb the active ingredient. Imigate moderately but thoroughly after application, allowing no leaching and runout from containers for at least three irrigations or 10 days whichever is longer.

INCORPORATION

Incorporation of MERIT 1 G Greenhouse and Nursery Insecticide can be achieved by cultivation, irrigation, rainfall, mechanical placement or by using mechanical soil or media mixing equipment.

WOODY PERENNIALS

Onset of protection is slower than in herbaceous species. A delay of 2 or more weeks should be expected. Longer delays may be expected with larger plants. Application should therefore be made in advance of expected insect activity.

BARK MEDIA

Media with 30% or more bark content when treated with MERIT 1 G Greenhouse and Nursery Insecticide may confer a shorter period of protection.

RESTRICTIONS

Do not apply to soils which are water-logged, or saturated. This condition will not allow the penetration of water into the root zone of the plants.

Do not over-irrigate or allow excessive runoff to occur following an application of MERIT 1 G. Do not apply to bedding plants intended to be used as food crops.

For outdoor ornamentals grown in beds, applications of MERIT 1 G cannot exceed a total of 40 lb per acre per year.

On plants with a production cycle of less than one year, application is not to exceed a frequency of more than once each 16 weeks for a particular plant. On stock plants and woody crops with a production cycle of greater than one year, application may not exceed once a year.



Beetles Jacebugs Jacebugs Jeafhoppers Leafminers Jeafminers Jeafmi			DOSAGE						
PEST VISE Size Portscut yd Bulk Rate Topical Rate To			Container	Bulk App	lication	Topical Ap	plication	REMARKS	
Note Septies Species	DECT		size				C		
Species Spec			 -					5 . 5	
Complex Comp	_								
Containerized 1 Gallon 200 to 160 1.5 to 1.0 1.5 to 1.0 2.73 to 1 2.7 to 4.1 1.0 to 1.3 1.0 to 0.8 1.0	Aphids	Species	3.5 to 5.0		3.9 to 3.0	1/4 to 1/3	1.0 to 1.4		
Suppression Containerized 1 Gallon 200 to 160 10.0 to 160 11.0 to 10.0 10.0 to 160 12.0 200 to 160 12.0 20.0 to 160			5.5 to 7.0		1.8 to 1.2	1/3 to 1/2	1.4 to 2.0	_	
Definite period Paints per Pot 11,0 to 200 to 160 80 to 60 80 to	(suppression)	,	8.0 to 10.0		1.0 to 0.8	1/2 to 2/3	2.0 to 2.7		
Ispanese Beeties Accounty	Elm leaf beetles	plants per pot)	11.0 to	900 to 400	0.8 to 0.5	2/3 to 1	2.7 to 4.1		
Apply 15 ounces per 1000 sq ft 1	Japanese		12.0	200 to 160		;	1	Mix thoroughly before	
Leafhorpers seafminers Species	Beetles	· · · · · · · · · · · · · · · · · · ·	<u></u>	80 to 60				filling pots and	
Species 5,5 to 7,0 1600 to 2,5 to 2,0 1/2 to 2/3 2,0 to 2.7 1,7 to 4.1 1,7 to 5.0 1,5 to 1.0 900 to 400 1,0 to 0.8 1,5 to 1.0 2/3 to 1 1,0 to 1.1	.acebugs	Woody	2.0 to 3.0	4500 to	9.0 to 7.0	1/4 of 1/3	1.0 to 1.3	transplanting.	
Realimens Meanybugs Saylids Suspited Substitution of the Complex Substitution of the C	_eafhoppers	Perennial	3.5 to 5.0	3200	6.0 to 4.0	1/3 to 1/2	1.3 to 2.0		
Mealybugs Posyllids Coor mealybugs C	_eafminers	Species	5.5 to 7.0	1600 to	25 to 20	1/2 to 2/3	2.0 to 2.7		
Complex (such as Black Vine Weevil, Apopha Weevil, Chrupes (such as Black Vine Weevil)** Complex (such as Black Vine Weevil, Apopha Weevil, Chrupession)** Mitte Grub Larvae (such as Beetle, Chafers, Deetle) Vhiteflies Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5.0 to 10.0 radic and incorporate into the root zone. Minimize leaching and runout for at least 3 irrigations or 10 days, whichever is longer. Chafers, Deetle, Asiatic Garden Beetle, Asiatic Garden 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon to active minimize and incorporate into the root zone. Minimize leaching and runout for a teash of the active ingredient into the root zone. Minimize leaching and runout for at least 3 irrigations or 10 days, whichever is longer. Containerized 1 Gallon 125 to 100 4.4 to 2.2 2.1/2 to 5.0 to 10.0 to 200 days, whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5.0 to 10.0 to 200 foot splanting. On established plants, irrigate moderately and incorporate into the root zone. Minimize leaching and runout for at least 3 irrigations or one dischained in corporate into the root zone. Minimize leaching and runout for at least 3 irrigations or one dischained in corporate into the root zone. Minimize leaching and runout for at least 3 irrigations or one dischained in corporate into the root zone. Minimize leaching and runout for at least 3 irrigations or to get stablished plants, irrigate moderately after application to a for the root zone. Plants 2 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15 to 10.0 to 20 to 4.4 to 2.2 2.1/2 to 5.0 to 10.0 to 20 to 4.4 to 2.2 2.1/2 to 5.0 to 10.0 to 20 to 4.4 to 2.2 2.1/2 to 5.0 to 10.0 to 20 to 4.4 to 2.2 2.1/2 to 5.0 to 3.0 to 4.4 to 2.2 2.1/2 to 5.0 to 3.0 to 4.4 to 2.2 3.4 to 5.0 t	Mealybuas			1300	1				
Root Weevil Complex (such as Black Vine Weevil, Apopta Root Weevil) Apopta Root Weevil, Carus Root Weevil) White Grub Iarvae (such as Japanese Beetle, Chafers, European Chafer, Oriental Reactie Root Root Root Root Root Root Root Roo		(Three or		900 to 400	1				
Root Weevil, Apopka Weevil, Chrus Root Weevil, Apopka Weevil, Chrus Root Weevil, Apopka Weevil, Chrus Root Weevil)* Soft scale Thrips (suppression)* White Grub Iarvae (such as Dagelle, Chafers, European Chafer, Onental Beetle, Asiatic Garden Plants 2 Gallon 70 to 60 3.3 to 1.7 3-23 to 7-1/3 15.0 to 3.0 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 2 Feeding on foliage only. Thrips in buds and flowers will not be suppressed. 4 Apply a single or split (halved) application as a 36-inch band (18" either side of the planted row).	=	more plants		200 to 160	1.0 10 0.6	1 (0 1-1/2	4.1 (0 0.1		
Complex (such as Black Vine Weevil, Apopka Weevil, Chrus Root Weevil)** Not Weevil, Chrus Root Weevil)** Nothite Grub larvae (such as Japanese Beetle, Chalers, European Chaler, Oriental Beetle, Asiatic Garden Beetle) **Containerized 1 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7.1/3 15.0 to 30.0 10 days, whichever is longer. **Containerized 1 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7.1/3 15.0 to 30.0 15.5 to 37.0 Thirps in buds and flowers will not be suppressed. **For use on non-bearing citrus nursery stock **Vinite grub larvae **For use on non-bearing citrus nursery stock **Vinite grub larvae **Such as Japanese **For use on non-bearing citrus nursery stock **Vinite grub larvae **For use on non-bearing citrus nursery stock **Vinite grub larvae **Such as Japanese **For use on non-bearing citrus nursery stock **Nite grub larvae **Such as Japanese **For use on non-bearing citrus nursery stock **Nite grub larvae **Such as Japanese **For use on non-bearing citrus nursery stock **Nite grub larvae **Such as Japanese **For use on non-bearing citrus nursery stock **Nite grub larvae **Such as Japanese **For use on non-bearing citrus nursery stock **Nite grub larvae **Such as Japanese **Beetle, Masked Chalers, Christian as 36-band (18" either side of the planted row). **Japanese Beetle, Masked Chalers, Christian as 36-band (18" either side of the planted row). **Japanese Beetle, Masked Chalers, Christian as 36-band (18" either side of the planted row). **Japanese Beetle, Masked Chalers, Christian as 36-band (18" either side of the planted row). **Japanese Beetle, Masked Chalers, Christian as 36-band (18" either side of the planted row). **Japanese Beetle, Masked Chalers, Christian as 36-band (18" either side of the planted row). **Japanese Beetle, Masked Chalers, Christian as 36-band (18" either side of the planted row). **Japanese Beetle, Asiatic Garden Beetle, Asiatic Garden Beetle) **Japanese Beetle, Asiatic Garden Beetle, Asiatic Garden Beetle) **Japanese Beetle, Asiatic Garden Beetle	, •	per pot)	,2.0						
(such as Japanese Beetle, Aslatic Garden Beetle) Containerized 1 Gallon 70 to 60 3 3 to 1.7 3-23 to 7.173 15.0 to 3.0 to 2.0 to 1.0 5 Gallon 70 to 60 3 3 to 1.7 3-23 to 7.173 15.0 to 3.0 to 2.0 to 1.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to 2.0 to 2.0 to 1.0 to 1.0 to 2.0 to				55.55					
Vine Weevil, Apopka Meevil)** Apopka Meevil, Citrus Root Weevil)** Soft scale Phrips Commental crops grown in flats, benches, or beds Japanese Beetle, Chafers, European Chafer, Offental Beetle, Asiatic Sarden Beetle, Asiatic Flants Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 to 5.0 to 10.0 to 20.0 to 3 imgations prior to the active ingredient into the roo zone. White grub larvae such as Japanese Beetle, Asiatic Flants 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 1 Level tsp of Merit 1 G = 4.1 grams 3 teaspoo	•	ł	l '						
Apopka Weevil, Citrus Root Weevil)** Not scale hittings (suppression)* White Grub Japanese Beetle, Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle, Plants 3 Gallon 70 to 60 3.5 to 1.0 2.5 1.11/4 to 2.11/2 to 9.0 18.5 to 37.0 10.0 y.g. whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1.11/4 to 2.11/2 5.0 to 10.0 days, whichever is longer. Minimize leaching and nunout for at least 3 irrigations or 10 days, whichever is core media before planting. On established plants, irrigate moderately after application to move the active ingredient into the roo zone. Minimize leaching and nunout for at least 3 irrigations or 10 days, whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1.11/4 to 2.11/2 5.0 to 10.0 days, whichever is longer. Containerized 1 Gallon 125 to 100 4.4 to 2.2 2.1/2 to 5.0 10.0 to 20.0 make applications for the days whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1.11/4 to 2.11/2 5.0 to 10.0 days, whichever is longer. Containerized 1 Gallon 125 to 100 4.4 to 2.2 2.1/2 to 5.0 10.0 to 20.0 make applications for the second of the days whichever is longer. To be a stablished plants, irrigate moderately after application of 10 days, whichever is longer. To be a stablished plants, irrigate moderately after application of 10 days, whichever is longer. The plants 2 Gallon 125 to 100 4.4 to 2.2 2.1/2 to 5.0 10.0 to 2.0 make applications for the transport of the gallon of 10 days, whichever is longer. The plants 3 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7.1/3 15.0 to 30.0 to 2.0 make applications for the gallon days application and the second of the plants o								ingredient into the root	
Root Weevil)** Soft scale Thrips (suppression)* White Grub larvae (such as Japanese Beetle, Chafers, Conental Beetle, Asiatic Garden Plants 2 Gallon 2 Gallon 3 Gallon 7 to 60 3 3 to 1.7 5 Gallon 5 to to 30 2 0 to 1.0 4 4-1/2 to 9.0 1 18.5 to 37.0 1 level tsp of MERNIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. * Fer use on non-bearing citrus nursey stock * Field and Apply a single or split (halved) application as a 36-inch band (18" either should be followed by rainfall or irrigation. * Apply as a broadcast treatment and incorporate into the so or media before planting. On established plants, irrigate moderately after application to the own to accomplication on the ord to accomplication of the target pest. Irrigate application for the target pest. Irrigate ap	Apopka								
Soft scale Thrips Compared to the property of the plants o	Weevil, Citrus	ł	'		ł				
Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 15.0 to 10.0 to 20.0 16.0 to 30.0 15.0 to 30.0 1	Root Weevil)**					•			
Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 to 5.0 to 10.0 For optimum control to the target peaks allowed by after application to move the active ingredient into the root zone. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 to 5.0 to 10.0 For optimum control to make application to move the active ingredient into the root zone. Minimize leaching and runout for at least 3 irrigations or 10 days, whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 to 5.0 to 10.0 For optimum control, make application to move the active ingredient into the root zone. Minimize leaching and runout for at least 3 irrigations or 10 days, whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 to 5.0 to 10.0 For optimum control, make applications or 10 days, whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 2-1/2 to 5.0 to 10.0 For optimum control, make application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid moderately after application to move the larget peaks impaid to the larget peaks in the l	Soft scale								
White Grub (such as Japanese Beetle, Chafers, Cornetal Beetle, Asiatic Garden Plants 1 Gallon 2 Gallon 3	Thrips								
India, benches, or beds Inalia, benches, or beds Incorporate into the so or media before planting. On established plants, irrigate moderately after application to move the active ingredient into the roc zone. Minimize leaching and runout for at least 3 irrigations to or move the active ingredient into the roc zone. Minimize Inalia, Dente	(suppression)*		İ	Apply	15 ounces per	1000 sq ft			
Greetle, Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) Whitefiles Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5.0 to 10.0 for glanting. On established plants, irrigate moderately after application to move the active ingredient into the root zone. Minimize leaching and runout fra at least 3 irrigations on 10 days, whichever is longer. Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5.0 to 10.0 days, whichever is longer. Containerized 2 Gallon 125 to 100 4.4 to 2.2 2-1/2 to 5.0 10.0 to 20.0 to 2	White Grub								
Japanese Beetle, Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) Whiteflies Containerized 1 Gallon 260 to 230 5.0 to 2.5 1.1/4 to 2-1/2 5.0 to 10.0 to 20.0 10 days, whichever is longer. Containerized 2 Gallon 125 to 100 4.4 to 2.2 2.1/2 to 5.0 10.0 to 20.0 10 days, whichever is longer. Containerized 3 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30.0 10 days, whichever is longer. Containerized 5 Gallon 50 to 30 2.0 to 1.0 4-1/2 to 9.0 18.5 to 37.0 For optimum control, make applications or the target peets. Irrigate and early after application to move the active ingredient into the suppressed. Terest such as Japanese Beetle, Asiatic Garden Beetle, Asiatic Garden Beetle, Asiatic Garden Beetle)	larvae								
Beetle, Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) Whitefiles Containerized Plants 2 Gallon 125 to 100 4.4 to 2.2 2-1/2 to 5.0 10.0 to 20.0 10 days, whichever is longer. Containerized 2 Gallon 125 to 100 4.4 to 2.2 2-1/2 to 5.0 10.0 to 20.0 to	•	Oi beas					•		
Containerized Plants 2 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30. Por use application to move the active ingredient into the root 20ne. Containerized Plants 2 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30. To egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root 20ne. Minimize leaching and runout for at least 3 irrigations or 10 days, whichever is longer. Containerized Plants 2 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30. To 10.0 to 20.0 3 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30. To 99 hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root 20ne. 1 level tsp of MERIT 1 G = 4.1 grams 3 teaspoons =1 tablespoon * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. * For use on non-bearing citrus nursery stock Vinite grub larvae such as Japanese Beetle, Asiatic Chafers, Cincipean Chafer, Oriental Beetle, Asiatic Garden Beetle) * For use on an on-bearing citrus nursery stock Field and Forest Surseries Side of the row) prior to egg deposition. Use 2 00 to 2.75 ib/1000 feet of row in a 36"-band (18" either side of the planted row).	•								
European Chafer, Oriental Beetle, Asiatic Garden Beetle) Whiteflies Containerized Plants 2 Gallon Plants 2 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30.0 To 4-1/2 to 9.0 To 10.0 to 20.0 Sallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30.0 To 20.0 To 20 Sallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30.0 To 20.0 To 20 Sallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30.0 To 20.0 To									
Chafer, Oriental Beetle, Asiatic Garden Beetle) Whitefiles Containerized 1 Gallon Plants 2 Gallon 3 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30.0 10.0 to 20.0 15.0 to 30 2.0 to 1.0 4-1/2 to 9.0 18.5 to 37.0 15.0 to 30.0 16.0 to 30.0 to 3			<u> </u>						
Oriental Beetle, Asiatic Garden Beetle) Whitefiles Containerized Plants 2 Gallon 125 to 100 4.4 to 2.2 2-1/2 to 5.0 to 10.0 For optimum control, a flarer such as Japanese Beetle, Asiatic Garden Beetle) White grub Beetle, Asiatic Chafers, Curiopean Chafer, Oriental Beetle, Asiatic Garden Beetle) Oriental Beetle, Asiatic Garden Beetle) Oriental Beetle, Asiatic Garden Beetle) Accordance A Gallon A Gallo	•							1	
Reetle) Whitefiles Containerized Plants 2 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5.0 to 10.0 Plants 2 Gallon 70 to 60 3.3 to 1.7 3.2/3 to 7-1/3 15.0 to 30.0 to 29.0 to 2.0 to 1.0 4.4 to 2.2 2-1/2 to 5.0 10.0 to 20.0 18.5 to 37.0 longer. Teeding on foliage only. Thrips in buds and flowers will not be suppressed. Feeding on foliage only. Thrips in buds and flowers will not be suppressed. Feeding on foliage only. Thrips in buds and flowers will not be suppressed. Fero use on non-bearing citrus nursery stock White grub larvae such as Japanese Beetle, Masked Chafers, Oriental Beetle, Asiatic Garden Beetle) Reaching and runout for at least 3 irrigations of 10 days, whichever is longer. For the containerized 1 Gallon 2 Gallon 125 to 100 4.4 to 2.2 2-1/2 to 5.0 10.0 to 20.0 10.0 to 20.0 10.0 to 20.0 15.0 to 30.0 to 20.0 to 20		}	ł						
Beetle) Whiteflies Containerized Plants 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5.0 to 10.0 10.0 to 20.0 16.0 to 30.0 15.0 to 30.0	Beetle, Asiatic								
Whiteflies Containerized Plants Containeri									
Vitite flies Containerized 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5.0 to 10.0 For optimum control, make applications price to egy hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone. I level tsp of MERIT 1 G = 4.1 grams 3 teaspoons =1 tablespoon I level tsp of MERIT 1 G = 4.1 grams 3 teaspoons =1 tablespoon I level tsp of MERIT 1 G = 4.1 grams 3 teaspoons =1 tablespoon I level tsp of MERIT 1 G = 4.1 grams 3 teaspoons =1 tablespoon I level tsp of MERIT 1 G = 4.1 grams 3 teaspoons =1 tablespoon Feeding on foliage only. Thrips in buds and flowers will not be suppressed. Field and Forest Apply a single or split (halved) application as a 36-inch band (18" either side of the planted row). Field and Forest Apply a single or split (halved) application. Use 2 00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row). Apply May through July. For optimum control, make applications price to egy deposition. Use 2 00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row).	Beetle)								
Containerized Plants Containerized Plants 1 Gallon 260 to 230 1.25 to 100 1	Whiteflies							1	
Plants 2 Gallon 3 Gallon 70 to 60 3 Gallon 5 Gallon 6 Gallon 7 Ga		Containerized	1 Gallon	260 to 230	5.0 to 2.5	1-1/4 to 2-1/2	5.0 to 10.0		
3 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15.0 to 30.0 18.5 to 37.0 18.5 to 37.5 to 37.	1	Diameter 1	,		l - · ·			make applications prior	
The level to be described and such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Masked Chafers, European Beetle) The level to be the planted for the planted row). Solo 30 2.0 to 1.0 4-1/2 to 9.0 4-1/2 to 9.0 18.5 to 37.0 1								to egg hatch of the	
The level tsp of MERIT 1 G = 4.1 grams 1 level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. ** For use on non-bearing citrus nursery stock White grub larvae such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) 1 level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon 4 Feeding on foliage only. Thrips in buds and flowers will not be suppressed. 5 Feeding on foliage only. Thrips in buds and flowers will not be suppressed. 5 Field and Forest Side of the row) prior to egg deposition. Use 2 00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row). 5 Apply May through July. For optimum control, tleasinent should be followed by rainfall or irrigation. 6 Chafer, Oriental Beetle, Asiatic Garden Beetle)									
# Feeding on foliage only. Thrips in buds and flowers will not be suppressed. **For use on non-bearing citrus nursery stock White grub larvae such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) ### Reeding on foliage only. Thrips in buds and flowers will not be suppressed. ### Apply a single or split (halved) application as a 36-inch band (18" either side of the planted row). ### Apply May through July. For of through side of the planted row). ### Apply May through July. For of through side of the planted row). ### Apply May through July. For of through side of the planted row). ### Apply May through July. For of through side of the planted row). ### Apply May through July. For of	:	1	Janon	JU 10 JU	2.0 10 1.0	→ 112 tO 9.0	10.5 (0 57.0		
the root zone. 1 level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. **For use on non-bearing citrus nursery stock White grub larvae such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) the root zone. 1 level tsp of MERIT 1 G = 4.1 grams 3 teaspoons = 1 tablespoon * Feeding on foliage only. Thrips in buds and flowers will not be suppressed. **Apply a single or split (halved) application as a 36-inch band (18" either side of the row) prior to egg deposition. Use 2 00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row). Apply May through July. For oftmum control, tleasment should be followed by rainfall or irrigation. **Control and the root zone. **Apply May through July. For oftmum control, tleasment should be followed by rainfall or irrigation. **Apply May through July. For oftmum control, tleasment should be followed by rainfall or irrigation. **Apply May through July. For oftmum control, tleasment should be followed by rainfall or irrigation. **Apply May through July. For oftmum control, tleasment should be followed by rainfall or irrigation. **Apply May through July. For oftmum control, tleasment should be followed by rainfall or irrigation. **Apply May through July. For oftmum control, tleasment should be followed by rainfall or irrigation. **Apply May through July. For oftmum control tleasment should be followed by rainfall or irrigation.									
* Feeding on foliage only. Thrips in buds and flowers will not be suppressed. ** For use on non-bearing citrus nursery stock White grub larvae Such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) ** Feeding on foliage only. Thrips in buds and flowers will not be suppressed. ** For use on non-bearing citrus nursery stock Apply a single or split (halved) application as a 36-inch band (18" either side of the row) prior to egg deposition. Use 2 00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row). Apply May through July. For of thrum control, treament should be followed by rainfall or irrigation. ** ** ** ** ** ** ** ** ** ** ** ** *		ļ	ļ						
* Feeding on foliage only. Thrips in buds and flowers will not be suppressed. ** For use on non-bearing citrus nursery stock White grub larvae Field and Forest Side of the row) prior to egg deposition. Use 2 00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row). Apply May through July. For on this intercept of row in a 36"-band (18" either side of the planted row). Apply May through July. For on this intercept of row in a 36"-band (18" either side of the planted row). Apply May through July. For on this intercept of row in a 36"-band (18" either side of the planted row).		 	4 (5):-1		C = 4.4 ==== :::	24		L	
** For use on non-bearing citrus nursery stock White grub larvae Forest Such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) ** For use on non-bearing citrus nursery stock Apply a single or split (halved) application as a 36-inch band (18" either side of the row) prior to egg deposition. Use 2.00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row). Apply May through July. For optimum control, iteament should be followed by rainfall or irrigation.		* Fooding of						OH	
White grub larvae Field and Forest Such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)		_		-		ot be suppressed			
larvae such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) Forest Nurseries side of the row) prior to egg deposition. Use 2 00 to 2.75 lb/1000 feet of row in a 36"-band (18" either side of the planted row). July. For oftimum control, fleatment should be followed by rainfall or irrigation.	White and		, _			n ac a 36 inch ha	nd (19" aitha-	Apply May through	
Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)									
Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)							S TOOU IEEL UI		
Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)									
Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)			1						
Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)								1 7	
European Chafer, Oriental Beetle, Asiatic Garden Beetle)			[
Chafer, Oriental Beetle, Asiatic Garden Beetle)	•	ļ						l lade	
Oriental Beetle, Asiatic Garden Beetle)			ĺ					1	
Garden Beetle)	•	ĺ	ĺ					1	
Beetle)	Beetle, Asiatic	•							
								,	
0000	Beetle)								
								,,,,,	

•

6/4

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Dispose of empty containers by incineration in a sanitary landfill or if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response telephone number is 800-414-0244, or contact Chemtrec at 800-424-9300.

M - 8944b

Bayer Corporation Garden and Professional Care Box 4913 Kansas City, MO 64120-0013 (800) 842-8020 http://www.nobugs.com

