Please sead instructions on s	reverse before completing form.		Approved. OMB No. 2070-0060. Approval expire	043 s 2-28-95	
\$EPA	United States Environmental Protectio Washington, DC 204	n Agency wolfold	Registration OPP Identifier Nu	mber	
	Applicatio	n for Pesticide - Se	ection I		
1. Company/Product Number 3125–457	•		wards. Jr.	on stricted	
 Company/Product (Name) Provado 1.6 Floring 		PM# 19		İ	
5. Name and Address of Applicant (Include ZIP Code) Bayer Corporation P.O. Box 4913, 8400 Hawthorn Rd. Kansas City, MO 64120-0013 Check if this is a new address		(b)(i), my produ to: EPA Reg. No	EPA Reg. No.		
		Product Nam	IE		
Section - II Amendment - Explain below. Resubmission in response to Agency letter dated "Me Too" Application. Notification - Explain below. Other - Explain below.					
Explanation: Use additional page(s) if necessary. (For section I and Section II.) APR 2 3 1996 See attached					
		Section - III			
1. Material This Product Wil	l Be Packaged In:				
Child-Resistant Packaging Yes* No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt. No. per container	Water Soluble Packaging Yes No If "Yes" No. p Package wgt Conta	Metal Plastic Glass		
3. Location of Net Contents Label	Information 4. Size(s) Ret	ail Container	5. Location of Label Directions On Label On Labeling accompanying product		
6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Continue IV					
Section - IV					
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Name Karen S. Cain Title Registration Specialist (816) 242-2838					
	Certifica ements I have made on this form and ny knowingly false or misleading star law.	all attachments thereto are			
2. Signature	- Cair	Manager, Regi	istrations	(((e)	
4. Typed Name of John S. Thornt	on	5. Date 3/26/96	######################################	- 2 . 5 = - 2 . 1 ×	
EPA Form 8570-1 (Rev. 8-9	4) Previous editions are obsolete.	i une compressed in the compression of the property of the compression of the property of the compression of	White - EPA File Copy (original) Yellow - Appl	icant Con	

Gallons

U.S. LABEL

Reason to Issue: To add pests to cotton, potato and cole crops.

Date of Pre-Reg Draft: 03/22/96 (C)

Provado® 1.6 Flowable

Insecticide

For control of certain insects infesting various crops. Shake well before using.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]- N-nitro-2-imidazolidinimine	17.4%
INERT INGREDIENTS	82.6%
	100 00/

Contains 1.6 pounds of imidacloprid per gallon EPA Reg. No. 3125-457

PA Reg. No. 3125-457 Net Contents:

STOP - Read The Label Before Use

STOP - Read The Label Before Use KEEP OUT OF REACH OF CHILDREN

CAUTION

AMENDMENT To Previously Registered Labeling

	RECOMMENDED APPLICATIONS				
CROP	PEST	RATE PER APPLICATION	REMARKS		
Cotton	Aphids Fleahopper Plant bugs	Foliar Application 3.75 fl oz/acre	Apply specified dosage per acre as pests begin to build. Two applications at a 7 to 10 day interval may be required to achieve control. Scout fields and retreat if needed.		
	Suppression: Whiteflies (including sweet potato/ silverteaf whitefly)	Foliar Application 3.75 fl oz/acre	Begin applications when whitefly adults appear prior to development of nymphs. Make application on a 7 day interval as long as pest pressure continues. For resistance management purposes, switch to a pesticide of alternative chemistry to continue control after a maximum of 5 applications.		

Foliar Applications: Thorough coverage with direct contact of the spray material to the target pests is required for optimum control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage. Applications made with less than 5 gallons per acre may result in slower activity and/or less overall control from a single application than an application made with higher gallonages. Allow 7 days between applications and 14 days between last application and harvest.

Regardless of formulation or method of application, apply no more than 0.5 lb active ingredient of ADMIRE or PROVADO per acre per season, including seed treatment, soil <u>and</u> foliar uses. Do not apply more than a total of 6 field applications per season. Do not graze treated fields after any application of PROVADO.

المتحدد فا		P =
- 2 '		A
· ~ .		~11 -2
26		

- 1 .	. —	- 1 -
-	• •	1

RECOMMENDED APPLICATIONS			
CROP	PEST	RATE PER APPLICATION	REMARKS
Potato Aphids . Colorado potato beetle Fleabeetles Potato leafhopper Psyllids		Foliar Application 3.75 fl oz per Acre	Foliar Application: For optimal control, good coverage of the foliage is needed.
	Potato leafhopper	For aphid; and leafhopper pysilid and fleabeetle control make applications by ground equipment only. For Colorado potato beetle control east of the Rockles, applications may be made by ground or by air. For Colorado potato beetle control west of the Rockles, make applications by ground equipment only. For aphids, and leafhoppers, psyllids and fleabeetles apply specified dosages as pests begin to build. Two applications at a 7-day interval may be required to achieve control. Scout fields and retreat if needed. Allow at least 7 days between foliar applications. Allow at least 7 days between last application and harvest. A total of 15 fl oz (0.2 lb active ingredient) of PROVADO 1.6 Flowable per acre per season may be applied as a foliar spray.	
		For resistance management purposes, a PROVADO foliar application following a soil application of ADMIRE in the same crop is not recommended.	
		Note: Regardless of formulation or type of application (soil or foliar), do not apply more than a total of 0.31 lb active ingredient of ADMIRE or PROVADO per acre per season.	
Broccoli Brussels sprouts Cabbage Cauliflower Broccoli, Chinese (gai lon) Broccoli raab, (rapini) Cabbage, Chinese (bok choy, napa) Cabbage, Chinese mustard (gal choy) Collards Kale	Fleabeetles Whiteflies (Including Sweetpotato	Foliar Application 3.75 fl oz per acre	FOLIAR APPLICATION Aphid Control: Apply specified dosage as needed for control. Good coverage of the foliage is necessary for optimal control. —A spray adjuvant may be used to improve coverage. For best results, time application before a damaging population becomes established.
	Silverleaf Whitefly Control: Apply specified rate on a 5 to 7 day schedule at the first appearance of the whitefly in the crop. Good coverage of the foliage is necessary for optimal control. A spray adjuvant may be used to improve coverage. Continue applications until whiteflies are no longer infesting the crop.		
		PROVADO 1.6 Flowable will not knock down heavy aphid or whitefly populations. This product may be tank mixed with other pesticides as recommended for control of other pests or for improved control of whitefly. Allow at least 7 days between last application and harvest. Do not apply to vegetables grown for seed. Allow at least 5 days between foliar applications.	
Kohlrabi Lettuce. (Head and Leaf)		Do not apply more than a total of 18.75 fluid ounces of PROVADO 1.6 Flowable per year as foliar sprays (See NOTE).	
Mustard greens Rape greens		For resistance management purposes, a PROVADO foliar application following a soil application ADMIRE in the same crop is not recommended.	

Bayer Corporation

Crop Protection Products

Box 4913

Kansas City, MO 64120-0013

