

Provado 1.6 F

PM 19

3125-457

1/2

457-8767.YLD

U.S. LABEL

Base Pre-Reg (8767)

Reason to Issue: To replace vegetables grown for seed restriction with exception statement.

Date of Draft: 03/29/96 (Pre-Reg) (C)

Provado® 1.6 Flowable Insecticide

For control of certain insects infesting various crops.
Shake well before using.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	17.4%
INERT INGREDIENTS	82.6%
	100.0%

Contains 1.6 pounds of imidacloprid per gallon


EPA Reg. No. 3125-457

Net Contents: 1 Gallons

**STOP - Read The Label Before Use
KEEP OUT OF REACH OF CHILDREN**

CAUTION

AMENDMENT To Previously Registered Labeling



2/2

RECOMMENDED APPLICATIONS			
CROP	PEST	RATE PER APPLICATION	REMARKS
VEGETABLES (Direct-seed or Transplant) (Except those grown for seed) Eggplant Pepper (Includes Bell, Chili, Cooking, Pimientos and Sweet) Tomato Ground cherry Pepinos Tomatillo	Aphids Colorado potato beetle Whiteflies (Including Sweetpotato or Silverleaf whitefly)	Foliar Application 3.75 fl oz per acre	FOLIAR APPLICATION Aphid or Colorado Potato Beetle Control: Apply specified dosage as needed for control. Good coverage of the foliage is necessary for optimal control. A spray adjuvant may be used to improve coverage. For best results, time application before a damaging population becomes established.
			Silverleaf Whitefly Control: Apply specified rate on a 5 to 7 day schedule at the first appearance of the whitefly in the crop. Good coverage of the foliage is necessary for optimal control. A spray adjuvant may be used to improve coverage. Continue applications until whiteflies are no longer infesting the crop.
			PROVADO 1.6 Flowable will not knock down heavy aphid or whitefly populations. This product may be tank mixed with other pesticides as recommended for control of other pests or for improved control of whitefly. Foliar applications may be applied up to and including day of harvest. Do not apply to vegetables grown for seed. Allow at least 5 days between foliar applications.
			Do not apply more than a total of 18.75 fluid ounces of PROVADO 1.6 Flowable per year as foliar sprays (See NOTE).
		For resistance management purposes, a PROVADO foliar application following a soil application of ADMIRE in the same crop is not recommended.	
VEGETABLES (Direct-seed or Transplant) (Except those grown for seed) Broccoli Brussels sprouts Cabbage Cauliflower Broccoli, Chinese (<i>gai lan</i>) Broccoli raab, (<i>rapini</i>) Cabbage, Chinese (<i>bok choy, napa</i>) Cabbage, Chinese mustard (<i>gai choy</i>) Collards Kale Kohlrabi Lettuce (Head and Leaf) Mustard greens Rape greens	Aphids Flea beetles Whiteflies (Including Sweetpotato or Silverleaf whitefly)	Foliar Application 3.75 fl oz per acre	FOLIAR APPLICATION Aphid Control: Apply specified dosage as needed for control. Good coverage of the foliage is necessary for optimal control. A spray adjuvant may be used to improve coverage. For best results, time application before a damaging population becomes established.
			Silverleaf Whitefly Control: Apply specified rate on a 5 to 7 day schedule at the first appearance of the whitefly in the crop. Good coverage of the foliage is necessary for optimal control. A spray adjuvant may be used to improve coverage. Continue applications until whiteflies are no longer infesting the crop.
			PROVADO 1.6 Flowable will not knock down heavy aphid or whitefly populations. This product may be tank mixed with other pesticides as recommended for control of other pests or for improved control of whitefly. Allow at least 7 days between last application and harvest. Do not apply to vegetables grown for seed. Allow at least 5 days between foliar applications.
			Do not apply more than a total of 18.75 fluid ounces of PROVADO 1.6 Flowable per year as foliar sprays (See NOTE).
		For resistance management purposes, a PROVADO foliar application following a soil application of ADMIRE in the same crop is not recommended.	
NOTE: Regardless of the type of application (soil or foliar) or type of formulation, do not apply more than 0.5 lb ai of ADMIRE or PROVADO per acre per year.			