264214/382



U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (R7505C)

401 "M" St., S.W. Washington, D.C. 20460

NOTICE OF PESTICIDE:

\_\_\_ Reregistration

(under FIFRA, as amended)

EPA Req Number

0 6 SEP 1995

3125-462

Term of Issuance:

Conditional

Name of Pesticide Product:

Tempo 20 WP Golf Course Insecticide

Name and Address of Registrant (include ZIP Code):

BAYER Corp.

PO Box 4913

8400 Hawthorn Rd.

Kansas City, MO 64120-0013

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
  - 2. Make the following label changes:
    - a. Revise the EPA Registration Number to read, "EPA Req. No. 3125-462".
    - b. Under the subheading NOTE add the additional exclusionary statement: "Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes."

Signature of Approving Official:	Date:

EFA Form 8570 6

- Change the buffer zone statement to read as follows: C. "...do not make applications within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds."
- Under recommended applications you should qualify the d. site "Turfgrass" with "(including golf courses, around home lawns, business and office complexes, multi-family residential complexes, airports, cemeteries, parks, playgrounds and athletic fields, and other landscape turfgrass).
- Under Environmental Hazards delete "For terrestrial е. uses." This phrase incorrectly implies that your product has aquatic uses.
- 3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca Product Manager (13) Insecticide-Rodenticide Branch Registration Division (H7505C)

CONCURRENCES							
SYMBOL							
SURNAME N		Ì		<b>,</b>			
DATE			•••••••	*************	•••••		
EPA Form 1320-1A	1.00		L	Paradad Barra	l	OF	FICIAL FILE CO

XXX-8487.YLD

U. S. LABEL

Base Pre-Reg. (8487)

Reason to Issue: To propose registration of product for golf course use.

Date of Draft: 04/04/95 Pre-Reg(S)

## RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms.

For retail sale to, and use only by, Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

# TEMPO 2 Golf Course Insecticide

FOR COMMERCIAL USE ONLY FOR BROAD-SPECTRUM CONTROL OF INSECT PESTS INFESTING TURF, TREES AND ORNAMENTALS ON RESIDENTIAL AND COMMERCIAL SITES INCLUDING **GOLF COURSES** 

**ACTIVE INGREDIENT:** 

Cyfluthrin, cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-24.3% dimethylcyclopropanecarboxylate ..... INERT INGREDIENTS: ..... 75.7% 100.0%

(This product contains xylene-range, aromatic solvents)

Contains 2 pounds cyfluthrin per gallon.

EPA Est. 11556-KS-1 EPA Reg. No. 3125-XXX

Net Contents:

Milliliters

STOP - Read the Label Before Use

## Keep Out of the Reach of Children WARNING

See Rear Panel for Statements of Practical Treatment and Other Precautionary Statements

## **AVISO**

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)



ACCEPTED with COMMENTS in EPA Letter Dates

0 6 SEP 1995

Under the Federal Insecticide, Forgicide, and Rodentic de Act as amended, for the petitie registered under EPA Rec. No.

Course Insecticide

3125-162

## 477

## PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## WARNING

May cause eye injury. Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on clothing. Avoid breathing vapors or spray mist.

Do not contaminate feed or food. Keep out of reach of children.

Wear goggles, face shield, or safety glasses, when handling the undiluted material. Wash hands, arms, and face thoroughly with soap and warm water after handling. Remove and wash contaminated clothing with soap and hot water before reuse.

## STATEMENTS OF PRACTICAL TREATMENT

If in eyes: Hold eyelids open and flush with plenty of water. Get medical attention. If swallowed: Call a physician or Poison Control Center. Do not induce vomiting. Vomiting may cause an aspiration hazard. If on skin: Wash thoroughly with soap and water. Get medical attention if irritation occurs.

NOTE TO PHYSICIAN: Treat the patient symptomatically, taking into consideration that this formulation contains 34% aromatic petroleum distillates.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply TEMPO 2 or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using TEMPO 2 Golf Course Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS AND/OR INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE

INJURY TO OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Do not formulate this product into other end-use products.

When used at recommended rates and as directed under RECOMMENDED APPLICATIONS, TEMPO 2 Insecticide will control the designated pests on trees, shrubs, foliage plants and flowers in outdoor landscaped and interior plantscapes where these plants are grown. TEMPO 2 Insecticide also controls designated pests in turf grasses.

NOTE: Not for use in commercial greenhouses, nurseries, sodfarms or grass grown for seed.

**APPLICATION TO ORNAMENTALS: TEMPO 2 Insecticide** mixes read'ly with water and may be used in all types of spray equipment. Mix product with the required amount of water and apply as a dilute spray application to the point of run-off. When spraying hard-to-wet foliage such as holly, pine or ivy, the addition of a spreader/sticker may enhance both knock-down and residual activity. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed, as would be used in a dilute application. Phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions and no phytotoxicity has been observed. It is advised to pre-spray a selection of ornamentals if local use experience is unavailable. Time applications to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

APPLICATIONS ON TURF: TEMPO 2 Insecticide mixes readily with water and/or various fertilizer solutions and

can be used in appropriate spray equipment. Applications should be made with accurately calibrated equipment and applied uniformly, using a coarse, large droplet, spray application procedure. Do not allow drift onto ponds, streams, or lakes. For optimum control of armyworms, cutworms and sod webworms, delay irrigation and mowing for at least 24 hours following the application. Keep children and pets off treated areas until lawn has dried following the application.

COMPATIBILITY: TEMPO 2 Insecticide has been found to be compatible with all commonly used fungicides, miticides, liquid fertilizers and other insecticides. Check physical compatibility using the correct proportion of products in a small jar if local experience is unavailable.

Do not apply this product through any type of irrigation system.

	RECOMMENDED APPLICATIONS			
CRO?	PEST	DOSAGE TEMPO 2	REMARKS	
Landscape Ornamentals (Including Trees, Shrubs, Evergreens, Flowrers, Foliage Plants)	Armyworms Azalea caterpillars Bagworms Bristly roseslugs California oakworms Cankerworms Crickets* Cutworms Elm spanworms Flies* Fungus gnats Gypsy moth larvae Lace bugs Leaf-feeding caterpillars Midges* Mosquitoes* Oleander moth larvae Pillbugs Pine shoot moths Pine tip moths Redhumped caterpillars Sawfly larvae Sowbugs Spiders* Striped oakworms Tent caterpillars Tussock moth larvae Webworms Yellownecked caterpillars	milliliters (1 fl oz) Per 100 Gal.	Applications should be started prior to establishment of high pest populations and reapplication made as necessary. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader/ sticker at recommended rates may enhance control of insects on certain species of ornamentals having hard-to-wet surfaces.  *The use of TEMPO 2 Insecticide for the control of ants, crickets, spiders, midges, wasps, flies and mosquitoes is limited to ornamental areas and areas adjacent to buildings. Apply as a general spray, concentrating on damp areas beneath shrubbery, around foundations and in areas of tall grass or weeds. Do not apply to any food crop. Remove any animal feeding dishes prior to treatment and do not allow spray or drift to contact fish-bearing water.	

6	37	7
---	----	---

	RECOMMENDED APPLICATIONS				
CROP	PEST	DOSAGE TEMPO 2	REMARKS		
Landscape Ornamentals (Continued) (Including Trees, Shrubs, Evergreens, Flowers, Foliage Plants)	Ants* Aphids Boxelder bugs Budworms Casebearers Clover mites Cockroaches** E!m leaf beetles Flea beetles Grasshoppers Japanese beetles (adult) June beetles (adult) Leafhoppers Leafrollers Leaf skeletonizers Mealybugs Orchid weevil Pear psylla Peppertree psyllid Plant bugs Scale insects (crawler stages) Spittlebugs Striped beetles Thrips Ticks (including deer tick) Tussock moth larvae Wasps* Whiteflies	45 milliliters (1.5 fl oz) Per 100 Gal.	Applications should be started prior to establishment of high pest populations and reapplication made as necessary. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having hard-to-wet foliage.  *The use of TEMPO 2 Insecticide for the control of ants, crickets, spiders, midges, wasps, flies and mosquitoes is limited to ornamental areas and areas adjacent to buildings. Apply as a general spray, concentrating on damp areas beneath shrubbery, around foundations and in areas of tall grass or weeds. Do not apply to any food crop. Remove any animal feeding dishes prior to treatment and do not allow spray or drift to contact fish-bearing water.  **Outdoor cockroach control (including Asian cockroaches) is obtained by spraying hiding or resting places such as under shrubbery, in ground covers and mulching materials, in hollow trees, etc.		
CROP	PEST	DOSAGE TEMPO 2	REMARKS		
Turfgrass	Ants Armyworms Crickets Cutworms Earwigs Grasshoppers Japanese beetles (adult) Sod webworms Ticks (including deer tick)	4 ml per 1000 sq ft or 5.9 fl oz per acre	Apply when pests first appear. Reapply when necessary based on pest reinfestations not exceeding 6 applications per year. Delay watering or mowing for 24 hours for optimum control of armyworms, cutworms, and sod webworms. Keep children and pets off treated areas until lawn has dried following the application.  *Aids in control only.		
	Bluegrass billbug (adult) Chinch bugs Chiggers Fire ants* Fleas Hyperodes weevil (adult) Mole crickets Scorpions	6 ml per 1000 sq ft or 8.75 fl oz per acre	And in control only.		

7 }

#### RESTRICTIONS FOR TURFGRASS APPLICATION

Do not apply this product by aerial application or by any type of chemigation system. Do not apply when the average windspeed is greater than 15 mph.

To avoid risk to sensitive aquatic areas, do not make applications within 25 feet of ponds, lakes, marshes, permanent streams, rivers, estuaries, and marine habitats.

Apply product using spray nozzles which produce a coarse droplet size, such as flood jet nozzles. Do not apply when turfgrass areas are water-logged or the soil is saturated with water (i.e., will not accept irrigation). Use this product only as directed on label.

### STORAGE AND DISPOSAL

Not for Storage in or Around the House

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

Container Disposal: Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Storage: Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is 'eaking invert to prevent leakage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides above. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response Team Telephone No. is 816-242-2582, or contact Chemtrec at 800-424-9300.

Sold by: Bayer Corporation Specialty Products Box 4913 Kansas City, MO 64120-0013