

KILLS	(ART: aphid, cabbage looper, tomato hornworm, whitefly)																																													
	<table border="0"> <tr> <td>Ants</td> <td>Japanese beetles (adult)</td> </tr> <tr> <td>Aphids</td> <td>Lacebugs</td> </tr> <tr> <td>Armyworms</td> <td>Leaf-feeding beetles</td> </tr> <tr> <td>Bagworms</td> <td>Leaf-feeding caterpillars</td> </tr> <tr> <td>Boxelder bugs</td> <td>Leafhoppers</td> </tr> <tr> <td>Cabbage loopers</td> <td>Leafminers</td> </tr> <tr> <td>Caterpillars</td> <td>Mealybugs</td> </tr> <tr> <td>Chinchbugs</td> <td>Midges</td> </tr> <tr> <td>Clover mites</td> <td>Mosquitoes</td> </tr> <tr> <td>Cockroaches</td> <td>Pillbugs</td> </tr> <tr> <td>Colorado potato beetles</td> <td>Plant bugs</td> </tr> <tr> <td>Corn borers</td> <td>Scale insects (crawler stages)</td> </tr> <tr> <td>Corn earworms</td> <td>Sowbugs</td> </tr> <tr> <td>Corn rootworms (adult)</td> <td>Spiders</td> </tr> <tr> <td>Crickets</td> <td>Spittle bugs</td> </tr> <tr> <td>Cutworms</td> <td>Stinkbugs</td> </tr> <tr> <td>Earwigs</td> <td>Sunflower beetles</td> </tr> <tr> <td>Elm leaf beetles</td> <td>Sunflower moths</td> </tr> <tr> <td>Flea beetles</td> <td>Thrips</td> </tr> <tr> <td>Flies</td> <td>Tomato Worms</td> </tr> <tr> <td>Garden webworms</td> <td>Wasps</td> </tr> <tr> <td>Grasshoppers</td> <td>Weevils</td> </tr> <tr> <td>Gypsy moth larvae</td> <td>Whiteflies</td> </tr> </table>	Ants	Japanese beetles (adult)	Aphids	Lacebugs	Armyworms	Leaf-feeding beetles	Bagworms	Leaf-feeding caterpillars	Boxelder bugs	Leafhoppers	Cabbage loopers	Leafminers	Caterpillars	Mealybugs	Chinchbugs	Midges	Clover mites	Mosquitoes	Cockroaches	Pillbugs	Colorado potato beetles	Plant bugs	Corn borers	Scale insects (crawler stages)	Corn earworms	Sowbugs	Corn rootworms (adult)	Spiders	Crickets	Spittle bugs	Cutworms	Stinkbugs	Earwigs	Sunflower beetles	Elm leaf beetles	Sunflower moths	Flea beetles	Thrips	Flies	Tomato Worms	Garden webworms	Wasps	Grasshoppers	Weevils	Gypsy moth larvae
Ants	Japanese beetles (adult)																																													
Aphids	Lacebugs																																													
Armyworms	Leaf-feeding beetles																																													
Bagworms	Leaf-feeding caterpillars																																													
Boxelder bugs	Leafhoppers																																													
Cabbage loopers	Leafminers																																													
Caterpillars	Mealybugs																																													
Chinchbugs	Midges																																													
Clover mites	Mosquitoes																																													
Cockroaches	Pillbugs																																													
Colorado potato beetles	Plant bugs																																													
Corn borers	Scale insects (crawler stages)																																													
Corn earworms	Sowbugs																																													
Corn rootworms (adult)	Spiders																																													
Crickets	Spittle bugs																																													
Cutworms	Stinkbugs																																													
Earwigs	Sunflower beetles																																													
Elm leaf beetles	Sunflower moths																																													
Flea beetles	Thrips																																													
Flies	Tomato Worms																																													
Garden webworms	Wasps																																													
Grasshoppers	Weevils																																													
Gypsy moth larvae	Whiteflies																																													
WHERE TO USE	Vegetable and flower gardens, shrubs, groundcovers, lawns and around homes																																													
WHEN TO USE	When insects or damage first appear For severe infestations, repeat every 7 to 14 days as necessary																																													
ART: GRAPHIC OF PHONE	Questions or comments, call toll-free (877) 229-3724 www.AdvancedGarden.com																																													

Ingredients

ACTIVE INGREDIENT:

Cyfluthrin	0.003 %
OTHER INGREDIENTS*	<u>99.997 %</u>
	100.000 %

* Other ingredients include: Water, organic solvent (less than 0.1%, to dissolve the active ingredient for mixing) and surfactants (to keep the active ingredient dispersed in water).

EPA Reg. No. 3125-503



EPA Est. No. XXXX-AA-1

[INSIDE BOOKLET]

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

- Harmful if swallowed or inhaled.
- Causes eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Avoid breathing spray mist.
- Wash thoroughly with soap and water after handling.



First Aid

The table below describes the first aid procedures for incidents involving TEMPO Ready-to-Use Insecticide.

IN THE CASE OF ACCIDENTAL...	IMMEDIATELY...
Swallowing	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Contact with eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
Inhalation	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
<p>NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free (877) 229-3763 for medical emergency information. Active Ingredient: 0.003% Cyfluthrin [CAS # 68350-37-5]</p>	

Note to Physicians: No specific antidote is available. Treat the patient symptomatically.

Environmental Hazards

- This pesticide is toxic to fish. Do not apply directly to water.
- This pesticide is highly toxic to bees. Do not apply TEMPO Ready-to-Use Insecticide when bees are active.
- Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For best results, read and follow all label directions.

Before You Use

Read and follow these directions when using:

- Apply the product to flowering plants during early morning or late evening when bees are not present.
- Do not spray the product into fish pools, ponds, streams, or lakes.
- Do not use in small, enclosed areas where spray can be inhaled.

How to Use

Shake well before using.

To apply TEMPO Ready-to-Use Insecticide:

WHEN TO USE	When insects or damage first appear.
HOW TO USE	Adjust nozzle to desired spray pattern. Hold sprayer about 12 inches from foliage. Spray leaves, stems, and tender new shoots. To kill insects that may hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray, but not dripping. (ART: spraying undersides of leaves)
HOW OFTEN TO USE	Kills insects on contact and keeps on killing for weeks. For severe infestations, repeat every 7 to 14 days as necessary.

FLOWERS, SHRUBS, GROUND COVERS		
PEST		HOW TO APPLY
Aphids	Leaf-feeding caterpillars	Spray foliage, tender new shoots and the underside of leaves thoroughly. To kill insects that hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray droplets, but not dripping. (ART)
Bagworms	Leaf-feeding beetles	
Boxelder bugs	Leafhoppers	
Clover mites	Mealybugs	
Elm leaf beetles	Pillbugs	
Grasshoppers	Scale (crawlers only)	
Gypsy moth larvae	Spittlebugs	
Japanese beetle (adults)	Thrips	
Lacebug	Whiteflies	

VEGETABLE	PEST	
Carrots Radishes Sweet corn Tomatoes No waiting. You may spray up to day of harvest. Do not spray more than 5 times per season.	Aphids Armyworms Aster leafhoppers Cabbage loopers Colorado potato beetles Corn borers Corn earworms Corn rootworm adults Cutworms	Flea beetles Grasshoppers Leafhoppers Leafminers Stinkbugs Thrips Tomato worms Weevils
Peppers After spraying, wait 7 days before harvesting peppers. Do not spray peppers more than 6 times per season.	Armyworms Cabbage loopers Corn borers Corn earworms Garden webworms	Leafhoppers Leafminers Thrips Weevils
Sunflowers After spraying, wait 30 days before harvesting. Do not spray more than 3 times per season.	Cutworms Grasshoppers Midges	Sunflower beetles Sunflower moths Weevils

FOR BEST RESULTS:
 Do not water the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply.

HOUSEHOLD NUISANCE PESTS OUTDOORS		
PEST		HOW TO APPLY
Ants Boxelder bugs Clover mites Cockroaches Crickets Earwigs Flies	Midges Mosquitoes Pillbugs Sowbugs Spiders Wasps	Spray around doors, window frames and other places where insects may enter house. Spray around foundations and porches. Spray on ant hills and ant trails. (ART)

LAWN PESTS	
PEST	HOW TO APPLY
Ants, Armyworms, Chinchbugs, Crickets, Cutworms, Grasshoppers, Japanese Beetles (Adult)	[For localized infestations,] spray to penetrate the lawn and reach the soil. Do not mow for one day after application. [For large areas, use Tempo® Concentrate Insecticide.]

Children and pets may re-enter the treated area after spray has dried.

IMPORTANT: Directions for Storage and Disposal

Storage Conditions

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

Disposal

- Do not reuse empty spray bottle.
- Rinse the empty product container thoroughly.
- Place container in trash.

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state or local guidelines.

NOTICE

Research and testing have determined that the "Directions for Use" are appropriate for the proper use of TEMPO Ready-to-Use Insecticide under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

