

3125-590

08/30/2002

1/13



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number

3125-590

Date of Issuance

AUG 30 2002

Term of Issuance: Unconditional

Name of Pesticide Product:

Tempo® Ultra WSP

NOTICE OF PESTICIDE:

☒ Registration☐ Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Bayer Corporation, Agriculture Division
8400 Hawthorn Road, P.O. Box 4913
Kansas City, MO 64120-0013

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3).

1. Make the following labeling changes:

- a. Revise the EPA Registration Number to read "EPA Reg. No. 3125-590".
- b. In the table "Application for Turf Pests" on page 8, revise the parenthetical statement after "Ticks" in the left column to read "(including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever)."
- c. Revise the Container disposal instructions to incorporate the recommendations in the revised (draft) Label Review Manual for water soluble packets: "Dispose of the empty outer foil pouch in the trash, as long as WSP is unbroken. If WSP is broken, dispose of empty pouch in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke."

Signature of Approving Official:

Date:

August 30, 2002

Page 2

EPA Reg. No. 3125-590

- d. In the last bullet of the table listing permitted areas of use in the Appendix on page 11, insert "(cargo areas only)" after "aircraft".
 - e. **Delete the statement near the beginning of the Directions for Use "Do not formulate this product into other end-use products." Our regulations do not allow such a restriction of trade. It also conflicts with an applicant's right to apply the formulator's exemption.**
2. Please submit two (2) copies of your final printed label before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this action, please contact Susan Stanton of my team at (703) 305-5218.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505C)

Enclosures

Bayer Professional Care

Tempo® Ultra WSP

Insecticide

Water Soluble Packets

For Pest Management Professionals and commercial use only.

For broad-spectrum control of crawling, flying, and wood-infesting insect pests on indoor and outdoor surfaces, as well as insect pests of trees, landscape ornamentals and residential and commercial lawns.

ACTIVE INGREDIENT:

β-Cyfluthrin,
cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropane-carboxylate..... 10%

OTHER INGREDIENTS:..... 90%
100%

EPA Reg. No. 3125-xxx

Net Wt. 14.1 Ounces
Eight 50 Gram Packets

STOP - Read the label before use.
Keep out of reach of children.

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and warm water after handling. Remove contaminated clothing and wash before reuse.

Wear safety glasses, goggles, or face shield when handling the undiluted material and wear a respirator when making general surface overhead treatments indoors.

Do not allow children or pets to enter treated areas until surfaces are dry.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

AUG 3 D 2002

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

3125-590

FIRST AID

If on skin

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

If inhaled

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. The Bayer Kansas City Emergency Response Telephone No. is 800-414-0244.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.



ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply TEMPO ULTRA WP Insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

PRO FACTS

- Quick Knockdown & Residual Control
- Low Odor
- Non-staining
- Ideal for Power Spray Applications
- No phytotoxicity on a wide range of ornamentals

TEMPO ULTRA WSP	<ul style="list-style-type: none"> • Water soluble packets containing 10% beta-cyfluthrin wettable powder
CONTROLS	<ul style="list-style-type: none"> • Outdoor/ Perimeter Pests • Indoor Pests <ul style="list-style-type: none"> • General Household Pests • Pantry and Stored Product Pests • Food Processing Pests • Wood-infesting Pests • Turf and Ornamental Pests <p><i>For list of specific insects, see inside.</i></p>
WHERE TO APPLY	<p>Apply where pests have been seen or found, or can find shelter.</p> <ul style="list-style-type: none"> • In and around buildings and structures • On residential, commercial and recreational areas of turf • On ornamentals in landscapes and interior plantscapes
COVERAGE	<ul style="list-style-type: none"> • 1 packet treats up to 10,000 sq ft outdoors. • 1 packet makes up to 5 gallons of finished spray solution for indoors.
MIXING INFORMATION	<ul style="list-style-type: none"> • The inner packet contains TEMPO ULTRA WSP insecticide and dissolves in water. • Remove the inner packet and drop into spray tank while adding water.
HOW TO APPLY	<ul style="list-style-type: none"> • General surface spot, mist, or crack & crevice application. • Broadcast application to turf and foliar application to ornamentals.
RE-APPLY	<ul style="list-style-type: none"> • Every 7 to 10 days, if needed. • For turf, reapply if needed, but not more than 6 times per year.
RE-ENTRY 	<ul style="list-style-type: none"> • Avoid contact with treated surfaces until dry. • People and pets may re-enter after the treated area is dry.
QUESTIONS 	<p>For questions or comments, call toll-free 1 800 842-8020 www.bayerprocentral.com</p>

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read entire label and Conditions of Sale before using this product.

Do not formulate this product into other end-use products.

STRUCTURAL PESTS

FOR RESIDUAL PEST CONTROL IN AND AROUND BUILDINGS, STRUCTURES, FOOD/FEED HANDLING ESTABLISHMENTS AND MODES OF TRANSPORT

Use TEMPO ULTRA WSP as a general surface, spot, mist, or crack and crevice treatment in and around buildings and structures. This includes schools, factories, stores, warehouses, hospitals, food processing facilities, and modes of transport. For a list of common facilities where this product may be used, see Appendix.

STRUCTURAL PESTS

OUTDOOR/PERIMETER PESTS

Ants Bees Boxelder bugs Carpenter ants Centipedes Cockroaches Cluster fly Crickets Elm leaf beetle	Fire ants Firebrats Fleas Flies Gnats Ground beetles Hornets Midges Millipedes	Moths Pillbugs Scorpions Silverfish Sowbugs Spiders Ticks Wasps Yellowjackets
--	--	---

WHERE TO APPLY	<ul style="list-style-type: none"> • Apply where pests have been seen or found, or can find shelter. • Apply to buildings, porches, patios, garages, fences and other areas where pests are found. • To prevent infestation, treat the building foundation, walls, and around doors, windows, and soffits where pests are active or may enter or hide; also treat the soil, turf, or other substrates adjacent to buildings
USE RATE	<ul style="list-style-type: none"> • One 50-g packet covers 2,500 sq ft (20 g/1000 sq ft) to 5,000 sq ft (10 g/1000 sq ft). See Table 1 for mixing directions. • For fire ants and flying pests 0.025% —one 50-g packet/ 5 gal water 0.05% — one 50-g packet/ 2.5 gal water • Always use the high rate for fire ants, bees, wasps, yellowjackets and hornets. • Use high rate for severe infestations, faster knockdown, and longer residual.
HOW TO APPLY (Also see APPLICATIONS FOR OUTDOOR/ PERIMETER PESTS)	<ul style="list-style-type: none"> • Perimeter Treatment: To help prevent infestation of structures, treat area adjacent to buildings in a band up to 10 feet wide. Also treat the building foundation to a height of 2 to 3 feet in areas where pests are active or can find entrance. • Apply just enough dilution to adequately cover the area without excessive dripping or runoff. Volume can vary depending on the surface type treated. For example, mulch requires more spray than bare soil. If mulch is heavy, rake it back to expose soil before treating. • Avoid contact with treated surfaces until dry.
WHEN TO APPLY	<ul style="list-style-type: none"> • When pests are seen or found • Re-apply every 7 to 10-days if needed.

STRUCTURAL PESTS (continued)	
MIXING INFORMATION	<ul style="list-style-type: none"> The inner packet contains TEMPO ULTRA WSP insecticide and dissolves in water. To prepare the spray mixture, drop the unopened inner packet into spray tank while adding water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>Note: Do not mix TEMPO ULTRA WSP packets with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre-dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> Diluted spray mixture may be stored overnight. Agitate before using. TEMPO ULTRA WSP can be tank mixed with pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions.

Table 1: Mixing Directions for Outdoor/Perimeter Applications

Application Volume: Gallons/ 1,000 sq ft	Use Rate: Grams/ 1,000 sq ft	Packets of TEMPO ULTRA WSP Diluted to These Volumes of Finished Spray	
		25 gal.	50 gal.
2.5	10	2	4
	20	4	8
5	10	1	2
	20	2	4
10	10	-	1
	20	1	2

Formula for Determining the Percent Concentration in the Finished Spray	
(Packets of TEMPO ULTRA WSP)	Percent Concentration
(Gallons of finished spray) X 7.57	=

APPLICATION FOR OUTDOOR/PERIMETER PESTS	
ANTS	Locate and treat ant nests, if possible. Follow trails of foraging workers, check under objects near the structure, and inspect other likely nest sites. If ants are nesting in the soil, thoroughly drench or inject the nest and surrounding soil. Treat ant trails around windows and doors, along sidewalks, or other areas where seen. Apply a perimeter treatment as described in the HOW TO APPLY box.
BEEES, WASPS, YELLOWJACKETS, HORNETS	Use Rate: 0.05% concentration. Make pin stream applications directly to the nests. To help avoid stings treat nests in the evening when the insects are inside the nest. For best results, check nests within one or two days to ensure complete kill. Remove and destroy treated nests to prevent emergence of newly hatched insects, secondary pest problems, and possible staining from honey. If removal is not possible, retreat nests if necessary.
BOXELDER BUGS	Apply directly to boxelder bugs that have congregated around doors and windows and similar areas where they may enter structures. Apply perimeter treatment as described in the HOW TO APPLY box. For best results, treat infested trees early in the season. Tree spraying may require ornamental license – check with your local regulator.
CENTIPEDES, MILLIPEDES, PILLBUGS, SOWBUGS	Apply perimeter treatment to reduce the number of insects from entering the structure. Thoroughly treat mulched areas. Reduce moisture areas from around the structure.
FIRE ANTS	Use Rate: 0.05% concentration Thoroughly drench or inject the entire mound. When using injection treatment, you must treat the bottom of the mound. Apply dilution or foam using a downward and upward motion of the injector tool. For complete distribution, inject in several points throughout the mound. Treat new mounds as they appear. Do not use in pastures or areas of food crop production.
FLEAS & TICKS	Make a perimeter treatment, concentrating on areas where pets rest or run. For ticks, treat likely harborage areas like along paths, tall grass and other vegetation. For best results, treat the entire area where ticks could be present. Begin treating in the spring. Ticks may be re-introduced from surrounding areas by host animals. Continue treating until frost to control both larvae and adult ticks.

APPLICATION FOR OUTDOOR/PERIMETER PESTS (continued)	
FLIES	<p>Use Rate: 0.025% to 0.05%</p> <p>Apply low-volume mist or spray to outside surfaces of buildings, porches, patios, garages, fences, and other areas where flies rest.</p> <p>CLUSTER FLY: Treat in late summer or fall, just before flies begin looking for winter shelter. Spray siding, eaves, around windows and doors, window screens, and attics. To help reduce infestations, treat cracks and crevices, attics and other indoor winter hiding places.</p> <p>Note: Cluster flies may appear during winter warm spells, while snow is still on the ground.</p>
MOSQUITOES	Spray around landscape plants, turf and ground cover, under decks, around building foundations and other places where mosquitoes may rest.
SCORPIONS	Treat around piles of firewood and other areas where scorpions may hide. When possible, remove piles of materials that can harbor the pests. Apply a perimeter treatment.

STRUCTURAL PESTS		
INDOOR PESTS		
General Household Pests	Pantry & Stored Product Pests	Food Processing Pests
<u>Crawling Pest</u> Ants (except Pharaoh) Bedbugs Carpet beetles Centipedes Clothes moths Cluster fly Cockroaches Crickets Darkling beetles Earwigs Firebrats Millipedes Pillbugs Silverfish Sowbugs Spiders <u>Flying Pests</u> Flies Gnats Hornets Midges Moths Wasps Yellowjackets	<u>Beetles</u> (exposed adults & immature stages) Cadelle Cigarette Confused flour Dermestid Drugstore Granary weevil Hide Larder Leather Lesser grain borer Lesser mealworm Merchant grain Mealworm Red flour Rice weevil Sawtoothed grain Warehouse <u>Moths</u> Indian meal Mediterranean flour	Fruit flies Phorid flies Sciariid flies Also includes: General Household Pests Pantry & Stored Product Pests
WHERE TO APPLY	Apply where pests have been seen or found, or can find shelter. Use a general surface, spot, mist, or crack and crevice application.	
USE RATE	0.025% —one 50 g packet/5 gal water 0.05% — one 50 g packet/2.5 gal water • Use high rate for severe infestations, faster knockdown, and longer residual. • Always use the high rate for bees, wasps, yellowjackets and hornets.	

INDOOR PESTS (continued)	
HOW TO APPLY (Also see APPLICATION FOR INDOOR PESTS)	<ul style="list-style-type: none"> Remove or tightly cover fish tanks and disconnect aerators during application. Remove pets during application and until treated surfaces are completely dry. Avoid contact with treated surfaces until dry. Wear safety glasses, goggles, or face shield and a dust/mist respirator when treating overhead areas. <p><u>General Surface Applications</u></p> <ul style="list-style-type: none"> Use a low-pressure system (do not exceed 50 psi at nozzle tip), with a fan-type nozzle to apply the dilution uniformly. In the home, all food processing surfaces and utensils should be covered or thoroughly washed following treatment. Cover exposed food or remove from area being treated. <p><u>Spot, Mist, or Crack and Crevice Applications</u></p> <ul style="list-style-type: none"> Apply directly into voids, cracks and crevices using spray or foam equipment. Spray: Use a low-pressure system (do not exceed 50 psi at nozzle tip), with a pinpoint or variable pattern nozzle or applicator equipment that delivers low volume treatments, such as the Micro-Injector® or Actisol®. Foam: Use appropriate foam generating equipment to supply a sufficient volume of foam. Mix the dilution with manufacturer's recommended volume of foaming agent.
WHEN TO APPLY	<ul style="list-style-type: none"> When pests are seen or found Re-apply every 7 to 10-days if needed.
MIXING INFORMATION	<ul style="list-style-type: none"> The inner packet contains TEMPO ULTRA WSP insecticide and dissolves in water. To prepare the spray mixture, drop the unopened inner packet into spray tank while adding water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>Note: Do not mix TEMPO ULTRA WSP packets with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre-dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> Diluted spray mixture may be stored overnight. Agitate before using. TEMPO ULTRA WSP can be tank mixed with pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions.

RESTRICTIONS	<ul style="list-style-type: none"> Do not treat entire area of floor or floor coverings. Do not apply where electrical short circuits can occur. Use a dust or dry bait in these areas. Do not apply in occupied hospital and/or nursing home rooms. Occupants may return when surfaces have dried. Do not apply in occupied classrooms. Do not apply to furniture or upholstery where prolonged contact by humans will occur. Do not apply as a space spray.
---------------------	---

ADDITIONAL DIRECTIONS FOR FOOD AND FEED HANDLING ESTABLISHMENTS	
WHERE TO APPLY	<p>Applications may be made to both Food/Feed and non-food areas of food and feed handling establishments.</p> <p>Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared, or served. Includes areas for:</p> <ul style="list-style-type: none"> receiving, storing and packing (canning, bottling, wrapping, boxing) food preparing food edible waste storage enclosed processing systems (mills, dairies, edible oils, syrups) of food. <p>Serving areas where food is exposed and the facility is in operation are also considered food areas.</p>
USE RATE	<ul style="list-style-type: none"> Apply as a general surface spot, mist, or crack & crevice application. 0.025% —one 50 g packet/5 gal water 0.05% — one 50 g packet/2.5 gal water Use high rate for severe infestations, faster knockdown, and longer residual.
HOW TO APPLY (Also see INDOOR PESTS)	<p><u>General Surface Applications</u></p> <p>Cover or remove all food processing and handling equipment before treatment. After applying where food is commercially prepared or processed, wash all handling and processing equipment, benches, shelving, and other surfaces which food will contact. Rinse thoroughly with clean, fresh water.</p> <p>Do not apply to surfaces in food-handling areas when facility is in use or when foods are exposed.</p> <p>Do not apply directly to food.</p> <p><u>Spot or Crack & Crevice Applications</u></p> <p>Spot or crack and crevice application is permitted while facility is in use. Cover or remove food from area being treated.</p> <p>Do not apply directly to food or food handling surfaces.</p>

APPLICATION FOR INDOOR PESTS	
GENERAL HOUSEHOLD PESTS Cockroaches, Crickets, Silverfish, Spiders, and others	<ul style="list-style-type: none"> Use a coarse, low-pressure spray. Treat entry points such as around water pipes, doors, windows, and eaves. Treat areas where pests normally feed or hide such as baseboards, corners, around water pipes, under refrigerators, cabinets, sinks, and stoves, being sure to spray cracks & crevices. Spot treat floor or rugs beneath furniture, in closets, and storage areas, but do not apply to entire floor area. Do not use as a "space spray". IGR Note: This product can be tank mixed with an insect growth regulator (IGR) for extended cockroach control.
ANTS	<ul style="list-style-type: none"> Apply to ant trails on walls, tile, baseboards, around pipes, under cabinets, and other places where ants are active. Treat entry points such as windows and doors and the outdoor perimeter. For best results, treat nests when possible.
PANTRY AND STORED PRODUCT PESTS	<ul style="list-style-type: none"> For best results, clean area before treating. Remove and destroy infested food. Discard food accidentally contaminated with spray. For cupboards, shelving, or food storage areas: Remove utensils, exposed food, shelf paper, and other items before spraying. When dried, cover shelves with clean paper before replacing items.
COMMERCIAL STORED PRODUCT PESTS	<ul style="list-style-type: none"> Commercial storage or transportation: Clean and treat warehouses, production facilities, storage areas, rail cars, truck beds, and other areas where products are stored <i>before</i> filling with the product.
PESTS IN AIRCRAFT	<ul style="list-style-type: none"> Do not use in aircraft cabins. May be applied to carpet or upholstery in the cargo area only where there is no prolonged human contact.
PESTS IN LIVESTOCK HOUSING STRUCTURES including poultry houses & pet kennels	<ul style="list-style-type: none"> Apply as a general surface and/or a crack and crevice spray. For best results, clean facilities before applying and treat outdoor perimeter areas. Do not apply indoors where animals other than cattle or horses are present. Do not apply to animal feedstuffs or watering equipment.
FLYING INSECTS	<ul style="list-style-type: none"> Apply to surfaces where pests collect or rest.

STRUCTURAL PESTS		
WOOD-INFESTING PESTS		
Beetles Ambrosia Deathwatch False powderpost Old house borer Powderpost	Carpenter ants Carpenter bees	Termites Subterranean Formosan Drywood
WHERE TO APPLY	<ul style="list-style-type: none"> Apply where pests have been seen or found, or can find shelter. 	
USE RATE	<ul style="list-style-type: none"> 0.05% – one 50-g packet/2.5 gal water 	
HOW TO APPLY (Also see Application section)	<ul style="list-style-type: none"> Apply spray or foam as described in application section. Avoid contact with treated surfaces until dry. 	
WHEN TO APPLY	<ul style="list-style-type: none"> When pests are seen or found Re-apply every 7 to 10-days, if needed. 	
MIXING INFORMATION	<ul style="list-style-type: none"> The inner packet contains TEMPO ULTRA WSP insecticide and dissolves in water. To prepare the spray mixture, drop the unopened packet into the spray tank while filling with water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. Note: Do not mix TEMPO ULTRA WSP packets with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre-dissolve packet(s) in a separate container of water and then pour mixture into the spray tank. Diluted spray mixture may be stored overnight. Agitate before using. TEMPO ULTRA WSP can be tank mixed with pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions. Foam applications: Mix End-use dilution with manufacturer's recommended volume of foaming agent for 0.05% concentration. 	

APPLICATION FOR WOOD-INFESTING PESTS	
CARPENTER ANTS	<p><u>Around houses and other structures:</u> Apply around doors and windows and other places where carpenter ants may enter. Spray or foam into cracks & crevices, or spray, mist or foam through small drilled holes into voids where ants or nests are present. Use enough spray or foam to cover the area. Do not exceed 1 gallon of dilution per 1000 sq ft.</p> <p>Inside trees, utility poles, fencing and decking materials and similar structural members: Drill to find the interior infested cavity and inject dilution or foam using an appropriate treatment tool with a splashback guard.</p> <p><u>Tunneling in soil:</u> Apply as a soil drench or inject the dilution or foam in the soil at intervals of 8 to 12 inches. Use 1 gallon of dilution per 8 sq ft. Establish a uniform vertical barrier at the edge of walls, driveways or other hard surfaces where ants are tunneling beneath the surfaces.</p> <p><u>Protecting firewood from carpenter ants:</u> Treat soil beneath firewood before stacking at 1 gallon of dilution per 8 sq ft. Do not treat firewood with this product.</p>
(Continued on next column)	

APPLICATION FOR WOOD-INFESTING PESTS (continued)	
CARPENTER BEES	<p><u>In structures:</u> Spray or foam directly into gallery entrance holes. Leave entrance open for 24 hours to be sure that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant.</p>
WOOD-INFESTING BEETLES	<p>On surfaces in homes, apartments, and other structures</p> <p>For small areas, brush or spray evenly on wood surfaces.</p> <p>For large or overhead areas, thoroughly cover wood surfaces with a coarse spray.</p> <p>Protect surfaces below the treated area from accidental dripping by covering with disposable plastic sheeting or other material that can be thrown away after treatment.</p> <p><u>In localized areas:</u> Apply dilution or foam to voids and galleries in damaged wood, in spaces between wooden structural members, and between the sill plate and foundation. Apply to inaccessible areas by drilling, and then injecting, spraying, or misting the dilution or foam into the damaged wood or wall voids.</p>
DRYWOOD AND ABOVE-GROUND SUBTERRANEAN TERMITES	<p>This type of application is intended to be a supplemental treatment for control of drywood and above-ground subterranean termites and other wood-infesting insects.</p> <p><u>In localized areas:</u> Apply dilution or foam to voids and galleries in damaged wood, to spaces between wooden structural members, and between the sill plate and foundation. Apply to inaccessible areas by drilling, and then injecting, spraying, or misting the dilution or foam into damaged wood or wall voids.</p> <p><u>Swarming or reproductive stages:</u> Apply dilution to the swarm or area on which the termites collect. Make applications to attics, crawl spaces, unfinished basements or man-made voids with a coarse fan spray to control exposed reproductive forms of termites.</p> <p>Above-ground termites inside trees, utility poles, fencing and decking materials and similar structural members: Drill to find the infested cavity and inject dilution or foam using an appropriate tool with a splashback guard.</p> <p><u>Termite carton nests:</u> Inject dilution or foam into building voids or tree cavities using a pointed injection tool. Several injection points to varying depths may be needed. For best results, remove carton nest material when possible.</p> <p><u>Protecting firewood from termites:</u> Treat soil beneath firewood before to stacking at 1 gallon of dilution per 8 square feet. Do not treat firewood with this product.</p>

TURF PESTS	
TEMPO ULTRA WSP controls designated pests on landscape and recreational turf.	
TURF PESTS	USE RATE
Ants Armyworms Crickets Cutworms Earwigs Grasshoppers Japanese beetles (adult) Sod webworms Ticks (including deer ticks)	50 grams (1 packet) per 5,000 to 10,000 sq. ft. Use high rate for severe infestations, faster knockdown, and longer residual.
Black turfgrass ataenius (adult) Bluegrass billbug (adult) Chinch bugs Chiggers Fire ants (aids in control) Fleas Hyperodes weevil (adult) Mole crickets Scorpions	50 grams (1 packet) per 5,000 sq. ft.
WHERE TO APPLY	<ul style="list-style-type: none"> For use on landscape and recreational turfgrass. Not for use on turf grown for research, commercial seed production, sale, or other commercial use. Not for use on golf course turf. Apply where pests have been seen or found, or can find shelter. Permitted areas include but are not limited to: home lawns, business complexes, apartment buildings, airports, cemeteries, parks, playgrounds, recreational fields, institutional grounds, athletic fields, and other landscape turfgrass.
HOW TO APPLY (Also see APPLICATION FOR TURF PESTS)	<ul style="list-style-type: none"> Use an accurately calibrated sprayer. Apply uniformly using spray nozzles with a coarse droplet size such as flood jet nozzles. Keep children and pets off treated areas until lawn has dried.
WHEN TO APPLY	Apply when pests first appear. Reapply as needed based on pest reinfestation. Do not treat more than 6 times per year. Consult your state Cooperative Extension Office for the specific timing in your area.
(continued on next page)	

TURF PESTS (Continued)	
MIXING INFORMATION	<ul style="list-style-type: none"> The inner packet contains TEMPO ULTRA WSP insecticide and dissolves in water. To prepare the spray mixture, drop the unopened packet into the spray tank while filling with water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>Note: Do not mix TEMPO ULTRA WSP packets with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre-dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> Diluted spray mixture may be stored overnight. Agitate before using. TEMPO ULTRA WSP can be tank mixed with pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions. COMPATABILITY: Compatible with commonly used fungicides, miticides, liquid fertilizers and other insecticides.
RESTRICTIONS	<ul style="list-style-type: none"> Do not apply this product by aerial application or by any type of chemigation system. Do not apply when the average wind speed is greater than 15 mph. Do not allow drift onto ponds, streams, or lakes. Do not apply when turfgrass areas are waterlogged or the soil is saturated with water (i.e., will not accept irrigation).

APPLICATION FOR TURF PESTS	
ARMYWORMS, CUTWORMS, SOD WEBWORMS	Do not water or mow for at least 24 hours following application.
BILLBUGS	Apply when adults first appear in the spring. In temperate climates, this application will also control overwintering chinch bugs.
CHINCH BUGS	For best results, water the turf immediately after application to aid in the penetration of the active ingredient into the thatch layer where chinch bugs dwell.
TICKS (Including ticks that transmit Lyme Disease and Rocky Mountain Spotted Fever)	For best results, treat the entire area where ticks could be present. Begin treating in the spring. Ticks may be re-introduced from surrounding areas by host animals. Continue treating until frost to control both larvae and adult ticks.

ORNAMENTAL PESTS		
TEMPO ULTRA WSP controls designated pests on trees, shrubs, foliage plants and flowers in outdoor landscaped areas and interior plantscapes where these plants are grown.		
Ants Aphids Armyworms Azalea caterpillars Bagworms Black vine weevil (adult) Boxelder bugs Bristly rose slugs Budworms California oakworms Cankerworms Casebearers Clover mites Cockroaches Crickets Cutworms Elm leaf beetles Elm spanworms Flea beetles Flies Fungus gnats	Grasshoppers Gypsy moth larvae Japanese beetles (adult) June beetles (adult) Lace bugs Leaf-feeding caterpillars Leafhoppers Leafrollers Leaf skeletonizers Mealybugs Midges Mosquitoes Oleander moth larvae Orchid weevil Pear psylla Peppertree psyllid Pillbugs Pine shoot moths Pine tip moths Plant bugs	Redhumped caterpillars Sawfly larvae Scale insects (crawler stages) Sowbugs Spiders Spittlebugs Striped cucumber beetles Striped oakworms Tent caterpillars Thrips Ticks (including deer ticks) Tussock moth larvae Walnut caterpillars Wasps Webworms Whiteflies Yellownecked caterpillars
WHERE TO APPLY	<ul style="list-style-type: none"> Use in landscapes and interiorscapes. Permitted areas include but are not limited to: parks, recreational areas, athletic fields, institutional grounds, hotels, malls, office buildings, and airports. Not for use on plants or turf grown for research, commercial seed production, sale, or other commercial use. Not for use in greenhouses or nurseries. Apply where pests are seen, have been found, or can find shelter among trees, shrubs, evergreens, ground covers, flowers, and foliage plants. No phytotoxicity has been observed on a wide range of ornamentals tested under various environmental conditions. If in doubt, first try this product on a small sample. 	
USE RATE	<ul style="list-style-type: none"> 1 to 3 packets TEMPO ULTRA WSP per 100 gallons of water Use high rate for severe infestations, faster knockdown, and longer residual. 	
HOW TO APPLY (Also see APPLICATION FOR ORNAMENTAL PESTS)	<p>Apply as a dilute spray just to the point of run-off but do not allow excess run-off. For hard-to-wet foliage such as holly, pine, or ivy, adding a spreader-sticker may enhance knockdown and residual activity. For best results, spray thoroughly for good coverage including the underside of leaves for pests that feed or rest there.</p> <ul style="list-style-type: none"> Spray plants in bloom at times when pollinating insects are not present, such as early morning or late evening. Remove any animal feeding dishes prior to treatment. <p>Avoid contact with treated surfaces until dry.</p>	

(continued on next column)

ORNAMENTAL PESTS (continued)	
WHEN TO APPLY	<ul style="list-style-type: none"> Apply when pests first appear. Reapply as needed based on pest infestation. Consult your state Cooperative Extension Office for the specific timing in your area.
MIXING INFORMATION	<ul style="list-style-type: none"> The inner packet contains TEMPO ULTRA WSP insecticide and dissolves in water. To prepare the spray mixture, drop the unopened packet into the spray tank while filling with water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>Note: Do not mix TEMPO ULTRA WSP packets with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre-dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> Diluted spray mixture may be stored overnight. Agitate before using. TEMPO ULTRA WSP can be tank mixed with pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions. COMPATABILITY: Compatible with commonly used fungicides, miticides, liquid fertilizers and other insecticides.
RESTRICTIONS	<ul style="list-style-type: none"> Do not apply to any food crop. Do not allow drift onto ponds streams or lakes.

APPLICATION FOR ORNAMENTAL PESTS	
ANTS, CRICKETS, SPIDERS, MIDGES, WASPS AND FLIES	Apply as a general spray, concentrating on damp areas beneath shrubbery, around foundations and in areas of tall grass or weeds in ornamental areas and areas adjacent to buildings.
BAGWORMS	Spray when bagworms first begin to hatch. If possible, apply directly to the larvae. Young larvae are easier to control.
BLACK VINE WEEVIL (adult)	<p>Use rate: 2 – 3 packets per 100 gallons of water.</p> <p>Apply as needed when adults and damage are first noticed in spring and at 3 to 4 week intervals while weevils and fresh leaf notching are observed. Addition of a spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having difficult to wet foliage.</p>
SCALE (CRAWLERS)	Spray thoroughly, including foliage, trunk, limbs, and twigs.

STORAGE AND DISPOSAL	
Do not contaminate water, food, or feed by storage or disposal.	
Storage:	<ul style="list-style-type: none"> • Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. • Protect from freezing. • Store in original containers and out of the reach of children, preferably in a locked storage area. • Handle and open container in a manner as to prevent spillage.
Pesticide Disposal:	<ul style="list-style-type: none"> • Pesticide wastes are hazardous. • Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. • If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
Container Disposal:	<ul style="list-style-type: none"> • Do not use container in connection with food, feed, or drinking water. • Completely empty container into application equipment. Triple rinse, or equivalent. • Dispose of container by offering for recycling or reconditioning. Or puncture and dispose of container in a sanitary landfill, or by incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.
Spills:	<p>If the container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material.</p> <ul style="list-style-type: none"> • Do not walk through spilled material. • Dispose of as directed for pesticides above. • Keep unauthorized people away. • You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance at 800-414-0244 or contact Chemtrec at 800-424-9300.

CONDITIONS OF SALE. Read Before Using.
<ul style="list-style-type: none"> • THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. • THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. • INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. • IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

APPENDIX

Use TEMPO ULTRA WSP as a general surface, spot, mist, or crack and crevice treatment.

Permitted areas of use in buildings and structures, and modes of transport include, but are not limited to:

- apartment buildings
- bakeries
- bottling facilities
- breweries
- buses
- cafeterias
- candy plants
- cereal processing and manufacturing plants
- correctional facilities,
- dairies and dairy product processing plants
- frozen food plants
- grain mills
- granaries
- greenhouses (structures only) (non commercial)
- hospitals
- hotels
- houses
- industrial buildings
- kitchens
- laboratories
- livestock housing
- pet kennels
- manufacturing establishments
- mausoleums
- meat and produce canneries
- nursing homes
- pickle factories
- poultry houses
- processing and packing plants
- restaurants
- schools
- spice plants
- stores
- warehouses
- wineries and similar structures
- transportation vessels such as aircraft, buses, trucks, trailers, rail cars, and marine vessels

D - 9701 052102

Bayer Corporation
Professional Care
Box 4913
Kansas City, MO 64120-0013
(800) 842-8020
www.bayerprocentral.com

