

This product contains lindane, the gamma isomer of benzene hexachloride in an approved carrier

CAUTION

Avoid inhalation. May cause irritation of nose and throat. May be absorbed through skin. Contact with skin and eyes may cause irritation. Wash with plenty of soap and water. For eyes get medical attention — flush with water for 15 minutes. Avoid contamination of food and feed stuffs

Avoid dusting feed troughs or utensils. Do not treat sows, within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated and sick animals.

NOTICE

VERY YOUNG ANIMALS may be injured by this treatment. Do not treat animals under 3 months of age. Treat very lightly and with caution. Do not apply within 30 days of slaughter. Do not use on cats or dairy animals. Do not apply to nursing mothers or nursing puppies.

NET WEIGHT 1 LB.
(16 ounces)

VITA PLUS CORPORATION
GENERAL OFFICES
MADISON, WISCONSIN 53713

TAKE TIME



OBSERVE LABEL
DIRECTIONS

© 1971 Vita Plus Corporation

471

VITA PLUS

MANGE DRY TREATMENT

3213-8

Sept 15 1972

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECG... IN REGISTERED UNDER...

FOR HOGS, BEEF CATTLE, SHEEP, HORSES & DOGS

ACTIVE INGREDIENTS:
Gamma isomer of benzene hexachloride from lindane 1%

INERT INGREDIENTS 99%

CAUTION
Out of Reach
See other cautions on...

DIRECTIONS

Mange Dry Treatment is recommended for the treatment of mange (Psoroptic and Sarcoptic), lice, ticks, and fleas on hogs, beef cattle, sheep, horses, and dogs.

Apply dust liberally over all parts of the animal's body using dust gun if available.

Dust bedding and walls of sleeping quarters for additional control. Repeat treatment as needed, but do not apply more often than once a week. Do not use inside the household.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Empty bags should be buried or burned. (Stay away from smoke!) If the bags are burned, the ashes should be buried in a safe place.

This product contains lindane, the gamma isomer of benzene hexachloride in an approved carrier.

CAUTION

Avoid inhalation. May cause irritation of nose and throat. May be absorbed through skin. Contact with skin and eyes may cause irritation. Wash with plenty of soap and water. For eyes get medical attention — flush with water for 15 minutes. Avoid contamination of food and feed stuffs.

Avoid dusting feed troughs or utensils. Do not treat sows, within 2 weeks of farrowing nor far at least 3 weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated and sick animals.

NOTICE

VERY YOUNG ANIMALS may be injured by this treatment. Do not treat animals under 3 months of age. Treat very lightly and with caution. Do not apply within 30 days of slaughter. Do not use on cats or dairy animals. Do not apply to nursing mothers or treat nursing ones.

NET WEIGHT 1 LB.
(16 ounces)

VITA PLUS CORPORATION
GENERAL OFFICES
MADISON, WISCONSIN 53713

TAKE TIME



OBSERVE LABEL DIRECTIONS



3213-8

REGISTERED
Sept 15 1972
UNDER THE FEDERAL INSECTICIDE ACT

**FOR HOGS, BEEF CATTLE,
SHEEP, HORSES & DOGS**

ACTIVE INGREDIENTS.

Gamma isomer of benzene
Hexachloride from lindane 1%
INERT INGREDIENTS 99%

CAUTION

Keep Out of Reach of Children

See other cautions on back panel

DIRECTIONS

Mange Dry Treatment is recommended for the treatment of mange (Psoroptic and Sarcoptic), lice, ticks and fleas on hogs, beef cattle, sheep, horses, and dogs.

Apply dust liberally over all parts of the animal's body using dust gun if available.

Dust bedding and walls of sleeping quarters for additional control. Repeat treatment as needed, but do not apply more often than once a week. Do not use inside the household.

Do not use on empty animal enclosures. Destroy by burying with waste or burning slowly with the animal's carcass.

Do not use on cattle being treated for fly away. Do not use on the head or horns of any animal to be treated.