

Calc-N-Rich/Rabon®

EPA Reg. No. 3213-37

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:

*2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate	0.60%
INERT INGREDIENTS**	99.40%
Total	100.00%

*Each pound contains 2.7 g. Rabon®

**Refers only to ingredients which are not larvicidal.

EPA REG. NO. 3213-37

EPA EST. NO. 3213-WI-1

GUARANTEED ANALYSIS:

Calcium (Ca.) (Min.)	19.0%	Potassium (K.) (Min.)	1.5%
Calcium (Ca.) (Max.)	21.6%	Sulfur (S.) (Min.)	2.5%
Phosphorus (P.) (Min.)	10.0%	Iodine (I.) (Min.)	0.007%
Magnesium (Mg.) (Min.)	2.8%	Selenium (Se.) (Min.)	0.0007%

EACH POUND CONTAINS NOT LESS THAN:

Vitamin A	300,000 U.S.P. Units	Vitamin E	100 Int. Units
Vitamin D3	100,000 U.S.P. Units		

FEED INGREDIENTS:

Monocalcium and Dicalcium Phosphate, Calcium Carbonate, Magnesium Sulfate, Potassium Sulfate, Magnesium Oxide, Zinc Oxide, Manganous Oxide, Sulfur, Copper Oxide, Iron Carbonate, Ethylenediamine Dihydriodide, Cobalt Carbonate, Sodium Selenite, Corn Distillers Dried Grains with Solubles, Condensed Fermented Corn Extractives, Cane Molasses, Bentonite, Mineral Oil, Brewers Dried Yeast, Vitamin A Supplement, D-Activated Animal Sterol (Source of Vitamin D3), Vitamin E Supplement, Niacin, Choline Chloride, Calcium Pantothenate, Riboflavin Supplement, Menadione Sodium Bisulfite Complex (Source of Vitamin K), Artificial Flavor.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place.

PESTICIDE DISPOSAL — Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL — Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE:

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Start feeding Vita Plus Calc-N-Rich/Rabon® early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Vita Plus Calc-N-Rich/Rabon® may be fed up to slaughter and to lactating dairy cows.

Vita Plus Calc-N-Rich/Rabon® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control Vita Plus Calc-N-Rich/Rabon® should be used in conjunction with other good management and sanitation practices.

MANUFACTURED BY:

VITA PLUS CORPORATION

GENERAL OFFICES
Madison, Wisconsin 53713

NET WEIGHT 50 LBS.

PC # 1061

FEEDING DIRECTIONS

Feed Vita Plus Calc-N-Rich/Rabon* at the rate of 5 ounces per 1200 pounds of body weight daily

The following table shows the correct amount of Vita Plus Calc-N-Rich/Rabon* to be mixed into a complete daily ration in order to provide the recommended average daily intake of 70 mg. Rabon* Oral Larvicide per hundred pounds of body weight.

Determine your cattle's average body weight, find this figure in the left column. Then determine the average consumption of your complete daily ration and find this figure in the top column. Cross reference and you will find amount of Vita Plus Calc-N-Rich/Rabon* to be mixed into your ration. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if Vita Plus Calc-N-Rich/Rabon* is dispersed uniformly with the feedstuff at the recommended level. Common feeding mixing equipment (i.e. vertical mixers, horizontal blender, mixer/feeder trucks) may be used to prepare formulated feed.

CONSUMPTION PER HEAD PER DAY OF FINISHED FEED (Pounds)

	Lbs.	4#	5#	6#	7#	8#	9#	10#	15#	20#	25#	30#	
AVERAGE WEIGHT OF ANIMALS (Pounds)	400	52	41	35	30	26	23	21	14	10	8	7	POUNDS OF VITA PLUS CALC-N- RICH/ RABON* TO BE MIXED PER TON OF FINISHED FEED
	500	65	52	43	37	32	29	26	17	13	10	9	
	600	78	62	52	44	39	35	31	21	16	12	10	
	700	91	73	61	52	45	40	36	24	18	15	12	
	800	104	83	69	59	52	46	41	28	21	17	14	
	900	117	93	78	67	58	52	47	31	23	19	16	
	1000	130	104	86	74	65	58	52	35	26	21	17	
	1100	143	114	95	81	71	63	57	38	29	23	19	
	1200	156	124	104	89	78	69	62	41	31	25	21	
	1300	169	135	112	96	84	75	67	45	34	27	22	
	1400	181	145	121	104	91	81	73	48	36	29	24	
	1500	194	156	130	111	97	86	78	52	39	31	26	
	1600	207	166	138	119	104	92	83	55	41	33	28	
	1700	220	176	147	126	110	98	88	59	44	35	29	
1800	233	187	156	133	117	104	93	62	47	37	31		
1900	246	197	164	141	123	109	99	66	49	39	33		
2000	259	207	173	148	130	115	104	69	52	41	35		

* Rabon is the registered trademark of Shell Oil Co. U.S. Pat. 3 102 842