



ACCEPTED
SEP 4 1984
32,3-38

Phos-N-Rich/Rabon®

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:

*2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate 1.00%

INERT INGREDIENTS**

Total 99.00%
100.00%

*Each pound contains 4.54 g. Rabon®
**Refers only to ingredients which are not larvicidal
EPA REG. NO. 3213-38
EPA EST. NO. 3213-WI-1

GUARANTEED ANALYSIS:

Calcium (Ca.)(Min.)	0.2%	Potassium (K.)(Min.)	1.0%
Calcium (Ca.)(Max.)	1.0%	Sulfur (S.)(Min.)	2.0%
Phosphorus (P.)(Min.)	20.0%	Iodine (I.)(Min.)	0.012%
Magnesium (Mg.)(Min.)	2.0%	Selenium (Se.)(Min.)	0.0012%

EACH POUND CONTAINS LESS THAN:

Vitamin A	500,000 U.S.P. Units	Vitamin E	175 Int. Units
Vitamin D3	175,000 U.S.P. Units		

FEED INGREDIENTS:

Mono-ammonium Phosphate, Magnesium Oxide, Magnesium Sulfate, Potassium Sulfate, Zinc Oxide, Manganese Oxide, Sulfur, Copper Oxide, Iron Carbonate, Ethylenediamine Dihydriodide, Cobalt Carbonate, Sodium Selenite, Corn Distillers Dried grains with Solubles, Condensed Fermented Corn Extractives, Molasses Products, Bentonite, Mineral Oil, Brewers Dried Yeast, Vitamin A Supplement, D-activated Animal Sterol (Source of Vitamin D3), Vitamin E Supplement, Niacin, Choline Chloride, Calcium Pantothenate, Riboflavin Supplement, Menadione Sodium Bisulfite Complex (Source of Vitamin K), Artificial Flavor.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling, and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a dry place.
PESTICIDE DISPOSAL — Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL — Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE: GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Start feeding Vita Plus Phos-N-Rich/Rabon® early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.
Vita Plus Phos-N-Rich/Rabon® may be fed up to slaughter and to lactating dairy cows.
Vita Plus Phos-N-Rich/Rabon® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.
Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.
In order to achieve optimum fly control Vita Plus Phos-N-Rich/Rabon® should be used in conjunction with other good management and sanitation practices

MANUFACTURED BY:
VITA PLUS CORPORATION
GENERAL OFFICES
Madison, Wisconsin 53713
NET WEIGHT 50 LBS

FEEDING DIRECTIONS

Feed Vita Plus Phos-N-Rich/Rabon* at the rate of 3.0 ounces per 1200 pounds of body weight daily.

The following table shows the correct amount of Vita Plus Phos-N-Rich/Rabon* to be mixed into a complete daily ration in order to provide the recommended average daily intake of 70 mg. Rabon* Oral Larvicide per hundred pounds of body weight.

Determine your cattle's average body weight, find this figure in the left column. Then determine the average consumption of your complete daily ration and find this figure in the top column. Cross reference and you will find amount of Vita Plus Phos-N-Rich/Rabon* to be mixed into your ration. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if Vita Plus Phos-N-Rich/Rabon* is dispersed uniformly with the feedstuff at the recommended level. Common feeding mixing equipment (i.e. vertical mixers, horizontal blender, mixer/feeder trucks) may be used to prepare formulated feed.

CONSUMPTION PER HEAD PER DAY OF FINISHED FEED (Pounds)

	Lbs.	4#	5#	6#	7#	8#	9#	10#	15#	20#	25#	30#	
AVERAGE WEIGHT OF ANIMALS (Pounds)	400	31	25	21	18	16	14	12	8	6	5	4	POUNDS OF VITA PLUS PHOS-N- RICH/ RABON TO BE MIXED PER TON OF FINISHED FEED
	500	39	31	26	22	19	17	15	10	8	6	5	
	600	46	37	31	26	23	21	19	12	9	7	6	
	700	54	43	36	31	27	24	22	14	11	9	7	
	800	62	49	41	35	31	27	25	17	12	10	8	
	900	69	56	46	40	35	31	28	19	14	11	9	
	1000	77	62	51	44	39	34	31	21	15	12	10	
	1100	85	68	57	48	42	38	34	23	17	14	11	
	1200	92	74	62	53	46	41	37	25	19	15	12	
	1300	100	80	67	57	50	45	40	27	20	16	13	
	1400	108	86	72	62	54	48	43	29	22	17	14	
	1500	116	93	77	66	58	51	46	31	23	19	15	
	1600	123	99	82	71	62	55	49	33	25	20	16	
	1700	131	105	87	75	66	58	52	35	26	21	17	
	1800	139	111	93	79	69	62	56	37	28	22	18	
	1900	146	117	98	84	73	65	59	39	29	23	19	
2000	154	123	103	88	77	68	62	41	31	25	20		

*Rabon is the registered trademark of Shell Oil Co. U.S. Pat. 3 102 842