

NEW
APPLY ANY
TIME



\$998


Net Wt. 16 lbs.

Treats 2,000 sq. ft.

200-12
ACCEP
file 1/1/61

total
for lawns

1. Prevents Crabgrass
2. Controls Living Crabgrass
3. Kills Broadleaf Weeds
4. Controls Many Lawn Insects (See Rear Panel for Specific Insects)
5. 10-6-4 Lawn Fertilizer
6. Contains Sulfur, Iron, Zinc, and Manganese



The most complete lawn product made



ACTIVE INGREDIENTS: N-oleyl 1,3-propylenediamine salt of 2,4-dichlorophenoxyacetic acid* 0.436%, dimethylamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid** 0.489%, dimethyl ester of tetrachloroterephthalic acid*** 2.87%, disodium methanearsonate hexahydrate*** 1.58%, Heptachloroepoxyhydro-4,7-methanoindene 0.374% plus 0.212% related compounds.

INERT INGREDIENTS: 93.837%

*Equivalent to 0.307% 2,4-dichlorophenoxyacetic acid, use as a herbicide, patented under U.S. Patent No. 2,900,411. **Equivalent to 0.405% of 2-(2-methyl-4-chlorophenoxy) propionic acid. ***Equivalent to 1.00% disodium methanearsonate anhydride. ****Use as a pre-emergent herbicide, patented under U.S. Patent No. 2,923,634

CAUTION: Keep out of reach of children.

Avoid contact with skin, eyes, or clothing. Harmful if swallowed. Keep children and pets off treated area until watered in lawn and allowed to dry. Avoid contamination of feed and foodstuffs. Birds feeding on treated lawns may be harmed. Contamination of ornamental ponds will be harmful to fish. Avoid application on shrubs, flowers, fruit, or vegetable plants. Wash thoroughly after using.



for lawns


For maximum effectiveness, apply before or during the early stages of growth of weeds. Apply to lawns where weeds are growing or about to grow.

To prevent crabgrass, chickweed, dandelion, and other susceptible weeds, apply before or during the early stages of growth.


To kill existing broadleaf weeds such as dandelion, chickweed, and others, apply to lawns where weeds are growing or about to grow. Apply before or during the early stages of growth.

To control existing crabgrass, apply to lawns where crabgrass is growing. Apply to lawns where crabgrass is growing or about to grow. Apply before or during the early stages of growth.


Three Easy Steps To A Beautiful Lawn



1 For maximum effectiveness water lawn thoroughly two days before application. Determine square foot area to be covered, measure if necessary. (For example: $40' \times 50' = 2000$ sq. ft.) One 16 lb. bag covers 2000 sq. ft. (PAX Total will not burn when applied at recommended rate.)



2 Before spreading PAX Total, moisten the lawn with a fine spray or a sprinkler system. Do not allow water to puddle or flood. The moist lawn insures maximum effectiveness of PAX Total. After application do not water for 24 hours, then resume normal watering practices.




3 Set fertilizer spreader as suggested below. For best results, use a crisscross pattern as shown above. Spreader settings below are for crisscross patterns.

Spreading rate may vary with the age, model and condition of the spreader; check your spreader by a trial over a known-size area. (16 lb. bag covers 2000 sq. ft.)

Lawn Beauty		Scott		Jackson		Modern		Central		Sears	
MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING
18-57	7 $\frac{1}{2}$	35-5	7 $\frac{1}{2}$	18 in.	6 $\frac{3}{4}$	18 in.	6 $\frac{3}{4}$	20 in.	8	452-19191	6 $\frac{3}{4}$
		35-3	7								

Above settings are for crisscross applications. If your spreader is not included above, set spreader opening at lowest setting which gives even flow.

usually about 1/3 open. Spread in one direction and then crisscross. Continue crisscrossing until proper amount is applied. Broadcast spreaders are not recommended.




3234-12

crabgrass
 yellow foxtail
 knotweed
 common chickweed
 johnsongrass from seed
 purslane
 barnyardgrass
 dodder
 green foxtail
 palmer amaranth
 carpetweed

stinkweed grass
 stitchwort
 black medic
 dandelion
 speedwell
 plantain
 buckhorn
 shepherdspurge
 red clover
 filaree
 spotted spurge

wright groundcherry
 browntop Panicum
 Florida Pussley
 white clover
 English daisy
 henbit
 annual bluegrass
 lambsquarters
 goosegrass
 redroot pigweed
 spiny pigweed

witchgrass
 fall panicum
 sesbania
 Texas millet
 prostrate spurge
 ground ivy
 yarrow
 red sorrel



lawn moth larvae
 sod webworms
 white grubs
 earwigs
 ants
 sowbugs
 chinch bugs

WATERING. After application do not water or mow lawn for 24 hours; then resume normal watering practice.

SEEDING. If spring seeding is desired, apply seed as early as possible, and wait until new grass has been mowed twice before applying PAX Total. After an application of this product wait 60 days before reseeding.

WORKING. Complete any planned raking or digging of the lawn before you apply PAX Total. Disturbing the soil surface after the application may lessen the effectiveness of the product.

SHRUBS AND FLOWERS. Avoid making applications on flowers, shrubs, or other ornamental plantings. Avoid applications on a windy day.

SOFT AND TENDER GRASS. Succulent, tender grass may be discolored temporarily by applications during periods of high humidity and heat.

Manufacturer makes no express warranties of any kind except to give crabgrass control. The warranty of merchantability and all other implied warranties are disclaimed by the manufacturer. Buyer assumes all risks of use or handling, whether in accordance with directions or not.

Ornamental Turf