
Ornamenta / List


MICHOMORA Weed In Real

10-0-0 plus iron



KILLS WEEDY GRASSES

Kills Licegrass, Typh annun fannunt bleegrass Chabgrass, and nicht other wordy grasses ketards bornieda. Dee rear pancliter under weeds,



KILLS BROADLEAF WEEDS

It "s Challs, white clover, Hack medic, childwerd, and many other weeds. Prevents sparge. (See rear panel for other weeds).



KILLS LAWN INSECTS

Kills lawn moth larvae (sod webworms, cutworms, grubs earwigs, ants.

For Established Dichondra Lawns Caly

CAB :UN: K. ...? DUT OF REACH OF CHILDREN. Harmful if swallowed May cause irritation of ever nose throat and skin. Avoid breathing dust and avoid contact with skin eyes and clothing. Wash thoroughly after use. Avoid contamination of feed and foodstuffs.

Reep children and pets off treated areas ufitif material is watered into soil and dichondra leaves are dry


This product is toxic to fish and wildlife. Do not apply where runoff is likely to occur. Keep out of takes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ŀ,		 	(***
[. :		• • :	
Ε,	. •	 (3. 3)	
į			. : :

Giranda birana V	
10:	
9.7, .3	
(4 .) in more or fitting	
The state of the state of	CTT
Constitute, they	C .
Iron C. F. *	1 5 %
last (۶۰۰۰
	•
\$;	1
t .	
Maria transfer of the second	

PAX Company - 580 West 13th South - Salt Lake City, Utah 84115 U. S. E. A. Frystration No. 3204-16

super In dichondra weed in feed



SUPER MURE Weed in Reed

GENERAL WELD CONTROL For East results on by this product consult to the action of months on by the grant may time of the control of the contr

FOR ESTABLISHED DICHONDRA LAWNS ONLY

SPOT TREATING. I Aprily to were in area of recommended rates which is a special point of the tracking Conservation of passible. The tracking recommended recommended rates, in the resolution of the rest of the resolution of the resolution of the resolution of the r

CRACGPASS First districts of religions roots file to be added to a property of the product of a paying a control of and acceptance of the product of the paying and acceptance of the paying acceptance of th

OXAUS, CLOVER BLACK MEDIC: Make and the action of the control of t

PERMUDA To retard to modulo good in the special organization and a pumped a drawn for a fine and the special organization of the Lemma to the special organization of the special organization or the special organization of the special organization or the special organization organ

BLUEGEASS: Amount Application in Geography with a program of a second of the country of the second of the country of the count

STOKER to be the the control of the amount for every and March. It is not to be a second or the control of the

ACCEPTED


MAL 2H 1973

JUNDER THE PEDERAL INSECTICIDE ACT
FOR FCONOMIC POISON REGISTENT
POR FCON

MOW TO APPLY for a better dichondra lawn:



in the second se





DAMMON WEEDS CONTROLLED:


12.7.551.3

5

TROADITAVES



1. Measure area to be covered (for example-40 x 501. One 12 lb. bag covers 2000 sq. ft. Dichondra foliage should be dry at time of application. Remove clippings, leaves and other objects so that Dichondra Weed'n feed may contact soil surface evenly.



2. Set fertilizer spreader as suggested below. For best control follow crisscross pattern as illustrated, Recommended spreader settings given below are for crisscross patterns. CAUTION: Never make sharp turns with spreader open. Always close spreader while changing directions.



3. Immediately after spreading, wash Dichardra Weed'n feed off the dichardra folioge with a medium spray from a handheld nozzle. Sprinkling systems and stationary sprinklers are less setisfactory. After application do not water for 24 hours, then resume normal watering practice. Do not allow water to puddle or flood.

COMMON WEEDS CONTROLLED:

GRASSES

crabgrass bluegrass ryagrass goosegrass

barnyard grass
johnsongrass
(from seed only)
stinkgrass

BROADLEAVES

black medic chickweed (common) chickweed (mouseear)

ear)
purstane
shepherds purse
spiny amaranth

groundsel
sandbur
corn spurry
Florida pusley
spotted spurge
white clover
pearlwort
knotweed

pigweed
lambsquarter
evening primrose
red sorrell
carpetweed
smartweed
cheat
peppergrass
crowfoot

INSECTS CONTROLLED:

bermudagrass

fall panicum

watergrass

witchgrass

annual sedge

foxtail

annual bluegrass

(Pod annua)

Lawn moth larvae (sod webwerms), Cutworms, white grubs, earwigs, ants,

SUGGESTED SPREADER SETTINGS: Spreading rate may vary with the age, model, and condition of the spreader. Check your spreader over an area of known size. (One 12 lb. bag covers 2000 sq. ft.). Recommended settings below are for crisscross applications. If your spreader is not included below, set spreader opening at lowest setting which gives an even flow, usually about one-third open. Spread in one direction and then crosswise. Continue crisscrossing until proper amount is applied.

Bandini 8½ Lawn Beauty

Scott 9½

Modern 7 Central 8½

Germain 8½

RESEEDING: Dichondra lawns cannot be reseeded for 60 days after application. Materials that provide effective control of weeds may affect seedlings which germinate after treatment. Proper fertilization and watering will encourage the existing dichondra to fill in the thin or bare spots formerly occupied by weeds.

TREES & SHRUBS:

Do not apply on or near young trees or young shrubs.

BLADEGRASS LAWNS:

Do not use on blade grass lawns.

SUMMER APPLICATIONS:

- 7. Water dichondra lawn thoroughly the day before PAX appli-
- 2. Do not apply on days when temperature is 85 degrees or above.

GUARANTEE: Manufacturer guarantees this product to give results as a outlined on this bag or the purchaser may get his manufacturer. No other express warranties of any kind are made, except to conform to the promises and affirmation of facts made on the label. The warranty of merchantability, all other implied warranties and liability of the manufacturer without fault are expressly disclaimed.

Best

81/2

Buyer assumes all risk of personal injury or damage to property, except for negligent mu-labeling of the product. All claims must be presented to manufacturer in writing, within thirty days from the date a breach is discovered and an action thereon must be commenced within one year after the cause of action has occurred.

super and dichondra weed in seed

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes, or clothing. Harmful if swallowed. Avoid contamination of feed and foodstuffs. Wash thoroughly after use.

Keep children and pets off treated areas until material is watered in lawn and allowed to dry.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not store near fertilizers, seeds, insecticides, or langicides. Do not reuse container. Destroy when empty.

