




DO NOT USE ON DICHONDRA
OR BENTGRASS



action

WEED 'N FEED

WHEN TO APPLY:

MAY BE APPLIED ANYTIME OF YEAR. FOR BEST RESULTS APPLY WHEN WEEDS ARE GROWING ACTIVELY.

HOW TO APPLY FOR A BETTER LAWN




HOW TO APPLY FOR A BETTER LAWN



1 Measure area to be covered (for example 50 x100 ft.). One 22 lb. bag covers 5000 sq. ft. Best results will be obtained if lawn is cut two to three days before the application. For maximum effectiveness water lawn thoroughly the day before application. Application in the evening or early morning will be most effective.



2 Immediately before spreading PAX ACTION, wet the lawn well. If sections dry before they can be reached, re-wet, but avoid washing off material already applied. Do not water lawn for at least 24 hours after application. Do not apply on days when temperature is over 85°.



3 Set spreader as recommended below. Fill spreader where spilled material will not fall on lawn. Shake spreader to settle the material. Make application in two crisscross coverages as illustrated. Spreader settings below are for crisscross applications.

SUGGESTED SPREADER SETTINGS

Spreading rate may vary with the age, model, and condition of the spreader; check your spreader by a trial over a known-size area. (22 lb. bag covers 5000 sq. ft.)

Lawn Beauty		Scott		Modern		Central		Sears	
MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING
18-57	7 $\frac{3}{4}$	35-8	7 $\frac{1}{2}$	18 in.	6 $\frac{3}{4}$	20 in.	8	452-19191	6 $\frac{3}{4}$
		35-3	7						

Above settings are for crisscross applications. If your spreader is not included above, set spreader opening at lowest setting which gives even flow, usually about 1/3 open. Spread in one direction and then crisscross. Continue crisscrossing until proper amount is applied. Broadcast spreaders are not recommended.

SOME COMMON WEEDS CONTROLLED

black medie
buttercup
chickweed
cinquifol
clover
dandelion
docks

English ivy
heal-all
henbit
knotweed
lambquarters
mallow
morning glory

(1) mustard
Oxalis
pennywort
pepperweed
pigweed
plantain

prostrate spurge
purslane (pusley)
ragweed
red sorrel
sheepsorrel
smartweeds
Spanish nettle

spotted spurge
thistle
velvet leaf
vervain
Yarrow

NOTICES

SEEDING:

Apply to established lawns only. May be used on reseeded lawns six weeks after reseeding.

SHRUBS AND FLOWERS:

Avoid making applications on flowers, shrubs, or other ornamental plantings. Do not apply on or near young trees. Prevent drift of product to desirable plants.

(2) May injure carpetgrass, St. Augustine, bentgrass, dichondra, centipede, or newly seeded lawns.

(3) Do not apply to areas underlaid by roots of desirable trees and shrubs.

Repeat applications may be necessary on difficult to kill weeds.

imum effectiveness water lawn thoroughly the day before application. Application in the evening or early morning will be most effective.

for at least 24 hours after application. Do not apply on days when temperature is over 85°.

ages as illustrated. Spreader settings below are for crisscross applications.

SUGGESTED SPREADER SETTINGS

Spreading rate may vary with the age, model, and condition of the spreader; check your spreader by a trial over a known-size area. (22 lb. bag covers 5000 sq. ft.)

Lawn Beauty		Scott		Modern		Central		Sears	
MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING	MODEL	SETTING
18-57	7 $\frac{3}{4}$	35-8	7 $\frac{1}{2}$	18 in.	6 $\frac{3}{4}$	20 in.	8	452-19191	6 $\frac{3}{4}$
		35-3	7						

Above settings are for crisscross applications. If your spreader is not included above, set spreader opening at lowest setting which gives even flow, usually about 1/3 open. Spread in one direction and then crisscross. Continue crisscrossing until proper amount is applied. Broadcast spreaders are not recommended.

SOME COMMON WEEDS CONTROLLED

black medic
buttercup
chickweed
cinqufoil
clover
dandelion
docks

English ivy
heal-all
henbit
knotweed
lambquarters
mallow
morning glory

(1) mustard
Oxalis
pennywort
pepperweed
pigweed
plantain

prostrate spurge
purslane (pusley)
ragweed
red sorrel
sheepsorrel
smartweeds
Spanish nettle

spotted spurge
thistle
velvet leaf
vervain
Yarrow

NOTICES

SEEDING:

Apply to established lawns only. May be used on reseeded lawns six weeks after reseeding.

(2) May injure carpetgrass, St. Augustine, bentgrass, dichondra, centipede, or newly seeded lawns. X

SHRUBS AND FLOWERS:

Avoid making applications on flowers, shrubs, or other ornamental plantings. Do not apply on or near young trees. Prevent drift of product to desirable plants.

(3) Do not apply to areas underlaid by roots of desirable trees and shrubs. X

Repeat applications may be necessary on difficult to kill weeds.

GUARANTEE: Manufacturer guarantees this product to give results as outlined on this bag or the purchaser may get his money refunded by the manufacturer. No other express warranties of any kind are made, except to conform to the promises and affirmation of facts made on the label. The warranty of merchantability, all other implied warranties and liability of the manufacturer without fault are expressly disclaimed.

Buyer assumes all risk of personal injury or damage to property, except for negligent mislabeling of the product. All claims must be presented to manufacturer in writing, within thirty days from the date a breach is discovered and an action thereon must be commenced within one year after the cause of action has occurred.



action WEED 'N FEED

18-4-4

KILLS DANDELIONS

CLOBBERS CHICKWEED

KILLS HENBIT

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Keep children off treated area until grass is allowed to grow. Avoid contamination of feed and food. Avoid application on shrubs, flowers, fruit or vegetable plants. Wash thoroughly after using. Do not store near fertilizers, seeds, insecticides or herbicides.

ACTIVE INGREDIENTS:

Alkanolamine salt of 2,4-dichlorophenoxyacetic acid 0.870%
Dimethylamine salt of dicamba (3,6-dichloro-p-anisic acid) 0.125%

INERT INGREDIENTS:

2,4-dichlorophenoxyacetic acid equivalent 0.500%
3,6-dichloro-p-anisic acid equivalent 0.300%

14

GUARANTEE

Total N 4.0
14.0
Available
Soluble
Residual
Calcium

RAY COMPANY, 2800 West 10th South, Salt Lake City, Utah 84115

A. S. D. A. Reg.



18-4-4

GLOBBERS CHICKWEED

KILLS HENBIT

BLASTS KNOTWEED

ACTIVE INGREDIENTS:

Diethylamine salt of 2,4-dichlorophenoxyacetic acid	0.870%
Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)**	0.125%
INERT INGREDIENTS:	99.005%
	100.00%

2,4-dichlorophenoxyacetic acid equivalent	50.520%
3,6-dichloro-o-anisic acid equivalent	0.10%

GUARANTEED FERTILIZER ANALYSIS:

Total Nitrogen (N)	18.0%
4.0% Ammoniacal Nitrogen	
14.0% from Urea	
Available Phosphoric Acid (P ₂ O ₅)	4.0%
Soluble Potash (K ₂ O)	4.0%
Potential Acidity 300 lbs.	
Calcium Carbonate Equivalent	

Salt Lake City, Utah 84115

U. S. D. A. Registration No. 102319

WEED 'N FEED

NI WT 22 LBS




P A



AGT 100

WEED 'N FEED



KILLS DANDELIONS



GLOBE AMARANTH




KILLS HENBIT



BLASTS KNOTWEED



KILLS DANDELIONS



CLOBBERS CHICKWEED



KILLS HENBIT



BLASTS KNOTWEED

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. Keep children and pets off treated area until grass is allowed to dry. Avoid contamination of feed and food stuffs. Avoid application on shrubs, flowers, fruit, or vegetable plants. Wash thoroughly after using. Do not store near fertilizers, seeds, insecticides, or fungicides.

Do not reuse empty container. Destroy when empty.

ACTIVE INGREDIENTS:

Alkanolamine salt of 2,4-dichlorophenoxyacetic acid*	0.870%
Dimethylamine salt of dicamba (3,5-dichloro-p-anisic acid)**	0.125%
INERT INGREDIENTS:	99.005%
	100.00%

*2,4-dichlorophenoxyacetic acid equivalent	0.520%
**3,5-dichloro-p-anisic acid equivalent	0.104%

GUARANTEED FERTILIZER ANALYSIS:

Total Nitrogen (N)	18.0%
4.0% Ammoniacal Nitrogen	
14.0% from Urea	
Available Phosphoric Acid (P ₂ O ₅)	4.0%
Soluble Potash (K ₂ O)	4.0%
Potential Acidity 300 lbs.	
Calcium Carbonate Equivalent per ton	

PAX COMPANY, 580 West 13th South, Salt Lake City, Utah 84115

U. S. D. A. Registration No. 3234



action

WEED 'N FEED

Nt. Wt. 22 LBS

KILLS DANDELIONS

GLOBBERS CHICKWEED

KILLS H

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. Keep children and pets off treated area until grass is allowed to dry. Avoid contamination of feed and food-stuffs. Avoid application on shrubs, flowers, fruit, or vegetable plants. Wash thoroughly after using. Do not store near fertilizers, seeds, insecticides, or fungicides.

Do not reuse empty container. Destroy when empty.

ACTIVE INGREDIENTS:

Alkanolamine salt of 2,4-dichlorophenoxyacetic acid*	0.870%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)**	0.125%

INERT INGREDIENTS:

99.005%

100.00%

2,4-dichlorophenoxyacetic acid equivalent	0.520%
(3,6-dichloro-o-anisic acid) equivalent	0.104%

PAX COMPANY, 580 West 13th South, Salt Lake City, Utah 84115



caution

CHICKWEED

KILLS HENBIT

BLASTS KNOTWEED

ACTIVE INGREDIENTS:

ethylamine salt of 2,4-dichlorophenoxyacetic acid*	0.870%
ethylamine salt of dicamba (3,6-dichloro-o-anisic acid)**	0.125%
INERT INGREDIENTS:	99.005%
	<u>100.00%</u>

2,4-dichlorophenoxyacetic acid equivalent	0.520%
3,6-dichloro-o-anisic acid) equivalent	0.104%

GUARANTEED FERTILIZER ANALYSIS:

Total Nitrogen (N)	18.0%
4.0% Ammoniacal Nitrogen	
14.0% from Urea	
Available Phosphoric Acid (P ₂ O ₅)	4.0%
Soluble Potash (K ₂ O)	4.0%
Potential Acidity 300 lbs.	
Calcium Carbonate Equivalent per ton.	

Salt Lake City, Utah 84115

U. S. D. A. Registration No. 3234.