

LIGHTNING BRAND

# 3.4% COPPER 90% SULFUR DUST

### DIRECTIONS FOR USE

Apply as a dust without dilution. Adequate and uniform coverage of the entire plant is essential. Dust should be applied when the wind velocity is less than 5 miles per hour. The best hours are usually between 5:00 P.M. and 9:00 A.M. For disease control begin application before diseases occur or appear and repeat at 7 - 10 day intervals as needed.

CROP & DISEASE	RATE PER ACRE
PEANUTS - Leaf Spot	35 - 50 lbs.

Apply at first sign of disease and repeat at 2 week intervals. Dust 3 or 4 times depending on how early disease appears. Do not apply close to harvest.

### ACTIVE INGREDIENTS

Copper (as metal) in the form of cuprous hydroxide	3.4%
Sulfur	90.0%
INERT INGREDIENTS	6.6%

Total 100%

Screen Test Sulfur through 325 mesh = 93%

NET CONTENTS (LB(S))

## CAUTION


KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed. Sulfur may burn foliage when temperatures are excessively high (90° F. or above). Do not apply any sprays containing oil, dinitro materials or Captan within 30 days to foliage dusted with sulphur.

Manufactured For  
**KERR-McGEE CHEMICAL CORP.**

Oklahoma City, Okla.

**WARRANTY:** Our merchandise is guaranteed as set forth on our labels. Kerr-McGee Chemical Corp. can accept no responsibility for injury to persons or animals or for loss or damage to crops, soil, or other property resulting from failure to follow directions for use or to observe cautions printed on this package.



U.S. PAT. NO. 32381