

**ACTIVE INGREDIENT:**  
Sulphur not less than 90%  
**INERT INGREDIENTS:**  
Not more than 10%

**MESH TEST**  
Mixtures containing dry Sulphur(S) when sold for DUSTING vegetation, of Contained Sulphur not less than following mesh test guaranteed: 325 and finer 93%.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

**DIRECTIONS**  
**CITRUS:** (Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines). *Rust Mite*—Use 5 to 10 pounds per 100 gallons water. During cool weather, use 10 lbs. of Standard Wettable Sulphur per 100 gallons water. At least 8 lbs. of Standard Wettable Sulphur should be used in miticidal sprays for the combined control of rust mite and purple mite.

**PEACHES:** *Brown Rot, Powdery Mildew, Scab*—4 to 6 lbs. per 100 gals. water. Make applications during pink, petal fall and cover spray periods. Also as ripening of fruit approaches for Brown Rot control.

**APPLES:** *Apple Scab, Powdery Mildew*—6 to 8 lbs. per 100 gals. water. Make application between periods of rainfall. Repeat as necessary to cover new foliage growth during pink, petal fall and cover spray periods.

**CAUTION:** Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality.

During periods of high temperature, sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use Standard Wettable Sulphur with oil or within 3 weeks after oil application.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends

to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

Do not re-use container. Destroy by burying in a safe place away from water supplies. Do not burn.

EPA Reg. No. 3238-12


NEW WEIGHT  
**50 lbs.**

Manufactured by

**Agrico**

Chemical Company

Crop Protection Chemicals Division  
P. O. Box 3451 • Tulsa, Oklahoma 74101



#### ACTIVE INGREDIENT:

Sulphur not less than . . . . . 90%

#### INERT INGREDIENTS:

Not more than . . . . . 10%

#### MESH TEST

Mixtures containing dry Sulphur(S) when sold for DUSTING vegetation, of Contained Sulphur not less than following mesh test guaranteed: 325 and finer 93%.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

#### DIRECTIONS

**CITRUS:** (Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines). *Rust Mite*—Use 5 to 10 pounds per 100 gallons water. During cool weather, use 10 lbs. of Standard Wettable Sulphur per 100 gallons water. At least 8 lbs. of Standard Wettable Sulphur should be used in miticidal sprays for the combined control of rust mite and purple mite.

**PEACHES:** *Brown Rot, Powdery Mildew, Scab*—4 to 6 lbs. per 100 gals. water. Make applications during pink, petal fall and cover spray periods. Also as ripening of fruit approaches for Brown Rot control.

**APPLES:** *Apple Scab, Powdery Mildew*—6 to 8 lbs. per 100 gals. water. Make application between periods of rainfall. Repeat as necessary to cover new foliage growth during pink, petal fall and cover spray periods.

**CAUTION:** Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality.

During periods of high temperature, sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use Standard Wettable Sulphur with oil or within 3 weeks after oil application.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends

to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

Do not re-use container. Destroy by burying in a safe place away from water supplies. Do not burn

EPA Reg. No. 3238-12

NEW WEIGHT

**50 lbs.**

Manufactured by

**Agrico**

Chemical Company

Crop Protection Chemicals Division  
P. O. Box 3451 • Tulsa, Oklahoma 74101