

STANDARD BRAND 5% SEVIN 75% SULFUR DUST

DIRECTIONS FOR USE

To be used as a dust without dilution in conventional ground or air dusters. Dusts should be applied when the wind velocity is less than 5 miles per hour. The best hours for dusting are usually between 5:00 p.m. and 9:00 a.m. For disease control begin application before diseases occur and repeat at 7-10 day intervals as needed.

Crop, Insects & Diseases	Rate per Acre
--------------------------	---------------

PEANUTS —Blister Beetles, Mexican Bean Beetles	20 lbs.
--	---------

Traps, and eggs of Noctuid	
-------------------------------	--

Traps, and eggs of cornered Alfalfa Hopper, Cucurbit Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Green Cloverworm, Leaf Spot	20 lbs.
---	---------

Apply at first sign of infestation and repeat at 2-week intervals. Dust 3 or 4 times, depending on how early infestation appears.

Armyworms, Webworms, Earworms	50 lbs.
----------------------------------	---------

Some leaf injury may occur if applied when tender foliage is wet.

ACTIVE INGREDIENTS

Carbaryl (1-Naphthyl N-methylcarbamate)	5.0%
Sulfur	75.0%

INERT INGREDIENTS

Screen Test, Sulfur, through 325 mesh 9.0%	
--	--

Registered Trademark of Union Carbide Corporation, New York, N.Y.

NET CONTENTS 50 LBS.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful.

Wash hands before eating. Take cover or bath if work. Wear regular long-sleeved clothes, clothing, change to clean clothing daily. Do not contaminate feet and feet.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

Application of the dust, which contains sulfur, should not be made during periods of excessively high temperatures. Do not apply dusts or sprays containing oil, dithionates or captan within 48 hours of foliage dusted with sulfur. Sulfur may burn foliage, especially if temperatures are excessively high (90 degrees or above).

Application of this dust to bees, especially at the time of flowering, may seriously injure the crop.

Bees are important pollinators and should be protected during such periods. Consult your State Agriculture Extension Service.

Manufactured By

Standard Spray and Chemical Company

NEW YORK, N.Y. CHICAGO, ILL. PHOENIX, ARIZ.

NOTICE TO USER

Standard Spray and Chemical Company can accept no responsibility for injury to persons or animals or for loss or damage to crops, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

