

# STANDARD BRAND

# 5% SEVIN® 4% COPPER 75% SULFUR DUST

## DIRECTIONS FOR USE

To be used as a dust without dilution on conventional ground or air drifters. Dusts should be applied when the wind velocity is less than 5 miles per hour. The best hours for dusting are usually between 5:00 P.M. and 9:00 A.M. For disease control begin application before diseases occur and repeat at 7-10 day intervals as needed.

CROP	INSECTS & DISEASES	RATE PER ACRE	INTERVAL
			Days Application to Harvest
PEANUTS	Blister Beetles, Mexican Bean Beetle, Grasshoppers	10-20 lbs	
	No time limitation		
	Thrips, Leafhoppers, Three-cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Green Cloverworm, Leaf Spot	20 lbs	
	Apply at first sign of disease and repeat at 2 week intervals. Dust 3 or 4 times depending on how early disease appears.		
	Armyworms, Webworms, Corn Earworm, Stink Bug	20-30 lbs	
	Some leaf injury may occur if applied when tender foliage is wet.		
SUGAR BEETS	Leafhoppers, Armyworms, Webworms, Grasshoppers, Spider Mites, Sugar Beet Leaf Spot	20-40 lbs	14 days
	Use in hot weather in desert areas for mites. May cause burn, but injury less severe following an irrigation.		
BEANS	Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Downy Mildew, Anthracnose, Powdery Mildew, Angular Leaf Spot, Rust, Leaf Spot, Bean Leaf Beetle, Cucumber Beetle, Western Bean Cutworm, Japanese Beetle, Velvet Bean Caterpillar, Armyworms	25-40 lbs	0 day
	Sulfur may injure some varieties. Do not feed foliage to dairy animals or animals being finished for slaughter.		
BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER			
	Fleabeetles, Harlequin Bug, Corn Earworm, Armyworms, Cabbage Caterpillars, Downy Mildew, Leaf Spot	25-40 lbs	3 days
CARROTS	Fleabeetles, Leafhoppers, Harlequin Bug, Six-spotted Leafhopper, aster yellows vector, Corn Earworm, Armyworms, Cabbage Caterpillars, Spittlebugs, Lygus Bugs, Stink Bugs	20-40 lbs	0 day
	Leaf Spot, Early & Late Blight		

### ACTIVE INGREDIENTS:

Carbaryl (1 Naphthyl N methylcarbamate)	5.0%
Copper, as metallic (in the form of copper oxide)	4.0%
Sulfur	75.0%
INERT INGREDIENTS	16.0%
Total	100.0%

Screen Test Sulfur through 325 mesh - 95%

\* Registered Trademark of Union Carbide Corporation for Carbaryl Insecticide

NET CONTENTS \_\_\_\_\_ LB(S)

## CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Do not take internally. Do not breathe dust. Avoid prolonged or repeated breathing of dust. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothes. Change to clean clothing daily. Keep out of reach of children and animals.

Application of this material which contains sulfur should not be made just prior to or during periods of excessively high temperatures. Do not apply any sprays containing oil, dinitro materials or captan within 30 days to foliage dusted with sulfur. Sulfur may burn foliage when temperatures are excessively high (90 F or above).

WARNING - For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

NOTE FOR PHYSICIAN: CARBARYL IS A MODERATE, REVERSIBLE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.

Mfg. by Standard Spray & Chemical Co.,  
Lakeland, Fla. — Severn, N. C.

## DIRECTIONS FOR USE

EGGPLANT Fleabeetles, Cucumber Beetles, Tomato Fruitworm, Corn Earworm, Tomato Hornworms, Tobacco Hornworms, Leafhoppers, Lygus Bugs, Fall Armyworms, European Corn Borer, Stink Bugs, Lace Bugs, Alternaria, Anthracnose, Phomopsis, Powdery Mildew

20-50 lbs 0 day

PEPPERS Fleabeetles, Cucumber Beetles, Tomato Fruitworm, Corn Earworm, Tomato Hornworms, Tobacco Hornworms, Leafhoppers, Lygus Bugs, Fall Armyworms, European Corn Borer, Stink Bug, Lace Bugs, Anthracnose, Cercospora (Leaf Spot), Bacterial Spot, Frog Eye Spot, Powdery Mildew

30-60 lbs 0 day

POTATOES Armyworms, Cucumber Beetles, Flea Beetles, Leafhoppers, Plant Bugs, Lygus Bugs, Colorado Potato Beetle, Fall Armyworms, European Corn Borer, Stink Bugs, Lace Bugs, Early & Late Blight

15-40 lbs 0 day

Foliage application only

TOMATOES Tomato Fruit Worm, Tomato Hornworms, Tobacco Hornworms, Fall Armyworms, Leafhoppers, Stink Bugs, Fleabeetles, European Corn Borer, Corn Earworm, Armyworms, Stink Bugs, Russet Mite, Lygus Bugs, Alternaria, Septoria Leaf Spot, Anthracnose, Late Blight, Leaf Mold

20-60 lbs 0 day

STRAWBERRIES Fleabeetles, Lygus Bugs, Meadow Spittle Bugs, Leafhoppers, Strawberry Leaf Roller, Strawberry Weevil, Leaf Spot, Powdery Mildew

20-40 lbs 1 day

GRAPES Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, Powdery Mildew, Black Rot, Downy Mildew, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Red-banded Leaf Roller

30-60 lbs 0 day

### NOTICE TO BUYER

Seller warrants that the within product will comply with the description of the goods contained on the label. If Buyer's remedy under this warranty shall be limited exclusively to the remedy of resale, this warranty is expressly limited to all other warranties and conditions, expressed or implied, including, but not limited to, and all other obligations or liabilities of the seller, in whole or in part of anyone in seller's line of title. If an exchange or refund is accepted under these conditions, Buyer may return it for full credit.

USDA REG. NO. 3238-16