

3238-36

04/26/1976

1/1

ACCEPTED

APR 26 1976

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. **3238-36**



Micro-Sperse[®] Wettable Sulphur

ACTIVE INGREDIENT:
SULPHUR 90%
INERT INGREDIENTS: 10%
Particle Size—Average 5 Micron.

**CAUTION: KEEP OUT OF
REACH OF CHILDREN**

DIRECTIONS

APPLE: Scab, Powdery Mildew—Use 6 to 8 lbs. per 100 gal. of water in the pre-bloom, bloom and petal-fall sprays; 4 to 6 lb. in the cover sprays.

PEACH: Brown Rot, Scab, Powdery Mildew—Use 6 to 8 lbs. per 100 gal. of water in the pink and bloom sprays; 4 to 6 lbs. in the petal-fall, shuck and cover sprays. A coarser, less adherent sulphur such as .325-mesh spraying or dusting sulphur preferred in the pre-harvest sprays.

BEANS, PEAS: Powdery Mildew, Rust—Use 6 to 8 lbs. per acre in sufficient water for thorough coverage. Begin applications at first sign of infection and repeat in 7 to 14 days as necessary. Keep new growth protected.

ROSES: Black Spot, Powdery Mildew—Use 4 to 8 lbs. per 100 gal. of water (3 to 6 tablespoons per gal.) Make first application as soon as growth starts in the spring; repeat application at 7 to 14-day intervals. Keep new growth protected.

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines):
Rust Mite—Use 5-6 lbs. per 100 gallons of water.

DIRECTIONS FOR USE

Pour recommended amount of Micro-Sperse[®] Wettable Sulphur into partially filled spray tank. Add balance of water to fill tank. Keep agitated while filling and spraying.

CAUTION: Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality.

During periods of high temperature sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use Micro Sperse[®] Sulphur with oil or within 3 weeks after oil application.

When using lead arsenate with Micro-Sperse[®] Wettable Sulphur, always add the same amount of hydrated lime, or other softener, as would be used when applying lead arsenate without sulphur.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with di-

rections under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonable foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

Do not re-use container. Destroy by burying in a safe place. Do not burn.

EPA Est. 3238-FL-1
EPA Reg. No. 3238-36

NET WEIGHT
30 lbs.

BEST DOCUMENT AVAILABLE

Manufactured by



Chemical Company

Crop Protection Chemicals Division
P. O. Box 3451 • Tulsa, Oklahoma 74101