

STANDARD BRAND

# Toxaphene-Sulphur 20-40 DUST

An Agricultural Insecticide-Fungicide

ACTIVE INGREDIENTS:

*Toxaphene	20.0%
Sulphur	40.0%

INERT INGREDIENTS

	40.0%
TOTAL	100.0%

\*Technical Chlorinated Camphene (Chlorine Content 67% to 69%)

## CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Do not contaminate food and feed.

### OBSERVE THE FOLLOWING USE PRECAUTIONS.

Observe the limitations given on the label for specific crops in order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water nor apply to any area not specified on the label.

Do not use this product on feed or forage to be fed to dairy animals or livestock being finished for slaughter.

Do not use Toxaphene or Sulphur on cucurbits (cantaloupe, cucumber, squash, melon) or young plants, as injury to plants may result.

**Do not re-use container. Destroy by burying in a safe place away from water supplies. Do not burn.**

Manufactured By



STANDARD SPRAY  
and chemical company

Lakeland, Florida    Severn, North Carolina    Columbia, Alabama

NET CONTENTS 50 POUNDS

**PEPPERS:** Army worms, Blister Beetle, Colorado Potato Beetle, Cutworm (climbing and surface-feeding), Flea Beetle, Hornworm, Serpentine Leaf Miner—Use 10 to 15 lbs. per acre. Fruitworm, Pepper Weevil—Use 15 lbs. per acre. Do not apply within 5 days of harvest.

**TOMATOES:** Armyworms, Blister Beetle, Cabbage Looper, Colorado Potato Beetle, Cutworms (climbing and surface-feeding) Flea Beetle, Pinworm, Russet Mite, Serpentine Leaf Miner, Tomato Fruitworm, Tomato Hornworm—Use 10 to 25 lbs. per acre. Do not apply within 3 days of harvest.

Sulphur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply Sulphur to sensitive crops when temperatures have exceeded or are likely to exceed 90° F.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information and for special rate applications needed with special cultural practices.

**NOTICE** The buyer warrants that the product conforms to the label description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use, but makes no warranty, express or implied, for a particular purpose, express or implied, or for the use, storage or handling of the product in violation of label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by the buyer as to the risk of any injury to humans, animals, birds, or plants not limited to those specifically mentioned, and assumes such the manufacturer's liability for damage caused by the use of the product on crops and animals.

EPA Reg. No. 3238-37

**DIRECTIONS FOR USE:** Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation and infection and repeat at 5- to 10-day intervals, depending upon the severity of the disease and insect conditions, but observe use limits given for specific crops.

For the control of diseases persisting after the prescribed time limits have expired, an effective fungicide should be used alone, in a manner which will not leave residues in excess of established tolerances.

**RECOMMENDATIONS: BEANS:** Armyworms, Bean Leaf Roller and Leafhopper, Climbing Cutworms (apply only in seedling stage), Corn Earworm, Cowpea Curculio, Lygus, Powdery Mildew, Rust, Salt-marsh Caterpillars, Thrips—Use 15 lb. per acre. Do not apply to green or snap beans within 7 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Cabbage Looper, Corn Earworm—Use 10 to 20 lb. per acre. Imported Cabbageworm, Cutworms (climbing and surface-feeding), Fall Armyworm, Serpentine Leaf Miner, Thrips—Use 10 lb. per acre. Flea Beetles—Use 15 lb. per acre. Do not apply after edible parts start to form or within 30 days of harvest. On CABBAGE, limit dosage to no more than 12.5 pounds per acre. For dosages above 12.5 pounds per acre, do not apply to cabbage after heads start to form.

**COTTON:** Overwintered Boll Weevils, Flea Beetles, Fleahoppers, Thrips (early-season control)—Apply 10 lbs. per acre. Make 2 to 4 applications at 7 to 10-day intervals. Cotton Leafworm, Fall Armyworm, Lygus and other Mirids—Use 10 to 15 lbs. per acre. Boll Weevil, Bollworm—Use 15 to 20 lbs. per acre. Apply at 4 to 5-day intervals until control is obtained. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

ACCEPTED  
Nov. 16, 1972  
3238-37

3238-37