

DIRECTIONS FOR USE

To be used as a dust without dilution in conventional ground or air dusters. Dusts should be applied when the wind velocity is less than 10 miles per hour. The best hours for dusting are usually between 5:00 P.M. and 9:00 A.M. Use of this material other than in accordance with this label may cause serious plant, crop, or person injury, excessive residues, unsatisfactory control, or other unintended consequences.

Peanuts: For the control of *Cercospora* Leaf Spot, apply at the rate of 25 lbs. per acre as soon as there is any evidence of leaf spotting. This material contains Borax and is for use only on peanuts for Boron deficiency. The application of 25 lbs. per acre of this material will supply 1/2-lb. of Boron per acre.

WARNING: USE IN EXCESS OF 25 LBS. PER ACRE MAY INJURE THE CROP. IT SHOULD NOT BE USED MORE THAN ONE TIME IN A SINGLE GROWING SEASON. ITS USE ON OTHER CROPS MAY RESULT IN SERIOUS INJURY.

For additional applications of material to control *Cercospora* leaf spot, applications should be made at 7 to 10-day intervals as needed with MICRO-SPERSE[®], Peanut Dust with Calcium Sulfate (MAGI-CAL[™] added) or MICRO-SPERSE[®] "S" Peanut Dust with Calcium Sulfate (MAGI-CAL[™] added) (these products do not contain Boron) until a minimum of at least five applications have been made.

MAGI-CAL[™] contains calcium which supplements your regular landplaster practice. Some leaf injury may occur if applied when tender foliage is wet.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the peanut dust and spraying program best suited to your local conditions.

MICRO-SPERSE[®] BORAX Peanut Dust

with Calcium Sulfate
(MAGI-CAL[™] added)

As a fungicide for control of leaf spot	44%
As a source of Boron for Boron deficiency	10%
As a source of Calcium for Calcium deficiency	36%
As a source of Sulfur for Sulfur deficiency	10%
Calcium Sulfate	36.0%
Boron as 10% equivalent to 1/2 lb. borax per 25 lbs.	2.0%

MICRO-SPERSE[®] is a registered trademark of Standard Spray and Chemical Company for premium grade insecticides and fungicides having an average particle size of less than 5 microns.

Pat. Pending

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Sulphur may burn foliage when temperatures are excessively high (90° F. or above). Do not apply any sprays containing oil, dinitro materials or Captan within 30 days to foliage dusted with sulphur. Do not apply to any area not specified on label.

Do not reuse container. Destroy by burying in a safe place. Do not burn.

Manufactured By

Standard Spray and Chemical Company
LAKELAND, FLORIDA SEVERN, NORTH CAROLINA
COLUMBIA, ALABAMA

Net Weight 50 Lbs.

Notice to User: Standard Spray and Chemical Company can accept no responsibility for injury to persons or animals or for loss or damage to crops, soil or other property resulting from failure to follow directions of use or to observe cautions printed on the package.

ACCEPTED

July 2-1971

3238-48

USDA Reg. No. 3238-48

5/71