

# MICRO-FLO™ SULPHUR

NET CONTENTS 5 GALLONS

## RECOMMENDATIONS

Use of this material in a manner other than in accordance with this label may cause serious plant, crop, or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

Unless otherwise specified for specific tree fruits, dosage rates are given as pints of MICRO-FLO Sulphur per 100 gallons water for use in a thorough coverage spray.

### APPLE:

Scab: Pre-bloom through Calyx sprays 1½ to 3½ pints  
Cover sprays 1 to 2 pints

### Powdery Mildew:

Pre-bloom through Calyx sprays 2½ to 3 pints  
Cover sprays 2 pints

### PEACH:

~~Powdery~~ Mildew, Brown Rot, Scab:  
Pink and bloom sprays 1½ to 3½ pints  
Petal fall, shuck and cover sprays 1 to 1¾ pints

### CITRUS: (Grapefruit, Lemons, Limes, Oranges,

Tangelos and Tangerines):  
Rust Mite: 1 to 4 pints  
Do not apply within 4 weeks of an oil application.

### PEANUTS:

Rust: Apply 2 to 4 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

### ACTIVE INGREDIENT:

Sulphur as S 52.0%

### INERT INGREDIENTS:

48.0%

Total 100.0%

ACCEPTED

June 16 1977

Contains 6.0 pounds Sulphur per gallon!

3035-

## CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not apply to any area not specified on label. Do not re-use empty container. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.

Manufactured By



STANDARD SPRAY  
and chemical company

Lakeland, Florida

Sevens, North Carolina

Columbia, Alabama

**CAUTION:** Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality.

During periods of high temperature sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use MICRO-FLO Sulphur with oil or within 4 weeks of an oil application.

### DIRECTIONS FOR MIXING BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operations. **DO NOT ALLOW MIXTURE TO STAND.** Failure to maintain agitation will cause MICRO-FLO Sulphur to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of MICRO-FLO Sulphur act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulphur in any form is corrosive material. **To reduce the effect, equipment should be flushed daily.**

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These include, but are not limited to damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants, crops, and personal injury.

EPA. Reg. No. 3238-57