

STANDARD BRAND

ACCEPTED

FEB 14 1973

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 3238-59

# Toxaphene 6 Emulsifiable Concentrate Insecticide

Standard Brand Toxaphene 6 Emulsifiable Concentrate Insecticide is a water emulsifiable insecticidal concentrate containing 6 lbs. actual Toxaphene per gallon.

## DIRECTIONS FOR USE

For best results premix with a small amount of water and thoroughly agitate before adding to sprayer tank. Rinse empty container with water and drain into spray tank. Repeat twice more.

## RECOMMENDATIONS

**COTTON:** For control of boll weevil, bollworm, cutworms, fall armyworms, lygus, leafworm, grasshoppers, use  $2\frac{2}{3}$  to  $5\frac{1}{3}$  pints of this material suitably diluted with water per acre. For control of thrips and fleahoppers use 1 pint suitably diluted in water per acre.

DO NOT graze dairy animals or animals being finished for slaughter in fields treated late in the season.

**SMALL GRAINS:** (Wheat, barley, oats, rye, rice, grain sorghum) For control of armyworms and cutworms use  $1\frac{1}{2}$  to 2 quarts of this product suitably diluted and applied as indicated for cotton.

Do not ensile treated sorghum forage.

Do not apply to sorghum within 40 days of harvest. Do not apply more than one treatment after heads start to form.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**SOYBEANS:** For control of velvetbean caterpillar, bollworm, corn earworm, bean leaf beetles, armyworms, grasshoppers, blister beetle, green cloverworm, use  $1\frac{1}{2}$  to  $2\frac{2}{3}$  quarts of this material suitably diluted with water per acre. This product may be applied with either aerial or ground sprayers. With aerial application, the desired volume of insecticide concentrate should be diluted with water and applied at 3 to 4 gallons per acre.

Avoid spray drift. Do not apply less than 3 weeks before harvest. Do not feed plants treated with toxaphene, or ensilage made from treated plants, to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

DO NOT USE, POUR, SPILL OR  
STORE NEAR HEAT OR OPEN FLAME

(Contains 6 pounds Toxaphene per Gallon)

## ACTIVE INGREDIENTS:

	By Weight
Toxaphene (Chlorinated Camphene containing 67-69% Chlorine)	58.4%
Aromatic Petroleum Derivative Solvents	36.6%

INERT INGREDIENTS	5.0%
	100.0%

**WARNING**  
**KEEP OUT OF REACH OF CHILDREN**

Read Entire Label for Additional Handling Precautions

**WARNING:** May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe spray mist. Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Do not use on cucurbits. Not for use or storage in or around the home.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Apply only as specified on the label. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NET CONTENTS: . . . . . GALLONS

Manufactured By



**STANDARD SPRAY**  
**and chemical company**

Lakeland, Florida    Columbia, Alabama

## ANTIDOTE

**Internal**—Give sodium chloride 15 g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately.

**External**—Wash with soap and water.

## CONTAINER DISPOSAL

Rinse empty container with water and drain into spray tank. Repeat twice more.

Do not re-use container. Make unusable by punching holes in top and bottom. Crush. Bury in isolated area away from water supplies.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

EPA Reg. No. 3238-59