

### RESTRICTED PESTICIDE

FOR USE ONLY BY APPLICATOR LICENSED UNDER APPLICABLE STATE LAW  
Standard Brand Toxaphene-Methyl Parathion 6-3 EC is a water emulsifiable insecticidal concentrate for application in water base sprays against certain insects as directed below. Use contrary to these directions may cause serious plant, crop or personal injury.

### DIRECTIONS FOR USE

For best results premix with a small amount of water and thoroughly agitate before adding to sprayer tank. Rinse empty container with water and drain into spray tank. Repeat twice more.

**USE IN POWER-OPERATED GROUND SPRAYERS**—Use specified dosage of the spray concentrate in 5-10 gallons of water (low volume) to 50-100 gallons (high volume) per acre.

**USE IN AERIAL APPLICATION**—Use specified dosage of the spray concentrate in 3-10 gallons of water per acre.

### RECOMMENDATIONS

**COTTON:** **Early Season Program:** Against Plant Bugs and Fleahoppers, use 1 to 1½ pints per acre. Make two or three applications at 7-day intervals, starting at the two-leaf stage. For Thrips control only, use ¾ to 1 pint per acre; against Cutworms, use 3½ pints per acre. **Midseason Insects:** Overwintering Boll Weevils—use 2½ pints per acre. Plant Bugs, Aphids, Garden Webworms, and Cotton Leafworms, use 2 pints per acre. Make repeat treatments at 7-day intervals as insect counts indicate they are necessary.

**Late Season Control:** Use 3 to 4 pints per acre to control Boll Weevils, Bollworms, Garden Webworms and Aphids. Use the higher rate for heavy Bollworm outbreaks. Repeat applications at 5 to 7-day intervals as insect counts indicate they are necessary.

**CAUTION:** Do not apply within 7 days of handpicking or harvest. (Workers entering fields within 24 hours after application should wear protective clothing.) Do not graze dairy animals or animals being finished for slaughter in treated fields.

**SOYBEANS:** Against Aphids, Armyworms, Bean Leaf Beetles, Corn Earworms, Grasshoppers, Green Cloverworms, Stink Bugs and Velvetbean Caterpillars, use 2¾ pints per acre. Apply suitably diluted with water. Do not apply more than twice during the growing season.

**CAUTION:** Avoid spray drift. Do not apply less than 21 days before harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

**SMALL GRAINS (Barley, Oats, Wheat, Rice)**—Against Armyworms, apply 2 pints per acre suitably diluted with water. Do not apply within 15 days of harvest. Use limitation on Rice—crabs, crayfish and shrimp may be killed. Do not apply when they are important resources.

### CONTAINER DISPOSAL — DO NOT RE-USE

1. Rinse empty container with water and drain into spray tank. Repeat twice more.
2. Decontamination: Pour the following mixture into empty container:  
**5 gallon drums:** 2 quarts water, ½ cup caustic soda (household lye), 2 tbsp. detergent  
**30 gallon drums:** 3 gallons water, 1 lb. caustic soda, (household lye), ½ cup detergent  
**55 gallon drums:** 5 gallons water, 2 lbs. caustic soda (household lye), 1 cup detergent

Rotate container carefully until all inner surfaces are thoroughly wet. Bury wash solution at least 18 in deep in an isolated location away from water supplies. (CAUTION: Handle caustic soda (household lye) with extreme care. Do not get on skin, eyes, or on clothing. Read all directions and follow them carefully.)

3. Disposal: Punch holes in top and bottom of container; crush container and bury in isolated area away from water supplies.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

EPA Reg. No. 3238-60



DANGER—POISON—PRECAUTIONS

### CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts: spray mist or dust may be fatal if swallowed.



### CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



### CAN KILL YOU IF BREATHED

This product can kill you if vapors, spray mist or dust are breathed.

### POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply when runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.


Do not contaminate water by cleaning of equipment or disposal of wastes.

### DO NOT USE, POUR, SPILL or STORE NEAR HEAT or OPEN FLAME

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS; WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE.

POST TREATED AREA

Consult your State Agricultural Extension Service Experiment Station regarding posting treated areas



# POISON

See side panels for antidote and precautions

# TOXAPHEI 6-3

(Contains 6 lbs. T)

ACTIVE INGREDIENTS:

\*Toxaphene  
\*\*O, O-Dimethyl O-p-Aromatic Petroleum

INERT INGREDIENTS ...

\*Technical Chlorinated

\*\*Methyl Parathion

NET

# NO



Lakeland, F

STOP! READ THE LABEL

**POISON**


See side panels  
for antidote  
and precautions



**CAN KILL YOU**

**DANGER**

Keep out of reach  
of children



STANDARD BRAND

**TOXAPHENE-METHYL PARATHION  
6-3 EC INSECTICIDE**

(Contains 6 lbs. Toxaphene and 3 lbs. Methyl Parathion per gallon)

ACTIVE INGREDIENTS:	By Weight
*Toxaphene .....	52.50%
**O, O-Dimethyl O-p-Nitrophenyl Phosphorothioate .....	26.20
Aromatic Petroleum Derivative Solvents .....	14.15
INERT INGREDIENTS .....	7.15
	TOTAL 100%

- \*Technical Chlorinated Camphene (67-69% Chlorine)
- \*\*Methyl Parathion

NET CONTENTS \_\_\_\_\_ GALLONS

**NOT FOR HOME USE**

Manufactured By



Lakeland, Florida

**STANDARD SPRAY  
and chemical company**

Columbia, Alabama

USE ONLY WITH PROTECTIVE EQUIPMENT AND CLOTHING.

Do not wear clothing that has been in contact with this material. Wash clothes and shoes separately. Do not use in enclosed spaces. Do not use in areas where children or animals are present. Do not use near water. Do not use in areas where food is stored. Do not use in areas where food is prepared. Do not use in areas where food is consumed. Do not use in areas where food is stored. Do not use in areas where food is prepared. Do not use in areas where food is consumed.

Do not rub eyes or mouth with hands if you feel sore in any way. STOP work and get help right away. Call a doctor (physician) clinic or hospital immediately. Explain that the victim has been exposed to parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

Do not rub eyes or mouth with hands if you feel sore in any way. STOP work and get help right away. Call a doctor (physician) clinic or hospital immediately. Explain that the victim has been exposed to parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

**IMPORTANT!** Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

**AFTER WORK**, take off all work clothes and shoes. Shower, using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

**POISON SIGNS (Symptoms)**

Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pinpoint pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

**NOTE TO PHYSICIAN**

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Antidote—administer atropine sulfate in large doses. TWO to FOUR mg intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 20 minutes intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

