

DIRECTIONS FOR USE

Use of this material in a manner other than in accordance with this label may cause serious plant, crop, or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

PEANUTS: For the control of Cercospora Leaf Spot. Begin application of 3 to 4 pints of Sulphur-Zineb FUNGI-SPERSE in 10-20 gallons of water* to cover one acre. Begin first application approximately 40 days after planting and repeat at 7 to 10 day intervals. Increase desage rate to 3 to 4 quarts of Sulphur-Zineb FUNGI-SPERSE if disease pressure is severe. Do not feed treated peanut tops to livestock.

The amount of water used depends on type of equipment used.

CITRUS (FLORIDA): (Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines) . . . Rust Mite - use 2 gallons per 500 gallons of water in dilute sprays cr equivalent ratio for concentrate applications.

CAUTION: Some fruits and other plants are suscentible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality. During periods of high temperature sulphur may burn foliage and fruit. Do not make sulphur applications at excessive high temperatures. Do not use Sulphur-Zineb FUNGI-SPERSE with oil or within 3 weeks after oil applicarion. Do not use Sulphur-Zineb FUNGI-SPERSE in combination with copper compounds on Citrus.

Consult your State Agriculture Experiment Station or State Extension Service Specialists for complete details on the scraving program best suited to your local conditions.

EPA Est. 3238-FL-1

Lot No.

ACTIVE INGREDIENTS:

Sulphur Zineb (Zinc ethylenebisdithiocarbam INERT INGREDIENTS

(Contains 51/4 lbs. Sulphur and 1 lb. Zineb* pengallon) *(Tote! Zine as metallic from Zineb

FUNGI-SPERSE® is a registered trademark of Standard Spray and Chemical Company for premium grade insecticides and fungicides.

CAUTION > KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Do not fake internally. Avoid breathing spray mist. May cause irritation of mose, throat, or skin. Avoid contact with eyes, skin and clothing. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothes. Change to clean clothing daily. Do ast contaminate feed and feedstuffs. Do not apply to any area not specified on labely

Do not re-use samply chartainer. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.


Manufactured By






Lakeland, Fla.

Severn, N.C.



	By Weight
	44.0%
note)	8.4%
	47.6%
	100.0%

Columbia, Ala.



NET CONTENTS 5 GALLONS

DIRECTIONS FOR MIXING BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operations. DO NOT ALLOW MIXTURE TO STAND. Failure to maintain agitation will cause Sulphur-Zineb FUNGI-SPERSE® to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of Sulphur-Zineb FUNGI-SPERSE® act as a sticker on the plant, and this sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulphur in any form is a corrosive material. TO REDUCE THE EFFECT EQUIP-MENT SHOULD BE FLUSHED DAILY.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

EPA Reg. No. 3238-73