


WATER SOLUBLE PIVALYN CONCENTRATE
 PACKET IN SEALED BOXES CONTAINING 100 4-GRAM PACKETS



DIRECTIONS

To prepare Pivalyn water bait, empty contents of one packet of Pivalyn Concentrate into one quart Mason jar and fill with water. Expose this solution in amounts up to one quart in chicken founts or other convenient receptacle. Place in locations frequented by rats or mice.

IMPORTANT

Effective use of Pivalyn water baits requires CONTINUOUS exposure over a periods of days. Replenish solution before completely consumed and replace if water bait becomes dirty. Rats must drink liquid baits for 6 to 14 days. For mice, expose in smaller amounts of 1/2 to 1 pint and place at 6 to 10 ft. intervals for at least 2 weeks. Baiting should be continued until all rodent activity stops.

Where reinfestation is likely, permanent bait stations should be used and replenished as often as necessary.

WATER SOLUBLE
PIVALYN[®]
 CONCENTRATE
*Kills Rats
 and Mice*

MOTOMCO, INC.
 89 TERMINAL AVENUE
 CLARK, NEW JERSEY

CAUTION

Pivalyn, the active ingredient of this Concentrate, is an anticoagulant chemical. Therefore, Pivalyn Concentrate and Pivalyn solutions must be placed in areas inaccessible to children, pets and domestic animals or in tamper proof bait boxes. If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS:

Administration of blood transfusions combined with intravenous injections and oral doses of Vitamin K₁ is indicated.

BOX LABEL

PACKET LABEL

PACKET LABEL
 (BACK)

100 PACKETS

USDA REG. NO. 3240-3

DIRECTIONS
To prepare Pivalyn water:
1. Empty contents of
one packet into a glass.
2. Add water to the liquid
Mixture and stir with a
teaspoon until the liquid
is completely dissolved.
3. The solution is ready
for use.

IMPORTANT
Pivalyn is a potent
insecticide. It should
be used only for the
control of insects.
Do not use it for any
other purpose.
Keep it out of the
reach of children.
Do not use it in
openings where
children or animals
can reach it.
Do not use it in
places where it
can be inhaled.
Do not use it in
places where it
can be ingested.
Do not use it in
places where it
can be absorbed.
Do not use it in
places where it
can be inhaled.
Do not use it in
places where it
can be ingested.
Do not use it in
places where it
can be absorbed.

WATER SOLUBLE
PIVALYN
CONCENTRATE

CAUTION

Keep out of the reach of children.
Do not use in places where children
or animals can reach it.
Do not use in places where it
can be inhaled or ingested.
Do not use in places where it
can be absorbed.
Do not use in places where it
can be inhaled.
Do not use in places where it
can be ingested.
Do not use in places where it
can be absorbed.

**NOTE FOR
PHYSICIANS**

ACTIVE INGREDIENT:
(1) Sodium salt of 2-Pivalyl-1, 3-Indandione _____ **1.5%**
INERT INGREDIENT: _____ **98.5%**
_____ **100.0%**
(1) is equivalent to 2-Pivalyl-1, 3-Indandione _____ **1.4%**

U. S. Patent No. 2,880,132

Manufactured By

**CAUTION: Keep Out of
Reach of Children!
See side panel for
additional cautions.**

**CONTAINS 100 PACKETS, NET WEIGHT 4 GRAMS EACH. TOTAL NET WEIGHT: 14 OUNCES.
THIS PACKAGE MAY NOT BE BROKEN FOR RESALE**

and Mice

Effective use of Pivalyn water baits requires CONTINUOUS exposure over a period of days. Replenish solution before completely consumed and replace if water bait becomes dirty. Rats must drink liquid baits for 6 to 14 days. For mice, expose in smaller amounts of 1/2 to 1 pint and place at 6 to 10 ft. intervals for at least 2 weeks. Baiting should be continued until all rodent activity stops.

Where reinfestation is likely, permanent bait stations should be used and replenished as often as necessary.



MOTOMCO, INC.
88 TERMINAL AVENUE
CLARK, NEW JERSEY

bait boxes. If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS:

Administration of blood transfusions combined with intravenous injections and oral doses of Vitamin K₁ is indicated.

BOX LABEL

PACKET LABEL (FRONT)

ONE PACKET USDA Reg. No. 3246-3

WATER SOLUBLE
PIVALYN®
CONCENTRATE

ACTIVE INGREDIENT:	
(1) SODIUM SALT OF 2-PIVALYL-1, 3-INDANDIONE	1.5%
INERT INGREDIENT	98.5%
(1) EQUIVALENT TO 2-PIVALYL-1, 3-INDANDIONE	1.4%

Keep out of reach of children
MANUFACTURED BY
MOTOMCO, INC.
TERMINAL AVE., CLARK, NEW JERSEY 07066
NET WEIGHT 4.0 GRAMS

ACCEPTED

MAY 21 1971

(UNDER THE FEDERAL INSECTICIDE ACT)
FUNGICIDE AND ROSENBERG ACT
FOR ECONOMIC PEST CONTROL
ED UNDER NO. 2422-20 SUBP. 1
TO ATTACHED COMMENTS.

PACKET LABEL (BACK)

SEE LABEL ON OUTSIDE OF BOX FOR DIRECTIONS FOR PREPARING AND PLACING PIVALYN WATER BAITS.

Pivalyn, the active ingredient of this Concentrate, is an anticoagulant chemical. Therefore, Pivalyn Concentrate and Pivalyn solutions must be placed in areas inaccessible to children, pets and domestic animals or in tamper proof bait boxes. If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS: Administration of blood transfusions combined with intravenous injections and oral doses of Vitamin K₁ is indicated.

TEAR HERE