

3-10-74
MAY 23 1974

EPA Reg. No. 3240-17

MOTOMCO
WATER SOLUBLE
DIPHACINONE
RODENTICIDE CONCENTRATE

Kills Rats and Mice

Active Ingredient:

Sodium Salt of Diphacinone --	0.106%
2-Diphenylacetyl-1, 3-Indandione	99.894%
Inert Ingredients	100.0%

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

CAUTION

Keep out of reach of children!

See right panel for additional cautions

MOTOMCO, INC
89 TERMINAL AVE
CLARK, N. J. 07066
MAY 23 1974

Net Contents

ACCEPTED
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED UNDER NO. 3240-17

DIRECTIONS

How To Prepare: Dissolve one tablespoonful in one quart of water. Pour liquid bait in glass, plastic or crockery dishes or chicken founts. Note: Metal containers are not recommended.

How To Use: Place bait in locations where rats and mice feed and travel. Baiting should be maintained continuously as long as any evidence of feeding is observed. Average kill times are from 6 to 14 days, but some feeding may continue to the fourth week.

FOR RATS, minimum of one pint of liquid should be placed in several locations, such as, along walls, in corners, and where they can drink unobserved. Liquid stations should be maintained in an uninterrupted supply for at least 10 days.

FOR MICE, minimum of one pint liquid station should be placed at 8 to 10 foot intervals. Mice travel shorter distances than rats and are more difficult to control. Liquid stations should be maintained in an uninterrupted supply for at least 15 days.

IMPORTANT

Bait placements should be examined regularly and replenished before completely empty. Should baits become sour or moldy, they must be replaced. Where reinfestation is likely, permanent bait stations should be used and bait placed in them as often as necessary.

CAUTION

Keep away from children, pets, domestic animals and wildlife. Do not expose bait where humans, livestock, poultry, pets, or wildlife can drink it.

If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

Treated baits should be placed in locations not accessible to children, or pets, wildlife and domestic animals, or in tamper-proof bait boxes. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.