

T

SEP 13 1982
3240-24

FRONT PANEL

MOTOMCO

CONTRAX-P
RAT AND MOUSE BAIT
READY TO USE PLACE PAC

Active Ingredient:		
2-Pivalyl-1, 3-indandione025%
Inert Ingredients		99.975%
	TOTAL	100.000%

CAUTION
Keep out of reach of children
(See left panel for additional precautionary statements)

NET WEIGHT 1½ OZ. (43 GM)

Manufactured by
MOTOMCO, LTD.
Madison, WI 53708 U.S.A.

EPA EST. NO. 3240-WI-1 EPA REG. NO. 3240-24

RIGHT PANEL

APPLICATION DIRECTIONS

MICE: Place 1 CONTRAX-P place pac at 8 to 12 foot intervals. Bait placements should be checked frequently. At points of very high mouse activity, bait with 2 CONTRAX-P place pacs. Maintain a continuous supply of fresh bait for at least 15 days.

RATS: Place 2 to 10 CONTRAX-P place pacs per placement. The amount of bait placed at a given location should be adjusted to the size of the rat infestation. Maintain continuous supply of fresh bait for at least 10 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DISPOSAL

DO NOT REUSE CONTAINER
WRAP CONTAINER AND PUT IN TRASH COLLECTION

LEFT PANEL

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD

Keep out of lakes, streams, or ponds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway Rats, Roof Rats, and House Mice in and around homes, industrial, and agricultural buildings, and similar manmade structures.

Treated baits must be placed in areas not accessible to children, pets, domestic animals, or wildlife or in tamper proof bait boxes. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.