

POISON:

ANTIDOTE

EXTERNAL—Wash with water

INTERNAL—Drink milk, thin flour or
or starch solution, or raw egg white,
induce vomiting (tablespoon of mus-
tard in glass of warm water).

CALL PHYSICIAN

Irritating to eyes, skin and lungs.

NOTE:

This product degrades with age. Use
a test kit and increase dosage as neces-
sary to obtain required level of avail-
able chlorine.

Rinse empty container thoroughly
with water and discard it.

STORE IN COOL DARK PLACE

a product of

A & L LABORATORIES, INC.

Minneapolis, Minnesota 55405

3276-13



AL-CLO

SANITIZER

For use in MILK PLANTS,
CREAMERIES, DAIRY FARMS, ETC.

Active Ingredients:

Sodium Hypochlorite .. 8.4%

Inert Ingredients91.6%

CAUTION

**KEEP OUT OF REACH
OF CHILDREN**

(see side panel for
additional cautions)

CONTENTS: ONE GALLON

Recommended procedure for sanitizing
hard surface equipment in:

**MILK PLANTS
DAIRY FARMS
EGG BREAKING PLANTS**

Clean all surfaces thoroughly before sani-
tizing. Rinse or circulate 200 PPM (1 oz.
AL-CLO to 3¼ gallons water) for at
least 2 minutes prior to using equipment.

NOTE:

In sanitizing any equipment the sani-
tizing solution should not be rinsed
off before use.

Consult local health department for
special requirements.

U.S.D.A. REG. No. 3276-13



3276-13

ACCEPTED