

P.M. 14 3282-9

1980 10/2

**d-CON® MOUSE-PRUFE  
KILLS MICE**

**Exclusive Natural Blend**

**Kills Norway Rats, Roof Rats, and House Mice**

**Clean, Easy To Use**

**ACTIVE INGREDIENT:**

Warfarin [3- (alpha-Acetylbenzyl) -  
4-hydroxycoumarin..... 0.054%

**INERT INGREDIENTS:..... 99.946%**

**TOTAL 100.0%**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY  
STATEMENTS**

**ACCEPTED**  
**SEP 29 1992**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 3282-9

THE d-CON COMPANY, INC.  
Montvale, New Jersey 07645

EPA Reg. No. 3282-9

EPA EST. No. 3282-OH-1

NET CONTENTS 2 oz.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not is use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled and unconsumed bait as specified on this label.

**USE RESTRICTIONS** - For control of Norway rats, roof rats, and house mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

**SELECTION OF TREATMENT AREAS** - Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

### APPLICATION DIRECTIONS

Apply one 2-oz Mouse-Prufe box at 8- to 12-foot intervals in areas where mouse activity is evident. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease. If bait is not touched for 3 or 4 days and mouse activity is still present, move boxes to appropriate new locations. Apply bait in tamper-resistant bait stations wherever necessary (see "IMPORTANT").

**BEST AVAILABLE COPY**

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

### STORAGE AND DISPOSAL

**Storage:** Store only in original container, in a dry place inaccessible to children and pets.

**Disposal:** Do not reuse empty container. Securely wrap in newspaper and discard in trash.

### PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

**NOTE TO PHYSICIAN** - If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

**FOR HUMAN CASES:** If bait is eaten by humans, call a physician at once. For 24 hours emergency assistance, call your local **POISON CONTROL CENTER**. If swallowed by humans, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, injection or oral administration of Vitamin K occasionally combined with blood transfusions, are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

**FOR ANIMAL CASES:** If bait is eaten by animals, call a Veterinarian at once. If swallowed by pets, domestic animals or wildlife, this material may reduce the clotting ability of the blood and cause bleeding. Treatment is same as in human cases.

### ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not apply this product to water, swamps, bogs or potholes.

**BEST AVAILABLE COPY**