3787-9

8/2/2006

Page 133

Revised: May 23, 2006

Page: 1 of 3

.....

d-CON® Mouse-Prufe Kills Mice EPA Reg. No.: 3282-9

d-CON® Mouse-Prufe KILLS MICE

Kills House Mice

Clean, Easy to Use

ACCEPTED

AUG 2 2006

Under the Federal Innerliside, Punchide, and Michael Innerliside, as amended, for the pesitoide regulatored under 1907-9

ACTIVE INGREDIENT:

 Warfarin [3-(alpha-Acetonylbenzyl)-4-hydroxycoumarin
 0.054%

 OTHER INGREDIENTS:
 99.946%

 TOTAL:
 100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION:

SEE BACK (SIDE) PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No.: 3282-9 EPA Est. No.: 475-MS-1

NET CONTENTS: 2 OZ. (56 g)

Questions? Comments? Call 1-800-228-4722

Distributed by: Reckitt Benckiser Inc. Parsippany, NJ 07054-0224 Made in U.S.A. © RBI {year}

d-CON® Mouse-Prufe Kills Mice

EPA Reg. No.: 3282-9

Revised: May 23, 2006

Page: 2 of 3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of house mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Apply one 2-oz. Mouse-Prufe box at 8 to 12-foot intervals in areas where mouse activity is evident. Maintain uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease. If bait is not touched for 3 or 4 days and mouse activity is still present, move boxes to appropriate new locations.

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

d-CON® Mouse-Prufe Kills Mice

EPA Reg. No.: 3282-9

Revised: May 23, 2006

Page: 3 of 3

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from human, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Wear gloves if bait is handled.

Have label with	FIRST AID h you when obtaining treatment advice. You may also contact 1-800-228-4722 for emergency medical treatment information.
if swallowed	 Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.
If on skin	Wash with plenty of soap and water.
	and Veterinarian: If ingested, administer Vitamin K ₁ intramuscularly or orally, as cycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin

ENVIRONMENTAL HAZARDS: This product is toxic to mammals and birds. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets. **DISPOSAL:** If empty: Do not reuse empty container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.