

3282-65

1/29/2002

Page 178

d-CON® Mouse-Prufe II
EPA REG. NO: 3282-65

revised:01/29/2
page: 1

1.5 oz and 3.0 oz sizes - Front Label

(GOOD HOUSEKEEPING SEAL)

d-CON® Mouse-Prufe II

KILLS MICE

House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one	0.005%
INERT INGREDIENTS:	99.995%

TOTAL 100.000%

ACCEPTED

JAN 29 2002
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 3282-65

NET CONTENTS 1.5 oz (43g) 3.0 oz (85g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

278

1.5 oz and 3.0 oz sizes - Back Label

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

USE RESTRICTIONS:

THIS IS NOT A TAMPER-PROOF OR TAMPER RESISTANT BAIT STATION. DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, OR NONTARGET WILDLIFE. DO NOT USE THIS PRODUCT OUTDOORS.

This product may be used to control house mice in homes, industrial, commercial, agricultural and public buildings. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Wedges of Mouse Prufe II may be placed indoors along walls, in corners and in concealed places where mice are active and where children, nontarget mammals and birds will not be exposed to the bait.

APPLICATION DIRECTIONS:

Place wedge in horizontal position. Pull tab, lift and twist off. Space wedges at intervals of 8 to 12 feet if mouse activity is not highly localized and placements can be made in areas not accessible to children and nontarget animals. Replace wedges as their contents are consumed. Wedges not touched after 3 days should be moved to new locations appropriate for bait placements (see **USE RESTRICTIONS:**) if mouse activity persists. Maintain a supply of fresh bait for at least 15 days. Collect and dispose of all dead animals and leftover bait properly.

3 7 8

1.5 oz and 3.0 oz sizes - Back Label

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K ₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K ₁ is antidotal at doses of 10 to 20 mg (<u>not</u> mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K ₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K ₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

4 7 8

d-CONO Mouse-Prufe II
EPA REG. NO: 3282-65

revised:01/29/2
page: 4

1.5 oz and 3.0 oz sizes - Back Label

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser, Inc. Wayne, NJ 07470

EPA Reg. No. 3282-65

EPA Est. No. 475-MS-1

MADE IN THE USA

5 7 8

d-CON® Mouse-Prufe II
EPA REG. NO: 3282-65

revised:01/29/2
page: 5

6oz (170g) Outer Carton (4/1.5oz wedges) - Front Label

(GOOD HOUSEKEEPING SEAL)

Graphic - Mouse

**d-CON®
Mouse-Prufe II**

KILLS MICE

House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one0.005%

INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WGT: 6 oz (170g)

NET CONTENTS: 4/1.5 oz

Multi-(Value) Pack

4 Pack of Mouse Prufe II

SATISFACTION GUARANTEED OR YOUR MONEY BACK

6oz (170g) Outer Carton (4/1.5oz wedges) - Back Label

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

USE RESTRICTIONS:

THIS IS NOT A TAMPER-PROOF OR TAMPER RESISTANT BAIT STATION. DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, OR NONTARGET WILDLIFE. DO NOT USE THIS PRODUCT OUTDOORS.

This product may be used to control house mice in homes, industrial, commercial, agricultural and public buildings. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Wedges of Mouse Prufe II may be placed indoors along walls, in corners and in concealed places where mice are active and where children, nontarget mammals and birds will not be exposed to the bait.

APPLICATION DIRECTIONS:

Place wedge in horizontal position. Pull tab, lift and twist off. Space wedges at intervals of 8 to 12 feet if mouse activity is not highly localized and placements can be made in areas not accessible to children and nontarget animals. Replace wedges as their contents are consumed. Wedges not touched after 3 days should be moved to new locations appropriate for bait placements (see **USE RESTRICTIONS:**) if mouse activity persists. Maintain a supply of fresh bait for at least 15 days. Collect and dispose of all dead animals and leftover bait properly.

6oz (170g) Outer Carton (4/1.5oz wedges) - Back Label

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID

Have label with you when obtaining treatment advice.

**If swallowed
by human**

- Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

**If swallowed
by pet or animal**

- Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

8 7 8

d-CONO Mouse-Prufe II
EPA REG. NO: 3282-65

revised:01/29/2
page: 8

6oz (170g) Outer Carton (4/1.5oz wedges) - Back Label

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser, Inc. Wayne, NJ 07470

EPA Reg. No. 3282-65

EPA Est. No. 475-MS-1

MADE IN THE USA
