

d-CON
GENERATION 2
Kills
Rats & Mice

From d-CON
 America's #1
 Rat Killer

Advanced New
 Anticoagulant
 Formula

Proved up to 50
 Times More Active
 (Weight for Weight)
 than Warfarin
 Against Rats
 and Mice

Can Kill
 in One Feeding

Satisfaction
 Guaranteed or
 Your Money Back

The d-CON Company Inc
 Subsidiary of Sterling Drug
 225 Summit Avenue Montvale
 New Jersey 07645

© 1971 M. V. ...

d-CON
GENERATION 2
Kills
Rats & Mice

IMPROVED ANTICOAGULANT FORMULA

*RATS AND MICE WILL DIE WITHIN 4 OR 5 DAYS

8 READY-TO-USE BAIT PACKS

***Can Kill in
 one feeding**

KILLS WARFARIN RESISTANT NORWAY RATS
 AND WARFARIN RESISTANT HOUSE MICE

CAUTION: Keep out of reach of children.

READ ADDITIONAL PRECAUTIONARY STATEMENTS
 ON BACK PANEL

ACTIVE INGREDIENT: Brodifacoum
 3-(3,4-bromophenyl)-1,1-biphenyl-4-yl
 1,2,3,4-tetrahydro-1-naphthyl-1-yl
 hydroxy-2H-1-benzopyran-2-one
 INERT INGREDIENTS

TOTAL 100.000

NET WT 16 OZ (453g)
 NET CONTENTS 8 2.0 OZ (57g)

d-CON
GENERATION 2

Kills Rats & Mice

IMPROVED ANTICOAGULANT FORMULA

KILLS WARFARIN RESISTANT NORWAY RATS
 AND WARFARIN RESISTANT HOUSE MICE

d-CON Kills Rats & Mice

GENERATION 2

DIRECTIONS FOR USE

d-CON Generation 2 can kill in one feeding* when used as directed. Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice

General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Use Restrictions

For control of rats and house mice in homes, industrial and commercial buildings, place bait packs in areas inaccessible to children, pets, domestic animals and wildlife

For indoor use only. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food

Selection of Treatment Areas

Place packs in or near points where you have seen rat or mice signs such as droppings, runways, burrows or gnawing marks. Once areas requiring baiting have been identified, proceed as follows:

APPLICATION DIRECTIONS

1. Place one ready-to-use bait pack in each potential feeding location. Pack should not be opened, as this protects bait from moisture and spoilage. Be sure to place packs in dark, out of the way locations, never out in the open

2. Cover the pack with a bait box or board

3. For best results, allow pack to remain undisturbed for at least 2 days. If high Rodent activity persists, place a new pack if original one has been removed or partially eaten. Place 2 new packs where original was entirely eaten. Continue checking every 2 days, adding new packs as needed until a complete lack of feeding is noted

4. If initial inspection shows no consumption, allow pack to remain 3 more days. If no feeding occurs after 5 days, move pack to a new location

For complete control, continue baiting for at least 15 days. If you are in an area where there is danger of reinfestation from adjoining property, permanent bait stations should be maintained. Stations should be checked periodically

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding

NOTE TO PHYSICIAN:

If ingested, administer Vitamin K₁ intramuscularly

or orally, as indicated in bishydroxycoumarin overdoses. Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg) and should be given and repeated as necessary (based on monitoring of prothrombin times)

ENVIRONMENTAL HAZARDS:

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds


DISPOSAL:

Do not reuse empty container. Wrap container and put in trash collection



ACCEPTED

NOV 15 1981



ENVIRONMENTAL HAZARDS.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

DISPOSAL.

Do not reuse empty container. Wrap container and put in trash collection.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Refer to label on package for full use directions.

NET WT 2.0Z (57g)
EPA Reg. No. 188-M-1
EPA Est. No. 280718P

TOTAL 100,000*

0.005%
99.995%

INERT INGREDIENTS

1,2,3,4-tetrahydro-1-naphthalenyl-4-hydroxy-2H-1-benzopyran-2-one

ACTIVE INGREDIENT Brodifacoum

3-(3,4-dibromophenyl) biphenyl-4-yl

1,2,3,4-tetrahydro-1-naphthalenyl-4-hydroxy-2H-1-benzopyran-2-one

READ ADDITIONAL PRECAUTIONARY STATEMENTS ON BACK PANEL

CAUTION: Keep out of reach of children

AND WARFARIN-RESISTANT HOUSE MICE

KILLS WARFARIN-RESISTANT NORWAY RATS

RODENTICIDE BAIT PACK (PELLETS)

Rats & Mice

Kills

Can Kill in one feeding.

APPROVED AND CONTROLLED BY EPA

GENERATION 2

D-GON

PRECAUTIONARY STATEMENTS, HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION:

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN:

If ingested, administer Vitamin K intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Vitamin K₁ is antidote, at doses of 10 to 20 mg (not mg/kg) and should be given and repeated as necessary (based on monitoring of prothrombin times).