

Front Panel
(Box)

d-CON
PELLETS GENERATION II

4 READY-TO-USE BAIT TRAYS
ADVANCED ANTICOAGULANT FORMULA
*Can kill in one feeding

Kills Rats & Mice
KILLS WARFARIN-RESISTANT NORWAY RATS
AND WARFARIN-RESISTANT HOUSE MICE.

*RATS AND MICE WILL DIE WITHIN 4 OR 5 DAYS

GOOD HOUSEKEEPING SEAL:
Good Housekeeping promises a limited warranty to
consumers. Replacement or refund if defective.

Flavor attractive to rats and mice
Palatable Formulation

ACTIVE INGREDIENT: Brodifacoum
3-[3-(4'bromo-[1,1'-biphenyl]-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-
hydroxy-2H-1-benzopyran-2-one0.005%
INERT INGREDIENTS 99.995%

TOTAL 100.000%

NET WT. 12 OZ. (340g)
NET CONTENTS 4/3.0 OZ. (85g)

CAUTION: Keep out of reach of children
may be harmful or fatal if swallowed
Read additional precautionary statements
on side panel

BOTTOM OF BOX
d-CON Kills Rats & Mice
Pellets Generation II

KILLS WARFARIN-RESISTANT NORWAY RATS
AND WARFARIN-RESISTANT HOUSE MICE

328266
This product is a restricted use pesticide
as defined under EPA Reg. 263.

91 1989

ACCOUNTING
INVOICE

ASB-66
PMA 14

1/9

BACK PANEL
(BOX)

d-CON^R Kills
PELLETS GENERATION II Rats &
ADVANCED ANTICOAGULANT FORMULA Mice

d-CON Pellets Generation II can kill in one feeding when used as directed. Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway Rats, Roof Rats and House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Pellets Generation II may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-resistant bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Place trays in or near points where you have seen rat or mice signs such as droppings, runways, burrows, or gnawing marks. Once areas requiring baiting have been identified proceed as follows:

APPLICATION DIRECTIONS:

1. Place one ready-to-use bait tray in each potential feeding location. Place trays in dark, out-of-the way locations, where rodents are likely to find them. Do not place trays in the open or in any location exposed to children or nontarget mammals or birds.



BACK PANEL
(BOX) (CONTINUED)

For House Mice: Use only one tray per location.

For Norway and Roof Rats: Start with one tray per location. Add trays, up to a maximum of four per location, if you determine that there is high rat activity at the location.

2. If bait must be applied in areas accessible to children or nontarget animals, trays must be placed in secured, sturdy, tamper-resistant bait stations which deny these nontarget species access to the bait.

3. For best results, leave trays undisturbed for at least two days after placement. However, if contents have been scattered or mostly consumed sooner, replace trays. Clean up spilled bait. Check bait every two days, replacing trays as needed or until signs of rodent activity cease. Trays not fed from for five consecutive days may be relocated as needed. When there are no longer any signs of rodent activity, dispose of trays properly.

For complete control, continue baiting for at least 15 days. If you are in an area where there is danger or reinfestation from adjoining property, permanent bait stations should be maintained. Stations should be checked periodically.

NOTE: Underline denotes bold face type



b/c

BACK PANEL (Continued)
(BOX)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait. IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24 hour emergency assistance, call your local Poison Control Center. IF BAIT IS EATEN BY ANIMALS OR PETS: Call your local veterinarian.

NOTE TO PHYSICIAN AND VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL:

Store in original container in areas inaccessible to small children and pets. Bait that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.

4/19

LEFT SIDE PANEL
(BOX)

d-CON

READY MIXED GENERATION II
ADVANCED ANTICOAGULANT FORMULA

Kills Rats & Mice

Kills
Warfarin-Resistant
Norway Rats and
Warfarin-Resistant
House Mice

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assumes the risk of any such use.

SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

EPA Reg. No. 3282-
EPA Est. No. 3282-OH-1 or 2393-WI-1
See Bottom of Box

Satisfaction
Guaranteed or
Your Money Back

5/11

RIGHT SIDE PANEL
(BOX)

d-CON

PELLETS GENERATION II
ADVANCED ANTICOAGULANT FORMULA

Kills

Rats & Mice

From d-CON
America's #1
Rat Killer

Advanced
Anticoagulant
Formula

Can Kill
in One Feeding

Satisfaction
Guaranteed or
Your Money Back

The d-CON Company Inc.
Subsidiary of Sterling Drug
225 Summit Avenue
Montvale, New Jersey 07645 U.S.A.

b/g

TOP PANEL
(BOX)

d-CON PELLETS GENERATION II

Kills Rats and Mice

4 Ready-To-Use Bait Trays

b11.

BACK PANEL
(BAIT TRAY)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Place bait in areas not accessible to children, pet, domestic animals or wildlife or in tamper resistant bait boxes.

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait. IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24 hour emergency assistance, call you local Poison Control Center. IF BAIT IS EATEN BY ANIMALS OR PETS: Call your local veterinarian.

NOTE TO PHYSICIAN AND VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

See outer box for complete Directions for Use.

THE d-CON COMPANY, INC.
Subsidiary of Sterling Drug, Inc., 225 Summit
Avenue, Montvale, NJ 07645 U.S.A.

FRONT PANEL
(BAIT TRAY)

d-Con_R

PELLETS GENERATION II

Kills Rats & Mice

READY-TO-USE BAIT TRAY

ADVANCED ANTICOAGULANT FORMULA

*Call kill in one feeding

KILLS WARFARIN-RESISTANT NORWAY RATS

KILLS WARFARIN-RESISTANT HOUSE MICE

CAUTION: KEEP OUT OF REACH OF CHILDREN
May be harmful or fatal if swallowed.
Read additional precautionary statements on side
panel.

ACTIVE INGREDIENT: Brodifacoum
3-[3-(4'-bromo-[1,1'biphenyl]-4-yl)-
1,2,3,4-tetrahydro-1-napthalenyl]-4-hydroxy-2H-1-
benzoy-an-2-one.....0.005%
INERT INGREDIENTS99.995%

TOTAL 100.000%

*First dead rodents will appear four or five days after
treatment begins.

NET WT. 3.0 Oz. Bait Tray (85 g)

3282-6C
MAY 1980
EPA
b/b