

PM 04 3282-66
01/05/99

Page 1 of 7

12 oz. BOX FRONT LABEL

(GOOD HOUSEKEEPING SEAL)

d-CON® PELLETS GENERATION II

KILLS RATS AND MICE

4 READY-TO-USE
BAIT FILLED TRAYS

Flavor attractive to rats and mice

CAN KILL IN ONE FEEDING*

*Rats and mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

ACCEPTED
JAN 5 1999
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 3282-66

Keep out of reach of children.
CAUTION: May be harmful or fatal if swallowed.
Read additional precautionary statements on
back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-[1,1'-	
biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-	
hydroxy-2H-1-benzopyran-2-one	0.005%
INERT INGREDIENTS	99.995%
	TOTAL 100.000%

NET CONTENTS 4/3.0 OZ. (85g) NET WT. 12 OZ. (340g)

BACK PANEL

d-CON PELLETS GENERATION II

KILLS RATS AND MICE

Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control Norway Rats, Roof Rats and House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Pellets Generation II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:")

APPLICATION DIRECTIONS: The amount of bait contained in the trays in this box is unlikely to be sufficient to kill more than 3 to 12 rats. More than one box may be needed to control larger infestations.

To Control Norway and Roof Rats: Place 1 - 4 bait trays per placement. Space placements at intervals of 15 - 30 feet in infested areas. If trays are not fed from for 5 consecutive days, relocate them to other places where rodent activity exists and where placements consistent with the requirements of this label can be made. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Open tray and apply 1/4 - 1/2 oz. (1 - 2 level tablespoons) of bait at 8 to 12 foot intervals in infested areas. Larger placements (up to 2 ounces) may be needed at points of extremely high mouse activity. Maintain a supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. Replace contaminated or spoiled bait immediately if rodent activity is still evident. Collect and dispose of all dead animals and leftover bait properly.

To prevent reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait **IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE.** For 24 hour emergency assistance, call your local Poison Control Center. **IF BAIT IS EATEN BY ANIMALS OR PETS:** Call your local veterinarian.

NOTE TO PHYSICIAN AND VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL:

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap contained and any unused bait in newspaper and discard in trash.

12 OZ. BOX RIGHT SIDE PANEL

d-CON®
PELLETS GENERATION II

KILLS RATS AND MICE

Kills
Warfarin-Resistant
Norway Rats and
Warfarin-Resistant
House Mice

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assumes the risk of any such use.

SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

EPA Reg. No. 3282-66
EPA Est. No. 475-MS-1; 2393-WI-1

SATISFACTION
GUARANTEED OR
YOUR MONEY BACK

12 OZ. BOX LEFT SIDE PANEL

d-CON®
PELLETS GENERATION II

KILLS RATS AND MICE

CAN KILL IN ONE FEEDING*

*Rats and mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins

MADE IN THE USA

d-CON BAITES:
KILLING RATS AND MICE IN AMERICA
FOR 50 YEARS

SATISFACTION
GUARANTEED OR
YOUR MONEY BACK

Distributed by:
Household Products Division
Reckitt & Colman Inc.
Wayne, NJ 07474

3 OZ. BAIT TRAY - FRONT PANEL

d-CON®
PELLETS GENERATION II


KILLS RATS AND MICE

Ready-To-Use Bait Tray

Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice

CAN KILL IN ONE FEEDING*

*Rats and mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.



Keep out of reach of children.
CAUTION: May be harmful or fatal if swallowed.
Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one0.005%
INERT INGREDIENTS99.995%
TOTAL 100.000%

NET WT. 3 OZ. BAIT TRAY (85g)

3 OZ. BAIT TRAY - BACK PANEL

d-CON® PELLETS GENERATION II

KILLS RATS AND MICE

Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label on the outer package before using this product. It is illegal to sell these bait trays individually.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Place bait in areas not accessible to children, pets, domestic animals or wildlife or in tamper-resistant bait boxes.

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait. **IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE.** For 24 hour emergency assistance, call your local Poison Control Center. **IF BAIT IS EATEN BY ANIMALS OR PETS:** Call your local veterinarian.

NOTE TO PHYSICIAN AND VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-

hydroxy-2H-1-benzopyran-2-one0.005%

INERT INGREDIENTS99.995%

TOTAL 100.000%

Distributed by:

Household Products Division

Reckitt & Colman Inc., Wayne, NJ 07474

EPA Reg. No. 3282-66

EPA Est. No. 475-MS-1; 2393-WI-1