

d-CON
LIM-N8
RAT KILLER
IN READY-TO-USE BAIT PACKS
(Pellets)

Read the entire label carefully and follow the directions for use. Do not use in areas where children, domestic animals, or pets are present. Do not use in areas where food is prepared or served. Do not use in areas where food is stored. Do not use in areas where food is consumed. Do not use in areas where food is prepared or served. Do not use in areas where food is stored. Do not use in areas where food is consumed.


d-CON
LIM-N8
RAT KILLER
IN READY-TO-USE BAIT PACKS
(Pellets)

Kills Rats and Mice. Improved Anticoagulant Formula.
Can kill in one feeding with first dead rodents appearing four or five days after treatment begins.

CAUTION Keep out of reach of children. Do not use in areas where children, domestic animals, or pets are present. Do not use in areas where food is prepared or served. Do not use in areas where food is stored. Do not use in areas where food is consumed.

NET CONTENTS 10.43 gm (1/5 OZ) BAIT PACKS

In Accordance with the provisions of the Federal Food, Drug, and Cosmetic Act, this product is labeled as follows:
1/16/96



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION May be harmful or fatal if swallowed. Keep away from children, domestic animals, and pets. Wash hands after handling.
IF BAIT IS EATEN BY HUMANS: Call a physician at once. For a hospital emergency, consult your local Poison Control Center.
IF BAIT IS EATEN BY ANIMALS OR PETS: Call a veterinarian at once. For a hospital emergency, consult your local Poison Control Center (215) 333-3611.

NOTE TO PHYSICIAN AND VETERINARIAN
This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If prothrombin activity, prothrombin time and administration of Vitamin K, are indicated, as in paragraph from user dose of dicumarol (D-50) (50 mg/kg).
FOR HUMAN CASES: Vitamin K₁ is indicated at doses of 10 to 20 mg (oral mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.
FOR ANIMAL CASES: Vitamin K₁ is indicated at 5 mg/kg orally. In severe cases, blood transfusions may be necessary.
ENVIRONMENTAL HAZARDS
This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to beds of prey and mammals that eat out of any body of water.

d-CON
LIM-N8
RAT KILLER
IN READY-TO-USE BAIT PACKS (Pellets)
KILLS RATS AND MICE
Satisfaction Guaranteed or your money back

DIRECTIONS FOR USE:
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Use Restrictions: For control of Norway Rats, Flood Rats, House Mice in and around homes, industrial, commercial, agricultural and public buildings. LIM-N8 may also be used in and around transport vehicles (ships, trains, aircraft) and related part of terminal buildings. Do not use in sewers. This product must be placed in tamperproof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.
Selection of Treatment Areas: Undermost areas where rats and mice will most likely feed and consume the bait. Generally these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places, between floors and walls or in locations where rodents or their signs have been observed.

APPLICATION DIRECTIONS
For Norway Rats: Apply 2 to 6 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.
For House Mice: Apply 1 bait pack at intervals of 8 to 12 feet per placement. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.
Rats and Mice: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to disposal paragraph. To prevent reinfestation eliminate food, water and harborage as much as possible. If reinfestation does occur repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL
Store in original container in areas inaccessible to small children and pets. Bait that cannot be used according to label instructions must be disposed of according to applicable federal, state or local procedures.

Distributed by THE d-CON COMPANY INC.
Subsidiary of Sterling Drug Inc. 225 SUMMIT AVENUE MONTVALE, NJ 07045



BEST AVAILABLE COPY