11/21/2001

Please read instructions on reverse be 05-31-98	ore completing form.			rom Approved, Or	VIB 140. 2070-0000, Approvar expires			
Ω EPA Fin	es	☐ Regist	tration	OPP Identifier Number				
En'	rironmental Prote	ction Agenc	y 🛛 Amen	dment	289122			
	Washington, DC	20460	☐ Other					
Application for Pesticide - Section I								
Company/Product Number			Product Manager		3. Proposed Classification			
	2-74		eacock					
Company/Product (Name) d-CON LIM-N8 Rat Killer	PM#			None Restricted				
Name and Address of Applicant (Reckitt Benckiser Inc.	Include ZIP Code)				ith FIFRA Section 3(c)(3)			
1655 Valley Road		(b)(i),			n composition and labeling			
Wayne, New Jersey 074	1	Reg. No.	NOTIFICATION	M				
		1	Product Name					
Check if this is a new a	nddress	11000	or Name	MAX -				
		Secti	on - II					
Amendment - Explain below.			Final printed la	bels in response to	Agency letter dated			
Resubmission in response to	Agency letter dated	·	Me Too" Applic	ation				
Notification - Explain below.			Other - Explain					
Explanation: Use additiona			ion I and Section I	l.)				
Response to EPA Letter D								
Change the Product's Bas	ic Name from d-C	ON LIM-N8 R	at Killer to d-CON	Bait Pellets II.				
		Socti	 on - III					
Material This Product Will Be Page	ockaged In:		011 - 111					
Child-Resistant Packaging	Unit Packaging		Water Soluble Pag	ckaging	2. Type of Container			
Yes*			Yes		Metal			
⊠ No	│ □ No		⊠ No		Plastic			
	If "Yes"	No. per	If "Yes"	No. per	Glass			
*Certification must	Unit Packaging wg	container 20	Package wgt.	container	Paper			
be submitted	1.002	20			Other (Specifiy)			
Location of Net Contents Inform		(s) Retail Contai		,	Label Directions			
Label 🔀 Con	z and 20/1.0o	Z	On Label					
				ng accompanying product				
6. Manner in Which Label is Affixed to Product Lithograph Other Print on carton								
☐ Paper glued ☐ Stenciled								
		Section	on - IV					
1. Contact Point (Complete items of	irectly below for identific	ation of individua	to be contacted, if nec	essary, to process	this application)			
Name Title Telephone No. (Incl								
Sean McNea	Sr. Regulatory Affairs Specialist			Code) .: 973-686-7390				
	Certific	cation		6. Date Application				
					Received			
acknowledge that any knowingly fa under applicable law.	lse or misleading statem	ent may be punis	hable by fine or impriso	onment or both	(Stamped)			
2. Signature	3. Title			3 / 34 /				
Jan Mail	Sr. Regulatory Affairs Specialist			(2) 新井行を必要(3) (2) (2) (2)				
4. Typed Name	5. Date							
Sean McNe	October 16, 2001			r fr. i				

1.0 oz - Bait Pack Front Panel (for Mice)

(GOOD HOUSEKEEPING SEAL)

page:

revised: 10/16/01

d-CON® Bait Pellets II

KILLS MICE

Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-

hydroxy-2H-1-benzopyran-2-one0.005%

TOTAL 100.000%

NET WT. 1.0 OZ. (28g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

revised: 10/16/01

d-CON® Bait Pellets II EPA REG. NO: 3282-74

EPA REG. NO: 3282-74 page: 2

1.0 oz - Bait Pack Back Label (for Mice)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

revised:10/16/01

page: 3

1.0 oz - Bait Pack Back Panel (for Mice)

APPLICATION DIRECTIONS:

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by mice for 5 consecutive days, they may be relocated to other places where mice activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

revised:10/16/01

age: 4

1.0 oz - Bait Pack Front Panel (for Mice)

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID								
IF SWALLOWED	• This material may reduce the clotting ability of the							
By HUMANS or	blood and cause bleeding. • Call a physician or Poison Control Center							
ANIMALS or PETS	immediately for treatment advice.							
	HOT LINE NUMBER							

Questions? Comments? Call 1-800-228-4722

NOTE TO PHYSICIAN AND VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K_1 are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K_1 is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K_1 is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K_1 should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

revised:10/16/01

page: 5

1.0 oz - Bait Pack Back Panel (for Mice)

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap container and any unused bait in newspaper and discard in trash.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

EPA Reg. No. 3282-74 EPA Est. No. 475-MS-1 ·

MADE IN THE USA

revised:10/16/01

page: 6

1.0 oz Bait Pack Front Panel (for Mice and Rats)

(GOOD HOUSEKEEPING SEAL)

d-CON® Bait Pellets II

KILLS RATS AND MICE

Rats and mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-

TOTAL 100.000%

NET WT. 1.0 oz. (28g) Bait Pack

SATISFACTION GUARANTEED OR YOUR MONEY BACK

revised:10/16/01 page: 7

1.0 oz Bait Pack Back Label (for Mice and Rats)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

revised:10/16/01

d-CON® Bait Pellets II EPA REG. NO: 3282-74

EPA REG. NO: 3282-74 page: 8

1.0 oz Bait Pack Back Label (for Mice and Rats)

APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Rats and Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rodents for 5 consecutive days, they may be relocated to other places where rodent activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

revised: 10/16/01

d-CON® Bait Pellets II EPA REG. NO: 3282-74

PA REG. NO: 3282-74 page: 9

1.0 oz Bait Pack Back Label (for Mice and Rats)

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID									
IF SWALLOWED	• This material may reduce the clotting ability of the								
	blood and cause bleeding.								
By HUMANS or	• Call a physician or Poison Control Center								
ANIMALS or PETS	immediately for treatment advice.								

HOT LINE NUMBER

Questions? Comments? Call 1-800-228-4722

NOTE TO PHYSICIAN AND VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K_1 are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K_1 is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K_1 is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K_1 should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap container and any unused bait in newspaper and discard in trash.

revised: 10/16/01

1.0 oz Bait Pack Back Label (for Mice and Rats)

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

EPA Req. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA

EPA REG. NO: 3282-74 page: 11

20/1.0 oz Bait Pack Outer Box - FRONT PANEL

(GOOD HOUSEKEEPING SEAL)

Graphic - Rodent

revised:10/16/01

d-CON® Bait Pellets II

KILLS RATS AND MICE

Rats and mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-

TOTAL 100.000%

NET WT. 20/1.0 oz (28g) bait packs (20 place packs)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

revised:10/16/01

d-CON® Bait Pellets II EPA REG. NO: 3282-74

EPA REG. NO: 3282-74 page: 12

20/1.0 oz Bait Pack Outer Box - BACK LABEL

Kills Rats and Mice

Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals
to rodenticides. To help prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

d-CON® Bait Pellets II revised:10/16/01 EPA REG. NO: 3282-74 page: 13

20/1.0 oz Bait Pack Outer Box - BACK LABEL

APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Rats and Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rodents for 5 consecutive days, they may be relocated to other places where rodent activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

revised: 10/16/01

d-CON® Bait Pellets II EPA REG. NO: 3282-74

EPA REG. NO: 3282-74 page: 14

20/1.0 oz Bait Pack Outer Box - BACK LABEL

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

	F	I	R	S	T	AID
--	---	---	---	---	---	-----

IF SWALLOWED

• This material may reduce the clotting ability of the blood and cause bleeding.

By HUMANS or PETS

• Call a physician or Poison Control Center immediately for treatment advice.

HOT LINE NUMBER

Questions? Comments? Call 1-800-228-4722

NOTE TO PHYSICIAN AND VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K_1 are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K_1 is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K_1 is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K_1 should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap container and any unused bait in newspaper and discard in trash.

revised: 10/16/01 page: 15

20/1.0 oz Bait Pack Outer Box - BACK LABEL

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

EPA Req. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA