

3282-74

1/29/2002

1/22

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised: 01/29/2
page: 1

1.0 oz - Bait Pack Front Panel (for Mice)

(GOOD HOUSEKEEPING SEAL)

**d-CON®
Bait Pellets II**

KILLS MICE

House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'-
biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-
hydroxy-2H-1-benzopyran-2-one0.005%

INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WT. 1.0 OZ. (28g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

ACCEPTED

JAN 29 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act, we warrant for the products

3282-74

1.0 oz - Bait Pack Back Label (for Mice)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "**IMPORTANT:**").

1.0 oz - Bait Pack Back Panel (for Mice)

APPLICATION DIRECTIONS:

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by mice for 5 consecutive days, they may be relocated to other places where mice activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

4/22

1.0 oz - Bait Pack Front Panel (for Mice)

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID

Have label with you when obtaining treatment advice.

**If swallowed
by human**

- Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

**If swallowed
by pet or animal**

- Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

5/22

1.0 oz - Bait Pack Back Panel (for Mice)

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA

1.0 oz Bait Pack Front Panel (for Mice and Rats)

(GOOD HOUSEKEEPING SEAL)

**d-CON®
Bait Pellets II**

KILLS RATS AND MICE

Rats and House Mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-(1,1'-
biphenyl)-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-
hydroxy-2H-1-benzopyran-2-one0.005%
INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WT. 1.0 oz. (28g) Bait Pack

SATISFACTION GUARANTEED OR YOUR MONEY BACK

1.0 oz Bait Pack Back Label (for Mice and Rats)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "**IMPORTANT:**").

8/22

1.0 oz Bait Pack Back Label (for Mice and Rats)

APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Rats and Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rodents for 5 consecutive days, they may be relocated to other places where rodent activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

9/22

1.0 oz Bait Pack Back Label (for Mice and Rats)

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K ₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K ₁ is antidotal at doses of 10 to 20 mg (<u>not</u> mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases; blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K ₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K ₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

10/22
revised: 01/29/2
page: 10

1.0 oz Bait Pack Back Label (for Mice and Rats)

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA

6/1.0 oz Bait Pack Outer Box - FRONT PANEL

(GOOD HOUSEKEEPING SEAL)

Graphic - Rodent

**d-CON®
Bait Pellets II**

KILLS MICE

House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one0.005%
INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WT. 6/1.0 oz (28g) bait packs (6 place packs)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

6/1.0 oz Bait Pack Outer Box - BACK LABEL

Kills Mice
Kills Warfarin-Resistant House Mice

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible.

13/22

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised: 01/29/2
page: 13

Use tamper-resistant bait stations wherever necessary (see
"IMPORTANT:").

6/1.0 oz Bait Pack Outer Box - BACK LABEL

APPLICATION DIRECTIONS:

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by mice for 5 consecutive days, they may be relocated to other places where mice activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

6/1.0 oz Bait Pack Outer Box - BACK LABEL

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID

Have label with you when obtaining treatment advice.

**If swallowed
by human**

- Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

**If swallowed
by pet or animal**

- Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised: 01/29/2
page: 16

6/1.0 oz Bait Pack Outer Box - BACK LABEL

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA

20/1.0 oz Bait Pack Outer Box - FRONT PANEL

(GOOD HOUSEKEEPING SEAL)

Graphic - Rodent

**d-CON®
Bait Pellets II**

KILLS RATS AND MICE

Rats and House Mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT: Brodifacoum 3-[3-(4'-bromo-{1,1'-
biphenyl}-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-
hydroxy-2H-1-benzopyran-2-one0.005%
INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WT. 20/1.0 oz (28g) bait packs (20 place packs)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

20/1.0 oz Bait Pack Outer Box - BACK LABEL

Kills Rats and Mice

Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative

d-CON® Bait Pellets II

EPA REG. NO: 3282-74

revised: 01/29/2

page: 19

19/22

food as possible. Use tamper-resistant bait stations wherever
necessary (see "**IMPORTANT:**").

20/1.0 oz Bait Pack Outer Box - BACK LABEL

APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Rats and Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rodents for 5 consecutive days, they may be relocated to other places where rodent activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

20/1.0 oz Bait Pack Outer Box - BACK LABEL

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID

Have label with you when obtaining treatment advice.

If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.

NOTE TO PHYSICIAN AND VETERINARIAN

The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised:01/29/2
page: 22

20/1.0 oz Bait Pack Outer Box - BACK LABEL

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA
