

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

June 30, 2003

305
12

S-631533

Reckitt Benckiser Inc.
1655 Valley Road
P.O. Box 943
Wayne, NJ 07474-0943

Attention: Mr. Sean McNear

Subject d-Con ® Bait Pellets II
EPA Reg. No. 3282-74
Your amended application of February 24, 2003
Revised Label

Purpose The purpose of your submission under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) is A) to add two additional package sizes: 1) 4, 1 oz place packs and 2) 5, 1 oz place packs and B) to add two claims to your label.

A. Packaging The 1.0-oz placepacks would not provide enough bait for a single placement at the labeled rate for controlling commensal rats. Therefore, it would be illegal to sell such placepacks individually (or in groups of less than 4) if rat directions appeared on the labeling.

B. Label The label submitted with the above letter, under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) is acceptable, provided that you make the following changes and submit one (1) copy of final printed labeling, including graphics, to us before you release product for shipment. A copy of the stamped label is enclosed for your records.

1. The only "Graphic - Rodent" that may appear is the one accepted in our January 26, 2003, letter. For mouse-only presentations, you may only use a house mouse graphic.

B. Label-continued

- 2. The proposed claims, "Mice (will) eat through (the) (place) (pack) pack." and "Ready-to-use place packs" are acceptable. The "Mice(& Rats)" version of the first of these claims may only be used on packages that bear claims for controlling commensal rats as well as house mice.
- 3. Packages of 4-14 of the 1-oz placepacks would have to bear appropriate qualifying statements advising customers and users of the limited numbers of rats that could be expected to be killed with the amounts of bait provided in such packages. At present, you are not marketing such packaging combinations, based upon our information.
- 4. Replace "Toxicity Category I Pesticide" with "Have label with you when seeking treatment advice."

Existing stocks

Stocks of existing labels may be used for eighteen (18) months.

Consequence for non-compliance

If these conditions are not complied with, the registrations will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Questions

If you have questions about this letter, please contact me at 703-305-5407 (by phone), 703-305-6596 (by fax), or peacock,dan@epa.gov (by E-Mail).

Sincerely yours,



Daniel B. Peacock, Biologist
Insecticide-Rodenticide Branch
Registration Division (7504C)

Letter location

Disk 107:A:\Brodifacoum\3282-74, revised label, add 2 labels, add 2 package sizes,6-30-2003wpd.wpd,June 5, 2003

3/22

d-CONO Bait Pellets II
EPA REG. NO: 3282-74

revised: 02/24/03
page: 1

1.0 oz - Bait Pack (for Mice)

(GOOD HOUSEKEEPING SEAL)

Mice (will) eat through (the) (place) (bait) pack

d-CONO Bait Pellets II **ACCEPTED** with **COMMENTS**
in EPA Letter Dated

KILLS MICE JUN 30 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under FIFRA, No.

House Mice may consume a lethal dose 38282-74 with first
dead mice appearing 4 or 5 days after feeding begins.

38282-74

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-(1,1'-biphenyl)-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-
benzopyran-2-one.....0.005%


INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT. 1.0 OZ. (28g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

Ready-to-use (1) place pack



4/22

1.0 oz - Bait Pack (for Mice)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").



5/22

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised: 02/24/03
page: 3


1.0 oz - Bait Pack (for Mice)

APPLICATION DIRECTIONS:

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by mice for 5 consecutive days, they may be relocated to other places where mice activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.



6/22

d-CONO Bait Pellets II
EPA REG. NO: 3282-74

revised: 02/24/03
page: 4

1.0 oz - Bait Pack (for Mice)


PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none"> • Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice. • Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none"> • Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
<p>This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (<u>not</u> mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.</p>	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.



d-CONO Bait Pellets II
EPA REG. NO: 3282-74

revised:02/24/03
page: 5

1.0 oz - Bait Pack (for Mice)

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.


Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

Questions? Comments? Call 1-800-228-4722

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA



8/22

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised:02/24/03
page: 6

1.0 oz Bait Pack (for Mice and Rats)

(GOOD HOUSEKEEPING SEAL)

Mice (& Rats) (will) eat through (the) (place) (bait) pack

**d-CON®
Bait Pellets II**

KILLS RATS AND MICE

Rats and House Mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-
benzopyran-2-one.....0.005%


INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT. 1.0 oz. (28g) Bait Pack

SATISFACTION GUARANTEED OR YOUR MONEY BACK

Ready-to-use (1) place pack



9/22

1.0 oz Bait Pack (for Mice and Rats)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "**IMPORTANT:**").

10/22

d-CONO Bait Pellets II
EPA REG. NO: 3282-74

revised: 02/24/03
page: 8

1.0 oz Bait Pack (for Mice and Rats)


APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Rats and Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rodents for 5 consecutive days, they may be relocated to other places where rodent activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.



1.0 oz Bait Pack (for Mice and Rats)

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
<p>This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (<u>not</u> mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.</p>	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

12/22
revised:02/24/03
page: 10

1.0 oz Bait Pack (for Mice and Rats)

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.


Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

Questions? Comments? Call 1-800-228-4722

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA



13/22

d-CONO Bait Pellets II
EPA REG. NO: 3282-74

revised:02/24/03
page: 11

4,5,6/1.0 oz Bait Pack Outer Box

(GOOD HOUSEKEEPING SEAL)

Graphic - Rodent

Mice (will) eat through (the)(place)(bait) pack

**d-CONO
Bait Pellets II**

KILLS MICE

House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-(1,1'-biphenyl)-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-
benzopyran-2-one.....0.005%


INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT 4/1.0 oz (28g) bait packs (4 place packs)
 5/1.0 oz (28g) bait packs (5 place packs)
 6/1.0 oz (28g) bait packs (6 place packs)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

Ready-to-use (4) (5) (6) place packs



4,5,6/1.0 oz Bait Pack Outer Box

Kills Mice

Kills Warfarin-Resistant House Mice

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.



15/22

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised: 02/24/03
page: 13

4, 5, 6/1.0 oz Bait Pack Outer Box


SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

APPLICATION DIRECTIONS:

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by mice for 5 consecutive days, they may be relocated to other places where mice activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.



16/22

4,5,6/1.0 oz Bait Pack Outer Box

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none"> • Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice. • Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none"> • Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
<p>This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (<u>not</u> mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.</p>	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

17/22

d-CONO Bait Pellets II
EPA REG. NO: 3282-74

revised:02/24/03
page: 15

4,5,6/1.0 oz Bait Pack Outer Box

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.


Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

Questions? Comments? Call 1-800-228-4722

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA



18/22

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised: 02/24/03
page: 16

20/1.0 oz Bait Pack Outer Box

(GOOD HOUSEKEEPING SEAL)

Graphic - Rodent

Mice (& Rats) (will) eat through (the) (place) (bait) pack

**d-CON®
Bait Pellets II**

KILLS RATS AND MICE

Rats and House Mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-
benzopyran-2-one.....0.005%


INERT INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT 20/1.0 oz (28g) bait packs (20 place packs)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

Ready-to-use (20) place packs



20/1.0 oz Bait Pack Outer Box

Kills Rats and Mice

Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.



20/1.0 oz Bait Pack Outer Box

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or signs of rodents have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").


APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Rats and Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rodents for 5 consecutive days, they may be relocated to other places where rodent activity exists and where bait placements consistent with the requirements of this label can be made.

Collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.



20/1.0 oz Bait Pack Outer Box

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
<p>This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.</p>	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

22/22

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

revised:02/24/03
page: 20

20/1.0 oz Bait Pack Outer Box

NOTICE TO BUYER AND USER: Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Distributed by: Reckitt Benckiser Inc. Wayne, NJ 07470

Questions? Comments? Call 1-800-228-4722

EPA Reg. No. 3282-74

EPA Est. No. 475-MS-1

MADE IN THE USA

