

3282-74

5/13/2009

10 of 19

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Heather Bjornson
Regulatory Consultant to Reckitt Benckiser
Technology Sciences Group, Inc
1150 18th St, N.W.
Washington, D.C. 20036

MAY 13 2009

Dear Ms. Bjornson:

Subject: Labeling Amendment to d-Con Bait Pellets II
EPA Registration No. 3282-74
Submission Date: February 11, 2009

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. If you have any questions regarding this label, please contact Jennifer Gaines at (703) 305-5967.

This product is not consistent with the Agency's May 28, 2008, "Risk Management Decision for Ten Rodenticides." EPA anticipates cancellation of those existing products that are not consistent with the Risk Mitigation Decision to occur no later than June 4, 2011. In the meantime, EPA is approving new registrations and amendments to existing registrations of rodenticide bait products, on a time-limited basis, so long as the registrations do not present greater risks of unreasonable adverse effects than existing products. Accordingly, this amendment is approved only subject to the condition that the registration shall expire on June 4, 2011. This condition of registration may be amended in conjunction with amendments to conform to the Risk Mitigation Decision. Absent such amendments, product released for shipment by the registrant after June 4, 2011, would be in violation of FIFRA.

Sincerely yours,

A handwritten signature in black ink, appearing to read "John Hebert".

John Hebert
Product Manager (07)
Insecticide-Rodenticide Branch
Registration Division (7505P)

2 of 19

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

Page 1 of 18
May 13, 2009

1.0 oz – Bait Pack (for Mice)


(GOOD HOUSEKEEPING SEAL)

House Mice (will) eat through (the)(place)(bait) pack

**d-CON®
Bait Pellets II**

KILLS MICE

CAN KILL IN ONE FEEDING*



* House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-

1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-

benzopyran-2-one.....0.005%

OTHER INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT: 1.0 OZ. (28 g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

Call 1-800-228-4722

Important: For directions for use and first aid instruction in Spanish, please call 1-866-648-1819

Ready-to-use

(1) (4) (5) (6) (ready-to-use) place packs

1.0 oz – Bait Pack (for Mice)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may only be used to control House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where house mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where their signs have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

4 of 19

1.0 oz – Bait Pack (for Mice)

APPLICATION DIRECTIONS:

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high house mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by house mice for 5 consecutive days, they may be relocated to other places where house mice activity exists and where bait placements consistent with the requirements of this label can be made.

While wearing gloves, collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

1.0 oz – Bait Pack (for Mice)**PRECAUTIONARY STATEMENTS:****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
<p>This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (<u>not</u> mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.</p>	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

1.0 oz – Bait Pack (for Mice)

STORAGE AND DISPOSAL

Do not contaminate water, food, feed by storage or disposal.

PESTICIDE STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE TO BUYER AND USER: To the extent consistent with applicable law, Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Questions? Comments? Call 1-800-228-4722

Distributed by: Reckitt Benckiser Inc. Parsippany, NJ 07054(-xxxx)

EPA Reg. No.: 3282-74

EPA Est. No.: 475-MS-1

MADE IN THE USA

Important: For directions for use and first aid instruction in Spanish, please call 1-866-648-1819

1.0 oz Bait Pack (for Mice and Rats)

(GOOD HOUSEKEEPING SEAL)

House Mice (& Norway and Roof Rats)(will) eat through (the)(place)(bait) pack

**d-CON®
Bait Pellets II**

KILLS RATS AND MICE

CAN KILL IN ONE FEEDING*

*Norway rats, Roof Rats and House Mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-

1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-

benzopyran-2-one.....0.005%

OTHER INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT: 1.0 OZ (28 g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

Call 1-800-228-4722

Important: For directions for use and first aid instruction in Spanish, please call 1-866-648-1819

Ready-to-use

(1) (4) (5) (6) (ready-to-use) place packs

1.0 oz Bait Pack (for Mice and Rats)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may only be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or house mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where their signs have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

1.0 oz Bait Pack (for Mice and Rats)

APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Norway Rats, Roof Rats, and House Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rats or house mice for 5 consecutive days, they may be relocated to other places where thier activity exists and where bait placements consistent with the requirements of this label can be made.

While wearing gloves, collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

1.0 oz Bait Pack (for Mice and Rats)

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
<p>This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.</p>	

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed by storage or disposal.

PESTICIDE STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

1.0 oz Bait Pack (for Mice and Rats)

NOTICE TO BUYER AND USER: To the extent consistent with applicable law, Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Questions? Comments? Call 1-800-228-4722

Distributed by: Reckitt Benckiser Inc. Parsippany, NJ 07054(-xxxx)

EPA Reg. No.: 3282-74

EPA Est. No.: 475-MS-1

MADE IN THE USA

4,5,6/1.0 oz Bait Pack Outer Box

(GOOD HOUSEKEEPING SEAL)

Graphic – Rodent

House Mice (will) eat through (the)(place)(bait) pack

**d-CON®
Bait Pellets II**

KILLS MICE

CAN KILL IN ONE FEEDING*

* House Mice may consume a lethal dose in one feeding with first dead mice appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-
benzopyran-2-one.....0.005%

OTHER INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT:	4/1.0 OZ. (28 g)	4 OZ. (112 g)
	5/1.0 OZ. (28 g)	5 OZ. (140 g)
	6/1.0 OZ. (28 g)	6 OZ. (172 g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK

Call 1-800-228-4722

Important: For directions for use and first aid instruction in Spanish, please call 1-866-648-1819

Ready-to-use

(4)(5)(6) (ready-to-use) place packs

4.5,6/1.0 oz Bait Pack Outer Box

Kills Mice

Kills Warfarin-Resistant House Mice

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may only be used to control House Mice in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

4,5,6/1.0 oz Bait Pack Outer Box

SELECTION OF TREATMENT AREAS: Determine areas where house mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where their signs have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

APPLICATION DIRECTIONS:

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high house mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by mice. If bait packs are not opened by house mice for 5 consecutive days, they may be relocated to other places where house mice activity exists and where bait placements consistent with the requirements of this label can be made.

While wearing gloves, collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none">• Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none">• Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K ₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K ₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K ₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K ₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.	

4,5,6/1.0 oz Bait Pack Outer Box

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed by storage or disposal.

PESTICIDE STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE TO BUYER AND USER: To the extent consistent with applicable law, Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Questions? Comments? Call 1-800-228-4722

Distributed by: Reckitt Benckiser Inc. Parsippany, NJ 07054(-xxxx)

Important: For directions for use and first aid instruction in Spanish, please call 1-866-648-1819

EPA Reg. No. 3282-74
EPA Est. No. 475-MS-1

MADE IN THE USA

20/1.0 oz Bait Pack Outer Box

16 of 19

d-CON® Bait Pellets II
EPA REG. NO: 3282-74

Page 15 of 18
May 13, 2009

(GOOD HOUSEKEEPING SEAL)

Graphic – Rodent

House Mice (& Norway and Roof Rats)(will) eat through (the)(place)(bait) pack

d-CON®
Bait Pellets II

KILLS RATS AND MICE

CAN KILL IN ONE FEEDING*

*Norway rats, Roof Rats and House Mice may consume a lethal dose in one feeding with first dead rodents appearing 4 or 5 days after feeding begins.

Keep out of reach of children.

CAUTION: May be harmful or fatal if swallowed.

Read additional precautionary statements on back panel.

ACTIVE INGREDIENT:

Brodifacoum 3-[3-(4'-bromo-{1,1'-biphenyl}-4-yl)-
1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-
benzopyran-2-one.....0.005%

OTHER INGREDIENTS:99.995%

TOTAL 100.000%

NET WEIGHT: 20/1.0 OZ. (28 g)
20 OZ. (560 g)

SATISFACTION GUARANTEED OR YOUR MONEY BACK
Call 1-800-228-4722

Important: For directions for use and first aid instruction in Spanish, please call 1-866-648-1819

Ready-to-Use
(20) (Ready-to-use) place packs

20/1.0 oz Bait Pack Outer Box

Kills Rats and Mice

Kills Warfarin-Resistant Norway Rats and Warfarin-Resistant House Mice

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: This product may only be used to control House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Bait Pellets II may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or house mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where their signs have been seen. Remove as much alternative food as possible. Use tamper-resistant bait stations wherever necessary (see "IMPORTANT:").

APPLICATION DIRECTIONS:

To Control Norway and Roof Rats: Use 4-16 bait packs per placement. Space placements at intervals of 15-30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days.

To Control House Mice: Use one bait pack per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 2 bait packs may be used at points of very high house mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

For Rats and Mice: Check bait locations every 2 days for signs of feeding. Replenish bait if more than 1/2 of the original amount has been removed by rodents. If bait packs are not opened by rodents for 5 consecutive days, they may be relocated to other places where their activity exists and where bait placements consistent with the requirements of this label can be made.

While wearing gloves, collect and dispose of all dead animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. This material may reduce the clotting ability of the blood and cause bleeding. Keep away from humans, domestic animals, and pets. Wash hands after handling bait.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed by human	<ul style="list-style-type: none"> • Call a Poison Control Center, doctor, or 1-800-228-4722 immediately for treatment advice. • Do not induce vomiting unless told to do so by the poison control center or doctor.
If swallowed by pet or animal	<ul style="list-style-type: none"> • Call a veterinarian immediately for treatment advice.
NOTE TO PHYSICIAN AND VETERINARIAN	
<p>This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). FOR HUMAN CASES: Vitamin K₁ is antidotal at doses of 10 to 20 mg (<u>not</u> mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. FOR ANIMAL CASES: Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases blood transfusions may be necessary.</p>	

20/1.0 oz Bait Pack Outer Box

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed by storage or disposal.

PESTICIDE STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE TO BUYER AND USER: To the extent consistent with applicable law, Buyer and User assume all responsibility for safety and risk associated with any use not in accordance with directions. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

Questions? Comments? Call 1-800-228-4722

Distributed by: Reckitt Benckiser Inc. Parsippany, NJ 07054(-xxxx)

EPA Reg. No. 3282-74
EPA Est. No. 475-MS-1

MADE IN THE USA

Important: For directions for use and first aid instruction in Spanish, please call 1-866-648-1819