

ACCEPTED  
 COMMENTS.

**DIRECTIONS FOR USE — Read entire label**

Add Staffel's 56% Malathion Emulsifiable concentrate to water at the Chart Rate and mix, to form a white milky emulsion. For foliage difficult to wet add a suitable spreader-sticker at the recommended rate. APPLICATION — Use a pressure spray and thoroughly spray entire plant. Be sure to COVER UNDERSIDE OF LEAVES.

PLANT	PEST	Per Gal. Water	Per 100 Gal.	Interval Last Appl. to Harvest
<b>ORNA- MENTALS</b> Flowers Vines Shrubs	Lace bugs, Euonymus Scale, Scurfy scale, Azalea Scale, Pine Leaf Scale, Magnolia Scale; Fletcher, Florida Red, Juniper, Monterey Pine, Soft, Pine Needle, and Wax Scale. Black Scale Crawlers. Aphids, Mealybugs, Spider Mites, Whitefly, Four-lined Leaf Bug, Japanese beetle adult, Potato Leafhopper, Tarnished Plant Bug, Thrips, Rose Leafhopper, European Pine Shoot Moth, Leaf Miner, Bagworms, Tent Caterpillar.	2 Tsp	2 Pts	
<b>VEGETABLES</b> Beans	Mexican Bean Beetle, Leafhoppers, Spider Mites, Japanese Beetle, Aphids, Cucumber Beetle	2 Tsp	2 Pts	1 Day
Tomatoes	Spider Mites, Aphids	2 Tsp	2 Pts	1 Day
Lettuce Leaf Head	Leafhoppers, Aphids, Mites	2 Tsp	2 Pts	14 Days 7 Days
<b>TREE FRUITS</b> Peaches	Oriental Fruit Moth, Plum Curculio, Aphids, Japanese Beetles, Terrapin & Cottony Peach Scale Apply at petal fall and every 10 to 14 days thereafter until control is achieved.	2 Tsp	2 Pts	7 Days
Pears	Mealybugs, Mites, Pear Psylla, Coddling Moth, Plum Curculio Apply at petal fall and every 10 to 14 days thereafter until control is achieved.	2 Tsp	2 Pts	1 Day

**DESTROY CONTAINER WHEN EMPTY.**

**WARNING**

Avoid prolonged breathing of spray mist. Avoid contact with skin. Wash thoroughly after handling. Do not contaminate food or feed products. Keep container closed and away from heat, sparks and open flame. Highly toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes of containers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

**8 OUNCES  
56%  
MALATHION**

(Premium Grade)  
**Active Ingredients:**  
 Malathion\* .....56%  
 Aromatic Petroleum  
 Solvent .....32%  
 Inert

**DIRECTIONS FOR USE — Read entire label**

Use only on plants listed on label. Do not use on plants not listed. Do not use on plants listed unless specifically directed. Do not use on plants listed unless specifically directed. Do not use on plants listed unless specifically directed. Do not use on plants listed unless specifically directed.

PLANT	PEST	Per Gal Water	Per 100 Gal Water	Interval Last Appl to Harvest
<b>ORNAMENTALS</b>				
Flowers	Whitefly, Spider Mite, Thrips, Aphid	2	2	14
Vines	Whitefly, Spider Mite, Thrips, Aphid	2	2	14
Shrubs	Whitefly, Spider Mite, Thrips, Aphid	2	2	14
<b>VEGETABLES</b>				
Beans	Bean Beetle, Aphid	2	2	14
Tomatoes	Tomato Pinworm, Aphid	2	2	14
Lettuce	Lettuce Miner, Aphid	2	2	14
<b>TREE FRUITS</b>				
Peaches	Peach Leaf Curler, Aphid	2	2	14
Pears	Pear Leaf Curler, Aphid	2	2	14

**DESTROY CONTAINER WHEN EMPTY**

**8 OUNCES  
56%  
MALATHION**

(Premium Grade)  
Active Ingredients:  
Malathion 56%  
Inert Ingredients 44%

**Inert Ingredients .....12%**  
\*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate.

**WARNING  
KEEP OUT OF REACH  
OF CHILDREN.  
HARMFUL IF  
SWALLOWED, INHALED  
OR ABSORBED  
THROUGH THE SKIN.**

**See back panel for  
additional cautions.  
FERD STAFFEL CO.  
San Antonio, Texas 78298  
USDA Reg. No. 3286-5**

**WARNING**

Avoid prolonged breathing of spray mist. Avoid contact with skin. Wear thoroughly after handling. Do not contaminate food or food products. Keep container closed and away from heat, sparks and open flame. Highly toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes of container. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**