

3286-7

6-27-1978

Staffel's

sevin*5 dust

FS1-61A

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl

N-methylcarbamate) 5%

INERT INGREDIENTS 95%

*Trade-mark of the Union Carbide Corp. for the active ingredient 1-naphthyl (N-methylcarbamate.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for additional cautions
Read all directions before using

EPA Reg. No. 3286-7

EPA Est. No. 3286-TX-1

Net Contents 1 lb.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL DIRECTIONS

Do not contaminate water, food, or feed by storage or disposal. Pesticide, spray mixture, or rinsate, that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of in an incinerator and landfill approved for pesticide containers, or bury in a safe place.

VEGETABLE, VINE & FIELD CROPS:

Apply when air is calm, using ground equipment, aircraft, or hand duster. Unless otherwise specified, apply at the first sign of infestation and repeat 7-10 day intervals when needed to maintain control. Observe use limits given for specific crops.

BEANS—Leafhoppers, lygus bugs, Mexican bean beetle, stink bugs. Apply 11 to 15 oz. per 1000 sq. ft., treating upper and lower leaf surfaces. Do not feed treated vines to dairy animals or meat animals.

CORN—Corn earworm. Apply 11 to 15 oz. per 1000 sq. ft. starting when the first silks appear. Repeat at 2-3 day intervals throughout the silking period. Direct duster outlets toward ear zone to insure thorough coverage of each silk. May be applied up to harvest on sweet corn ears for human consumption. To avoid excessive residues on fodder or forage, do not apply within 7 days of harvest. Reapply immediately after rain. Applications of Sevin to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

COTTON—Early season control of cotton fleahopper, thrips. Apply 7½ to 11 oz. per 1000 sq. ft. (20 to 30 lbs. per acre). Mid and late-season control of: Boll weevil, boll worm, cotton leaf perforator, cotton leafworm, lygus bug, stink bugs. Apply 15 oz. per 1000 sq. ft. (20 to 40 lbs. per acre). Repeat at 5-7 day intervals schedule. Salt-marsh caterpillar: Apply 1 lb. 2 oz. per 1000 sq. ft. (50 lbs. per acre). Do not graze treated fields or feed trash to dairy or meat animals.

CUCUMBER, SUMMER SQUASH—Flea beetle, melonworm, pickleworm, spotted cucumber beetle, squash bugs. Apply 7½ oz. per 1000 sq. ft.

EGGPLANT, PEPPER, TOMATO—Fall armyworm, flea beetle, tomato fruitworm. Apply 11 to 15 oz. per 1000 sq. ft. European corn borers, stink bugs, tomato hornworm. Apply 1 lb. 6 oz. to 1 lb. 14 oz. per 1000 sq. ft.

GRAPES (Western U.S.)—Grape leaf folder, grape leafhopper. For early season control use 11 oz. per 1000 sq. ft. and 11 to 15 oz. for summer control. For the control of first brood grape leaf folder, apply when maximum number of eggs have hatched. For leafhopper control, apply at first sign of infestation.

POTATOES—Colorado potato beetle, flea beetle, leafhopper. Apply 11 to 15 oz. per 1000 sq. ft. May be applied up to harvest on the above vegetable, vine and field crops.

STRAWBERRIES—Meadow spittlebug, strawberry leaf roller. Apply 15 oz. per 1000 sq. ft. May be applied up to 3 days before harvest.

ORNAMENTALS—Bagworm, blister beetle, boxelder bug, certain aphids, certain leaf miners, elm leaf beetle, exposed thrips, flea beetle, Japanese beetle, leafhoppers, leaf rollers, orange tortrix, periodical cicada, plant bugs, scale insects, tent caterpillars. Apply 11 to 15 oz. per sq. ft. Repeat when necessary to maintain control. In test, Sevin has been used at recommended rates without damage to andromeda, arbor vitae, ash, aster, azalea, bamboo, beech, begonia, birch boxwood, camellia, carnation, ceanothus cedar, chinese lem, chrysanthemum, cypress, dahlia, delphinium, dogwood, elm, eunonymus, fir, fuchsia, gladiolus, hemlock, hickory, holly, hornbeam, hydrangea, inkberry, iris, juniper, laurel, lilac, locust, magnolia, maidenhair fern, maple, marigold, nasturtium, oak, orchid, ornamental mulberry, petunia, pine, poplar, privet, rhododendron, rose, rose of Sharon, sassafras, spirea, verbena, viburnum, willow, wisteria, yew, zinnia.

FOR FLEAS, TICKS, MITES, LICE:

FOR FLEA AND TICK CONTROL ON DOGS AND CATS—Rub thoroughly in the skin of the animal, avoiding the eyes. Dust sleeping quarters, runways, pens, etc. weekly. DO NOT use on puppies or kittens under 4 weeks of age.

FOR MITES, LICE, FLEAS AND TICKS ON CHICKENS—One pound is enough dust for 100 chickens. Can also be used on pheasant, pigeons and partridge. Do not repeat applications to poultry within four weeks. Dust chicken house using one pound per 40 square feet. Do not house birds in treated poultry houses within seven days of slaughter. Do not dust birds within 7 days of slaughter. Do not dust nest nor eggs.

FERD STAFFEL COMPANY, San Antonio, TX 78298

ACCEPTED
JUN 27 1978
Federal Insecticide,
Fungicide, and Rodenticide Act,
under the Pesticide
Reg. No. 3286-7

11