

Best Available Copy

HOW TO USE STAFFEL'S SEVIN 50 WETTABLE:

Please Read Directions Carefully

DIRECTIONS FOR USE

Use 2 level tablespoons SEVIN-50 in 1 gallon water, unless otherwise directed. Mix proper amount SEVIN in a little water to make a paste then add balance of water to make a gallon of spray mixture.

Make first application when insects or their damage first appear. Repeat applications at 7 to 10 day intervals to maintain effective control, unless otherwise directed. Keep spray mixture well agitated.

ORNAMENTALS

ROSES, OTHER FLOWERS, SHRUBS, SHADE TREES. To control June Bugs, apple aphid, bagworm, blister beetles, boxwood leaf miner, elm leaf aphid and beetle, flea beetles, gypsy moth, lace bugs, leaf hoppers, leaf rollers, mealy bugs, webworms, oak leaf miners, periodical cicada (17 year locust), plant bugs, puss caterpillars, psyllids, rose aphid and slug, scale, insects, tent caterpillars, exposed thrips, and Japanese Beetles. DO NOT USE ON BOSTON IVY.

(For larger quantities use 2 lbs. Sevin-50W per 100 gallons water).

LAWNS

To control Chinch bugs, ants, earwigs, fall armyworm, fleas, leathoppers, millipedes, mosquitoes, and webworms (lawn moths). Use 1 level cupful SEVIN-50 in 15 to 20 gallons water and apply evenly to 500 square feet of established lawn area.

VEGETABLES

Can be used on day of harvest unless otherwise indicated.

BEANS. For Mexican bean beetle, bean leaf beetle, cutworms, cucumber beetles, flea beetles, Japanese beetles, leathoppers, velvet bean caterpillar, armyworms, corn earworm, stinkbugs, tarnished plant bug, lima bean pod borer, typhus and stink bugs.

CARROTS, LETTUCE. 2 to 4 tablespoons. Do not use on lettuce within 14 days of harvest. For flea beetles, leathoppers, harlequin bug, six-spotted leathopper (later yellows vector), armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bugs, stink bugs. For best control of small cutworms, armyworms on lettuce, use every 5 to 7 days after heads form.

CORN. Use 2 to 4 tablespoons. For Corn earworm, apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals. Re-apply immediately after rain. corn rootworm adults, European corn borer (apply thoroughly into the whorls when plant is about knee high). Repeat at weekly intervals until control is achieved. fall armyworm, flea beetles, Japanese beetles, Sap beetles, leathoppers.

CUCUMBERS, SQUASH, PUMPKINS, MELONS. For pickleworm, melon worm, cucumber beetles, flea beetles, leathoppers, squash bugs.

TOMATOES, POTATOES, EGGPLANT, PEPPERS. Use 2 to 4 tablespoons. For Colorado potato beetle, flea beetles, leathoppers, European corn borer,

fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA. For corn earworm and stink bugs.
CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI. Use 2 to 4 tablespoons. Do not use within 3 days of harvest. For flea beetles, harlequin bug, armyworms, corn earworm. For control of small cabbage loopers use every 5 to 7 days.

FRUITS

Use up to 1 day before harvest unless otherwise directed.

APPLES, PEARS. 1 tablespoon for Codling moth, apple mealybug, green apple aphid, white apple leathopper, 2 tablespoons for apple maggot, bagworm, Eastern tent caterpillar, European apple sawfly, eye-spotted bud moth, red-banded and fruit tree leaf rollers, green fruitworm, Japanese beetle, pear psylla, periodical cicada (17 year locust), plum curculio, rosy and wooly apple aphids, apple and pear rust mites, pear leaf blister mites, tarnished plant bug, tentiform leaf miners, Lecanium, oystershell and San Jose scales.

APRICOTS, PEACHES, NECTARINES. For Cucumber beetles, European earwig, Japanese beetle, June beetles, lesser peach tree borer, peach twig borer, periodical cicada, red-banded and fruit-tree leaf rollers, tarnished plant bug, tussock moth, Lecanium, San Jose and olive scales, Codling moth, oriental fruit moth, plum curculio. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not use 3 days before harvest on apricots, nectarines.

CITRUS. For citrus cutworm, fruit tree leaf roller, Western tussock moth, black scale, brown soft scale, citricola scale, yellow scale. Do not use within 5 days of harvest.

GRAPES. Can be used on day of harvest. For European fruit Lecanium, grape leaf folder, grape leathopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles, red-banded leaf roller.

STRAWBERRIES. Use up to 1 day before harvest. For cutworms, fruitworms, Japanese beetle, leathoppers, spittlebug, strawberry leaf roller, strawberry weevil.

PLUMS. Use up to 1 day before harvest. For mealy plum aphid, cherry fruit fly, eye-spotted bud moth, red-banded and fruit tree leaf roller, Japanese beetle, lesser peach twig borer, prune leathopper, brown soft scale, Lecanium, oystershell and San Jose scales, peach twig borer, Eastern tent caterpillar, Codling moth, tussock moth. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

NUTS

Can be used on day of harvest.

PECANS. For pecan nut case bearer. Do not use after husks split.

SEVIN* 50 Wettable

FOR CONTROL OF INSECTS ON VEGETABLES, FRUITS, PECANS, ORNAMENTALS, FLOWERS, LAWNS & TREES

Active Ingredient	By Wt.
Carbaryl (1-naphthyl-N-methylcarbamate)	50%
Inert Ingredients	50%

WARNING!

KEEP OUT OF REACH OF CHILDREN

U.S. Patent No. 2,903,473

CAUTION!

CAUTION — HARMFUL IF INHALED OR SWALLOWED. Do not breathe Dust. Do not take internally. Avoid Breathing of Spray.

SKIN CONTACT MAY BE HARMFUL. Avoid Skin Contact. Wash thoroughly after using.
FOR PHYSICIANS: Carbaryl is a moderate, reversible, Cholinesterase Inhibitor. Atropine is antidote.

AVOID CONTAMINATION OF FOOD.

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to direct treatment or residues on crops. For more information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: The Ferd Staffel Company makes no warranty, express or implied, concerning this material, except that it conforms to the chemical description on the label. Neither The Ferd Staffel Company, nor its supplier of this chemical, nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate

Put up by THE FERD STAFFEL COMPANY SAN ANTONIO, TEXAS 78206
12 OUNCES NET WEIGHT

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

USDA Reg. No. 3286-9

3286-9

03/04/1971

1/2

ACCEPTED
MAR 4 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC INSECT PESTICIDES
ED UNDER NO. ~~5286-9~~ SUBJECT
TO ATTACHED COMMENTS.

2/2