

HOW TO USE STAFFEL'S SEVIN 50 WETTABLE:

Please Read Directions Carefully

DIRECTIONS FOR USE

Use 2 level tablespoons SEVIN-50 in 1 gallon water, unless otherwise directed. Mix proper amount SEVIN in a little water to make a paste; then add balance of water to make 1 gallon of spray mixture.

Make first application when insects or their damage first appear. Repeat applications at 7 to 10 day intervals to maintain effective control, unless otherwise directed. Keep spray mixture well agitated.

ORNAMENTALS

ROSES, OTHER FLOWERS, SHRUBS, SHADE TREES. To control June Bugs, apple aphid, bagworm, blister beetles, boxwood leaf miner, elm leaf aphid and beetle, flea beetles, gypsy moth, lace bugs, leaf hoppers, leaf rollers, mealy bugs, webworms, oak leaf miners, periodical cicada (17 year locust), plant bugs, puss caterpillars, psyllids, rose aphid and slug, scale, insects, tent caterpillars, exposed thrips, and Japanese Beetles. **DO NOT USE ON BOSTON IVY.**

(For larger quantities use 2 lbs. Sevin-50W per 100 gallons water).

LAWNS

To control Chinch bugs, ants, earwigs, fall armyworm, fleas, leafhoppers, millipedes, mosquitoes, sod webworms (lawn moths). Use 1 level cupful SEVIN-50 in 15 to 20 gallons water and apply evenly to 500 square feet of established lawn area.

VEGETABLES

Can be used on day of harvest unless otherwise indicated.

BEANS. For Mexican bean beetle, bean leaf beetle, cutworms, cucumber beetles, flea beetles, Japanese beetles.

TOMATOES, POTATOES, EGGPLANT, PEPPERS. Use 2 to 4 tablespoons. For Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA. For corn earworm and stink bugs.

CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI. Use 2 to 4 tablespoons. Do not use within 3 days of harvest. For flea beetles, harlequin bug, armyworms, corn earworm. For control of small cabbage loopers use every 5 to 7 days.

FRUITS

Use up to 1 day before harvest unless otherwise directed.

APPLES, PEARS. 1 tablespoon for Codling moth, apple mealybug, green apple aphid, white apple leafhopper. 2 tablespoons for apple maggot, bagworm, Eastern tent caterpillar, European apple sawfly, eye-spotted bud moth, red-banded and fruit tree leaf rollers, green fruitworm, Japanese beetle, pear psylla, periodical cicada (17 year locust), plum curculio, rosy and wooly apple aphids, apple and pear rust mites, pear leaf blister mites, tarnished plant bug, tentiform leaf miners, Lecanium, oystershell and San Jose scales.

APRICOTS, PEACHES, NECTARINES. For Cucumber beetles, European earwig, Japanese beetle, June beetles, lesser peach tree borer, peach twig borer, periodical cicada, red-banded and fruit-tree leaf rollers, tarnished plant bug, tussock moth, Lecanium, San Jose and olive scales, Codling moth, oriental fruit moth, plum curculio. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not use 3 days before harvest on apricots, nectarines.

CITRUS. For citrus cutworm, fruit tree leaf roller, Western

ACCEPTED

HOW TO USE STAFFEL'S SEVIN 50 WETTABLE:

Please Read Directions Carefully

DIRECTIONS FOR USE

Use 2 level tablespoons SEVIN-50 in 1 gallon water unless otherwise directed. Mix proper amount SEVIN in a little water to make a paste, then add balance of water to make a gallon of spray mixture. Make first application when insects or their damage first appear. Repeat applications at 7 to 10 day intervals to maintain effective control, unless otherwise directed. Keep spray mixture well agitated.

ORNAMENTALS

ROSES, OTHER FLOWERS, SHRUBS, SHADE TREES. To control: June Bugs, apple aphid, bagworm, blister beetles, boxwood leaf miner, elm leaf aphid and beetle, flea beetles, gypsy moth, lace bugs, leaf hoppers, leaf rollers, mealy bugs, webworms, oak leaf miners, periodical cicada (17 year locust), plant bugs, puss caterpillars, psyllids, rose aphid and slug, scale, insects, tent caterpillars, exposed thrips, and Japanese Beetles. **DO NOT USE ON BOSTON IVY.**

(For larger quantities use 2 lbs. Sevin-50W per 100 gallons water.)

LAWNS

To control: Chinch bugs, ants, earwigs, fall armyworm, fleas, leafhoppers, millipedes, mosquitoes, sod webworms, lawn moths. Use 1 level cup of SEVIN-50 in 15 to 20 gallons water and apply evenly to 1,000 square feet of established lawn area.

VEGETABLES

Can be used on day of harvest unless otherwise indicated.

BEANS. For Mexican bean beetle, bean leaf beetle, cutworms, cucumber beetles, flea beetles, Japanese beetles,

TOMATOES, POTATOES, EGGPLANT, PEPPERS. Use 2 to 4 tablespoons. For Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bug, stink bugs.

OKRA. For corn earworm and stink bugs.

CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI. Use 2 to 4 tablespoons. Do not use within 3 days of harvest. For Flea beetles, harlequin bug, armyworms, corn earworm. For control of small cabbage looters use every 5 to 7 days.

FRUITS

Use up to 1 day before harvest unless otherwise directed.

APPLES, PEARS. 1 tablespoon for Codling moth, apple mealybug, green apple aphid, white apple leafhopper. 2 tablespoons for apple maggot, bagworm, Eastern tent caterpillar, European apple sawfly, eye-spotted bud moth, red-banded and fruit tree leaf rollers, green fruitworm, Japanese beetle, pear psylla, periodical cicada (17 year locust), plum curculion, rosy and wooly apple aphids, apple and pear rust mites, pear leaf blister mites, tarnished plant bug, tentiform leaf miners, Lecanium, oystershell and San Jose scales.

APRICOTS, PEACHES, NECTARINES. For Cucumber beetles, European earwig, Japanese beetle, June beetles, lesser peach tree borer, peach twig borer, periodical cicada, red-banded and fruit-tree leaf rollers, tarnished plant bug, tussock moth, Lecanium, San Jose and olive scales, Codling moth, oriental fruit moth, plum curculion. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not use 3 days before harvest on apricots, nectarines.

CITRUS. For citrus cutworm, fruit tree leaf roller, Western

ACCEPTED