

ACCEPTED

MAY 30 1978

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3286-9

Staffel's

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid breathing of spray. SKIN CONTACT MAY BE HARMFUL. Avoid skin contact. Wash thoroughly after using. AVOID CONTAMINATION OF FOOD AND FEED.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM Oplatea, or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARD

BEEF CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative agricultural extension service. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until 1 week application or to take other equally effective precautions.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL DIRECTIONS

Do not reuse container. Destroy when empty.

Use 2 level tablespoons SEVIN-50 in 1 gallon water, unless otherwise directed. Mix proper amount SEVIN in a little water to make a paste; then add balance of water to make a gallon of spray mixture. Make first application when insects or their damage first appear. Repeat applications at 7 to 10 day intervals to maintain effective control, unless otherwise directed. Keep spray mixture well agitated.

ORNAMENTALS, ROSES, OTHER FLOWERS, SHRUBS, SHADE TREES — To control June Bugs, apple aphid, bagworm, blister beetles, boxwood leaf miner, elm leaf aphid and beetle, flea beetles, gypsy moth, lace bugs, leaf hoppers, leaf rollers, mealy bugs, webworms, oak leaf miners, periodical cicada (17 year locust), plant bugs, puss caterpillars, psyllids, rose aphid and slug scale, insects, tent caterpillars, exposed thrips, and Japanese Beetles. **DO NOT USE ON BOSTON IVY.** (For larger quantities use 2 lbs. Sevin-50W per 100 gallons water)

LAWNS: To control Chinch bugs, ants, earwigs, fall armyworm, fleas, leafhoppers, millipedes, mosquitoes, sod webworms (lawn moths). Use 12 level tablespoons SEVIN-50 in 15 to 20 gallons water and apply evenly to 500 square feet of established lawn area.

VEGETABLES: Can be used on day of harvest unless otherwise indicated. **BEANS** — For Mexican bean beetle, bean leaf beetle, cutworms, cucumber beetles, flea beetles, Japanese beetles, leafhoppers, velvet bean caterpillar, armyworms, corn earworm, stinkbug, tarnished plant bug, lima bean pod borer, and Lygus. **CARROTS, LETTUCE** — 2 to 4 tablespoons. Do not use on lettuce within 14 days of harvest. For flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper (aster yellows vector), armyworms, cabbage caterpillars, corn earworm, sp. thrips, tarnished plant bugs, stink bugs. For best control of small cabbage loopers, corn earworm on lettuce, use every 5 to 7 days after heads form. **CORN** — Use 2 to 4 tablespoons. For Corn earworm, (apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals. Re-apply immediately after rain) corn rootworm adults, European corn borer (apply thoroughly into the shanks when plant is about knee high. Repeat at weekly intervals until control is achieved), fall armyworm, flea beetles, Japanese beetles, Sap beetles, Leafhoppers.

CUCUMBERS, SQUASH, PUMPKINS, MELONS — For pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers, squashbugs. **TOMATOES, POTATOES, EGGPLANT, PEPPERS** — Use 2 to 4 tablespoons. For Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA — For corn earworm and stink bugs. **CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI** — Use 2 to 4 tablespoons. Do not use within 3 days of harvest. For flea beetles, harlequin bug, armyworms, corn earworms. For control of small cabbage loopers use every 5 to 7 days.

FRUITS: Use up to 1 day before harvest unless otherwise directed. **APPLES, PEARS** — 1 tablespoon for Codling moth, apple mealybug, green apple aphid, white apple leafhopper, 2 tablespoons for apple maggot, bagworm, Eastern tent caterpillar, European apple sawfly, eye-spotted bud moth, red-banded and fruit tree leaf rollers, green fruitworm, Japanese beetle, pear psyllid, periodical cicada (17 year locust), plum curculio, rosy and woolly apple aphids, apple and pear rust mites, pear leaf blister mites, tarnished plant bug, tentiform leaf miners, Lecanium, cyster-shell and San Jose scales. **APRICOTS, PEACHES, NECTARINES** — For Cucumber beetles, European earwig, Japanese beetle, June beetles, lesser peach tree borer, peach twig borer, periodical cicada, red-banded and fruit tree leaf rollers, tarnished plant bug, tussock moth, Lecanium, San Jose and olive scales. Codling moth, oriental fruit moth, plum curculio: Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not use 3 days before harvest on apricots, nectarines. **CITRUS** — For citrus cut worm, fruit tree leaf roller, Western tussock moth, black scale, brown soft scale, citrus scale, yellow scale. Do not use within 5 days of harvest. **GRAPES** — Can be used on day of harvest. For European fruit Lecanium, grape leaf folder, grape leafhopper, grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles, red-banded leaf roller, **STRAWBERRIES** — Use up to 1 day before harvest. For cutworms, fruitworms, Japanese beetle, leafhoppers, spittlebug, strawberry leaf roller, strawberry weevil. **PLUMS** — Use up to 1 day before harvest. For mealy plum aphid, cherry fruit fly, eye-spotted bud moth, red-banded and fruit tree leaf roller, Japanese beetle, lesser peach twig borer, prune leafhopper, brown soft scale, Lecanium, cyster-shell and San Jose scales, peach twig borer. Eastern tent caterpillar, Codling moth, tussock moth. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

NUTS: Can be used on day of harvest. **PECANS** — For pecan nut case bearer. Do not use after husks split.

NOTICE: Because the Ferd Staffel Co. has no control over storage, handling, and conditions of use, which are of critical importance. Ferd Staffel Co. makes no representation of warranty, either express or implied, for results or residues greater than any tolerance established by applicable governmental agencies, nor due to misuse, improper handling or storage of this material. Nor does Ferd Staffel Co. assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

FERD STAFFEL COMPANY, San Antonio, TX 78298

Staffel's

sevin* 50
wettable

FS1-64A

FOR CONTROL OF INSECTS ON VEGETABLES, FRUITS, PECANS, ORNAMENTALS, FLOWERS, LAWNS & TREES

ACTIVE INGREDIENTS: Carbaryl (1-naphthyl-N-methycarbamate) ... 50%
INERT INGREDIENTS ... 50%
*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl-N-methycarbamate.

KEEP OUT OF REACH OF CHILDREN CAUTION

Read all directions before using
See back panel for additional cautions

U.S. Patent No. 2,903,473

EPA Reg. No. 3286-9 EPA Est. 3286-TX-1
Net Weight 12 oz.

3286-9

5-30-1978

11