

## FOR DAIRY AND BEEF CATTLE

### DIRECTIONS FOR APPLICATION BY HAND

FOR DIRECT APPLICATION TO DAIRY CATTLE AND BEEF CATTLE, FOR CONTROL OF HORNFLIES: Apply 2 heaping tablespoons (2 ounces) to the backs and flanks of infested animals. Rub lightly into hair. Repeat application no more than once in every 3 week period, if necessary.

Do not apply within 5 hours of milking or during milking.

FOR CONTROL OF LICE AND TICKS ON BEEF CATTLE. Use as above. Treat animals thoroughly. Repeat treatments as necessary.

AS AN AID FOR THE CONTROL OF FACE FLIES ON BEEF CATTLE: Dust the faces of cattle thoroughly. Frequent repeated applications may be necessary.

DO NOT TREAT ANIMALS LESS THAN ONE MONTH OF AGE.

FOR CONTROL OF HOG LICE: Apply to hogs, pens, and litter. One treatment may be sufficient. Treat animals thoroughly. Repeat in 10 days or thereafter as needed.

LICE AND KEDS ON SHEEP: Treat animals thoroughly, preferably following shearing. Repeat as necessary.

### DIRECTIONS FOR USE IN A DUST BAG

BEEF CATTLE: To aid in the control of horn flies and lice, hang where cattle

tend to congregate. For BEEF CATTLE a good location is near salt blocks, mineral blocks, or water supply. (Do not hang DUST BAG directly over feed, mineral, salt or water supply). For cattle on range or pasture, hang DUST BAG between fixed locations such as between 2 posts, buildings or trees. DUST BAG should be hung so bottom of bag is 4 to 6" below topline of cattle.

Allow one DUST BAG for each 10-15 animals.

Do not use the DUST BAG treatment for dairy cattle. For application to dairy cattle, follow directions given under "DIRECTIONS FOR APPLICATION BY HAND."

CAUTION: Harmful if swallowed. Do not apply within 5 hours of milking or during milking. Do not contaminate feed or water. Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.


Do not use on poultry or in poultry houses. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

NET WEIGHT 1 POUND

# DAIRY—BEEF CATTLE DUST

Contains METHOXYCHLOR and MALATHION



NET WEIGHT 1 POUND

# DAIRY—BEEF CATTLE DUST

Contains METHOXYCHLOR and MALATHION

For Dairy and Beef Cattle — Hogs, Sheep

**FOR HORNFY CONTROL ON DAIRY CATTLE  
AND HORNFLIES, LICE AND TICKS ON BEEF CATTLE**

**ACTIVE INGREDIENTS:**

Technical Methoxychlor*	5.0%
Malathion**	4.0%

INERT INGREDIENTS	91.0%
	100.0%

## CAUTION!

**KEEP OUT OF REACH OF CHILDREN!**

See side panel for additional caution

\*Equivalent to 4.4% 2,2-bis(p-methoxyphenyl)-1,1,1-trichlorethane and  
0.6% other isomers and related compounds.

\*\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**PREMIUM GRADE MALATHION ONLY**

**FERD STAFFEL CO.**

331 BURNET ST.

SAN ANTONIO, TEXAS 78298

EPA-Reg. NO. 3286-36